

The Efficacy of Patient With Untracentral Non-Small Cell Lung Cancer After Stereotactic Body Radiotherapy

Zhiyong Yuan ¹

1. Department of Radiation Oncology and Cyberknife Center, Tianjin Medical University Cancer Institute and Hospital, Tianjin, CHN

Corresponding author: Zhiyong Yuan, zhangyuhan2371@163.com

Categories: Radiation Oncology

Keywords: untracentral lung cancer, stereotactic body radiotherapy (sbrt), efficacy

How to cite this abstract

Yuan Z (April 02, 2020) The Efficacy of Patient With Untracentral Non-Small Cell Lung Cancer After Stereotactic Body Radiotherapy . Cureus 12(4): a542

Abstract

Objectives: To present the outcomes of patients with primary and untracentral non-small-cell lung cancer (NSCLC) after CyberKnife(CK) in 56Gy in our series.

Methods: We studied untracentral NSCLC patients, all with primary tumor, treated with SBRT between 2010 and 2018 retrospectively. The item, untracentral (UC)lung cancer defined as planning target volume (PTV) touching or overlapping tracheae, main bronchi, esophagus, heart or major vessels Within 2 cm around the bronchi. SBRT was delivered to total doses of 56 Gy in 6 fractions (n=2), 7 fractions (n=31), 8 fractions (n=25), 9 fractions (n=1) in consecutive days. We evaluate the results of clinical toxicity, overall survival (OS), progression-free survival (PFS) and local failure(LC).

Results: The gross of 59 patients were included in our study whose median age was 68 years ,74.6% of them accepted any therapy before. The median PTV was 77.8cc(ranger, 43.3, 91.8)and the median PTV volume was 59.3 cm³ (range,12.9-257.4). With a median follow -up of 57 months (range, 6-90 months), the median OSwas 32 months (range,2-105 months) and 5-year survival was 42.1%. On our univariate analysis, os was associated with tumor volume (=53.1 vs <53.1 cm³, p=0.016) and PTV dose (<85.9 vs=85.9cGy, p=0.039). On multivariate analysis, tumor volume which smaller than 53.1 cm³ has better OS (p=0.035). The median time of LC and PFS were 14 months and 20 months respectively. On the aspect of radiation toxicity, there were two cases (3.39%) with grade=3 COPD and one was intolerable.

Conclusions: SBRT using 56Gy delivered 7 or 8 fractions is available. On the other hand, a smaller tumor volume was associated with better OS.

Open Access

Abstract

Published 04/02/2020

Copyright

© Copyright 2020

Yuan. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Distributed under

Creative Commons CC-BY 4.0