

Evaluation of Target Motion During Stereotactic Body Radiotherapy of Ventricular Tachycardia

Lukas Knybel ¹, Jakub Cvek ², Radek Neuwirth ³, Otakar Jiravsky ³, Marek Sramko ⁴, Josef Kautzner ⁵

¹. Department of Oncology, University Hospital Ostrava, CZECH REPUBLIC ². Department of Oncology, University Hospital Ostrava ³. Internal Medicine, Cardiology, Podlesí Trinec, Třinec, CZE ⁴. Cardiology, Institute for Clinical and Experimental Medicine, Prague, CZE ⁵. Cardiac/thoracic/vascular Surgery, Institute for Clinical and Experimental Medicine, Prague, CZE

Corresponding author: Lukas Knybel, lukas.knybel@fno.cz

Categories: Radiation Oncology, Other

Keywords: ventricular tachycardia, cyberknife, motion management

How to cite this abstract

Knybel L, Cvek J, Neuwirth R, et al. (April 02, 2020) Evaluation of Target Motion During Stereotactic Body Radiotherapy of Ventricular Tachycardia. *Cureus* 12(4): a544

Abstract

Objectives: To evaluate the motion of a target in the heart using implantable cardioverter defibrillator (ICD) lead as surrogate.

Methods: We analyzed the log files of surrogate motion of 4 patients treated with SBRT of ventricular tachycardia. All clinical target volumes (CTV) were in direct contact with ICD lead. Targets were localized anterolateral (3) and inferolateral (1) in the heart. We evaluated the ICD lead motion amplitudes; intrafraction amplitude variability; correlation error between the ICD lead and external markers.

Results: In the superior-inferior (SI), latero-lateral (LL), and anterior-posterior (AP) directions, respectively, the median motion amplitudes were 5.5 mm (range 3.9 - 6.3 mm), 2.9 mm (range 2.1 - 3.2 mm), and 3.0 mm (range 2.1 - 4.9 mm). The median intrafraction amplitude variability was 2.2 mm (range 2.0 - 4.6 mm), 1.6 mm (range 0.4 - 2.4 mm), and 1.7 mm (range 1.6 - 4.4 mm) in the SI, LL, and AP directions, respectively. The median correlation error was 1.9 mm (range 1.8 - 2.6 mm).

Conclusions: Tracking of ICD lead as surrogate is feasible method to compensate breathing motion during SBRT of VT. Data from online tracking indicate motion irregularities and correlation errors consistently > 1mm what indicates the influence of the heartbeat.

Open Access

Abstract

Published 04/02/2020

Copyright

© Copyright 2020

Knybel et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Distributed under

Creative Commons CC-BY 4.0