

# Clinical outcomes of Stereotactic Body Radiation Therapy (SBRT) for the treatment of early-stage Non-Small Cell Lung Cancer (NSCLC) in the Community Hospital Setting

Luqman Dad <sup>1</sup>, Ryan Bathras <sup>2</sup>, Cody Tidwell <sup>2</sup>, Angel Torano <sup>3</sup>, Mary Young <sup>4</sup>

1. Radiation Oncology, DeCesaris Cancer Institute, Anne Arundel Medical Center, Annapolis, USA 2. Radiation Oncology, DeCesaris Cancer Institute, Anne Arundel Medical Center 3. Anne Arundel Medical Center, DeCesaris Cancer Institute, Anne Arundel Medical Center 4. DeCesaris Cancer Institute, Anne Arundel Medical Center

✉ **Corresponding author:** Luqman Dad, luq.dad@gmail.com

**Categories:** Radiation Oncology

**Keywords:** stereotactic body radiotherapy, non-small-cell lung cancer

## How to cite this abstract

Dad L, Bathras R, Tidwell C, et al. (June 16, 2016) Clinical outcomes of Stereotactic Body Radiation Therapy (SBRT) for the treatment of early-stage Non-Small Cell Lung Cancer (NSCLC) in the Community Hospital Setting. Cureus 8(6): a74

## Abstract

**Objectives:** SBRT is an effective treatment for medically inoperable patients with early-stage NSCLC. We evaluated outcomes with regards to local control, overall survival, and toxicity of SBRT for a medically inoperable patient population treated at Anne Arundel Medical Center. We determined clinical efficacy of definitive SBRT in medically inoperable early-stage NSCLC and compared outcomes to published data.

**Methods:** Between 2012 and 2015, 44 patients received a first course of SBRT for early-stage NSCLC. Five of the 44 patients received multiple course of SBRT, though the second, third or fourth courses were not included in this study due to a disproportionately high risk of failure and toxicity. All 44 tumors were treated on the Novalis Tx with cone-beam computed tomography (CT) localization. Simulation was conducted with 4D evaluation of the target and treatment was delivered to a static volume. There were two dosing schemes utilized: 54 Gy in 3 treatments and 50 Gy in 5 treatments, dependent on location being peripheral or central, as per RTOG guidelines.

**Results:** At a median follow-up of 19 months, the overall survival is 86%. Of the 44 patients, 36 (82%) had surveillance imaging done at least once, which found 31 (86%) patients were controlled locally, 2 (6%) had local progression, 2 (6%) had regional nodal, and 1 (3%) had distant failure to the brain. Radiation Therapy Oncology Group (RTOG)/Common Toxicity Criteria were reported in 43 patients, one was lost to follow-up. Of the 43 patients seen post-treatment, 3 experienced toxicity: 1 (2%) Grade 2 chest wall pain, 1 (2%) Grade 3 chest wall pain, and 1 (2%) Grade 3 pneumonitis.

**Conclusions:** Though we are limited by overall short follow-up time, SBRT provides good local control and limited toxicity, and therefore is an excellent treatment option for patients with medically inoperable NSCLC, our results are comparable to those reported in recently concluded clinical trials, RTOG 0236 and RTOG 0813.

## Open Access

### Abstract

Published 06/16/2016

### Copyright

© Copyright 2016

Dad et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Distributed under

Creative Commons CC-BY 3.0

