

## Open Access

## Abstract

Published 04/02/2023

## Copyright

© Copyright 2023

Trejo et al. This is an open access abstract distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Distributed under

Creative Commons CC-BY 4.0

## A Cumulative-Dose Optimization Approach of Volumetric Modulated Arc Therapy for Volume-Staged Stereotactic Radiosurgery to Treat Large Cerebral Arteriovenous Malformation

Orlando Trejo<sup>1</sup>, Fwang1 Wang<sup>2</sup>, Dima Soultan<sup>3</sup>

1. Radiation Oncology Physics, Kansas University Medical Center, Olathe, USA 2. Radiation Oncology, University of Kansas School of Medicine - Wichita, Wichita, USA 3. Radiation Oncology, University of Kansas School of Medicine, Kansas city, USA

**Corresponding author:** Orlando Trejo, [otrejo@kumc.edu](mailto:otrejo@kumc.edu)**Categories:** Medical Physics, Radiation Oncology**Keywords:** radiotherapy treatment planning, volumetric modulated arc therapy (vmat), arteriovenous malformation**How to cite this abstract**

Trejo O, Wang F, Soultan D (April 02, 2023) A Cumulative-Dose Optimization Approach of Volumetric Modulated Arc Therapy for Volume-Staged Stereotactic Radiosurgery to Treat Large Cerebral Arteriovenous Malformation. Cureus 15(4): a928

### Abstract

**Objectives:**

Large-volume, arteriovenous malformation (AVM) has conventionally been treated using gamma knife surgery (GKS) due to its high conformity and low dose-spillage values. Here we describe a cumulative-dose, volumetric modulated arc therapy (VMAT) optimization technique for such treatments.

**Methods:**

10 cases with large AVM (>10cc) were planned on a linear accelerator equipped with high-definition multi-leaf collimators and an external image-guided radiotherapy verification system. The AVMs were contoured and divided into 2-3 sub-targets to be treated sequentially in a volume-staged approach at two-week intervals. The prescription dose ranged from 18 Gy to 20 Gy for each stage, depending on the sub-target number, volume, and location. The first stage was optimized to minimize the dose to the adjacent sub-target. For every subsequent stage, the dose distribution from the previous stage was used as a base dose in the optimization process. Cumulative plans were evaluated in terms of target coverage (V100%), conformity index (CI100%), gradient index (GI), homogeneity index (HI), and normal brain V12Gy(%)

**Results:**

The VMAT plans resulted in an average planning target volume coverage of 96.7% ±1.2. The resulting CI100%, GI, HI, and V12Gy values were 1.19±0.2, 3.25±0.4, 1.27±0.1 and 2.52%±1.13, respectively.

**Conclusion(s):**

The linac-based plan optimization described can deliver multi-staged conformal doses to treat large AVMs. The results indicate similar conformity, gradient, and normal brain V12Gy to well-established GKS treatments.