

Review began 03/17/2023
Review ended 03/24/2023
Published 03/28/2023

© Copyright 2023
Qadeer et al. This is an open access article
distributed under the terms of the Creative
Commons Attribution License CC-BY 4.0.,
which permits unrestricted use, distribution,
and reproduction in any medium, provided
the original author and source are credited.

The Journey From Knee to Heart: A Case of Methicillin-Sensitive *Staphylococcus aureus* Infective Endocarditis Secondary to Septic Prepatellar Bursitis

Hafiza Qadeer¹, Amar Suwal¹, Ibiyemi O. Oke¹

1. Internal Medicine, Reading Hospital - Tower Health, Reading, USA

Corresponding author: Hafiza Qadeer, hafizaqadeer91@gmail.com

Abstract

Staphylococcus aureus is a cause of life-threatening gram-positive bacteremia and the most common causative organism of septic bursitis. Although there are several case reports of infective endocarditis complicated with septic bursitis and other rheumatic complications, there are very few cases reports of septic bursitis leading to infective endocarditis (IE). We present a patient with a history of recurrent prepatellar bursitis requiring catheter drainage, who presented with sepsis and IE.

Categories: Internal Medicine, Infectious Disease, Rheumatology

Keywords: mssa bacteremia, staphylococcus aureus, infective endocarditis, sepsis, pre-patellar bursitis

Introduction

Staphylococcus aureus is a common cause of life-threatening gram-positive bacteremia and is associated with a substantial risk of infective endocarditis (IE), especially in patients with underlying predisposing factors. A transesophageal echocardiogram, in some cases, is important in the evaluation of patients with gram-positive bacteremia for prognostication and therapeutic reasons [1]. The prepatellar bursa is located superficially in the knee, where it is prone to trauma and subsequent infection due to thin overlying skin. The most common causative organism of septic bursitis is *S. aureus* in 80% of cases; less commonly reported organisms include coagulase-negative *Staphylococci*, *Enterococcus spp*, *Escherichia coli*, *Pseudomonas aeruginosa*, and anaerobes [2].

Case Presentation

A 67-year-old with a past medical history of hypertension, moderate aortic stenosis, type 2 diabetes mellitus, hyperlipidemia, iron deficiency anemia, and a 10-year history of recurrent prepatellar bursitis with hemorrhagic effusion requiring catheter drainage (last drainage was done one week prior to this presentation), who worked as a bricklayer for more than 10 years, presented to the ER with two weeks history of worsening right knee swelling, pain, and difficulty with ambulation. There was associated redness of the skin overlying the swelling, loss of appetite, and chills but no fever, nausea, vomiting, night sweats, or malaise (Figure 1). No swelling was observed in other joints in the body.

How to cite this article

Qadeer H, Suwal A, Oke I O (March 28, 2023) The Journey From Knee to Heart: A Case of Methicillin-Sensitive *Staphylococcus aureus* Infective Endocarditis Secondary to Septic Prepatellar Bursitis. Cureus 15(3): e36806. DOI 10.7759/cureus.36806



FIGURE 1: An image of the patient's right knee at presentation showing swelling and overlying erythema

In the ER, the patient had a temperature of 36.7 °C, heart rate of 63 bpm, respiratory rate of 16/min, and oxygen saturation (SPO₂) of 99% on room air. There was marked swelling of the right knee with overlying erythema, differential warmth, and mild tenderness but no significant limitation of passive or active range of motion across the knee joint. He had a known 3/6 systolic ejection murmur in the aortic area but no physical stigmata of IE. His laboratory studies revealed a white cell count of $12.4 \times 10^3/\mu\text{L}$, hemoglobin 9.9g/dL, platelet 311, uric acid 7.0 (4.4mg/dL-7.6mg/dL), sedimentation rate 100 (0mm/hr-20mm/hr), C-reactive protein (CRP) 15.67 (<1.00mg/dL), creatinine 1.58, and an otherwise unremarkable basic metabolic panel. The X-ray of the right knee showed multicompartamental loss of joint space with osteophytosis and significant prepatellar soft tissue swelling (Figure 2).



FIGURE 2: Anteroposterior view of patient's right knee X-ray showing significant prepatellar swelling

The right knee joint aspirate was negative for crystals but grew methicillin-sensitive *S. aureus* (MSSA) and *Streptococcus mitis* group (Table 1).

Variable (unit)	Result	Normal value
Appearance	Bloody	Clear
Red blood cell (cmm)	11,000	0-2,000
White blood cell (cmm)	5,86,160	0-20
Segmented cells (%)	97	0 -50
Lymphocytes (%)	2	0-50
Monocytes (%)	1	0 -50
Gram stain	Few polymorphonuclear leukocytes and many gram-positive cocci	Negative
Culture and sensitivity	Methicillin-sensitive <i>Staphylococcus aureus</i> & <i>Streptococcus mitis</i> group	Negative
Fungal and anaerobic culture	Negative	Negative
Acid-fast Bacilli	Negative	Negative
Crystal analysis	Negative	Negative

TABLE 1: Synovial fluid analysis

One out of two blood culture bottles also grew MSSA, and the subsequent blood culture was negative. Transthoracic echocardiogram (TTE) was technically difficult due to his body habitus, but transesophageal echocardiogram (TEE) revealed a small mass measuring 0.8cm in its largest diameter on the aortic side of the right coronary cusp, concerning for valvular vegetations (Figure 3).




FIGURE 3: Transesophageal echocardiogram showing an aortic mass

The patient was taken to the OR for incision and drainage of right knee effusion with wound vacuum-assisted closure (VAC) placement. His large prepatellar infected hematoma was drained with residual lateral and medial pockets, and necrotic tissue was debrided. A culture of necrotic tissue grew MSSA and *S. mitis*. He was started on IV cefazolin which was switched to IV daptomycin at discharge by an infectious disease specialist to enable once-daily antibiotic administration for a total of six weeks. His follow-up TTE was done after the completion of antibiotics and revealed no obvious vegetation within the limits of the study. He completed six weeks of antibiotics and regained the full function of his knee with physical therapy.

Discussion

Septic prepatellar bursitis constitutes one-third of the total cases of prepatellar bursitis [3]. *Staphylococcus aureus* is the causative organism in 80% of the cases. In 55% of the cases, the predisposing factor is an overlying skin lesion related to occupational or recreational trauma. Immunosuppressive states like steroid use and diabetes mellitus also increase the risk of septic bursitis [4]. Typically, it presents as erythema, pain, swelling, and warmth of the prepatellar area. Careful examination is essential as the limitation of joint movement should raise suspicion for concurrent septic arthritis. Initial blood work shows leukocytosis with neutrophilia and elevation of sedimentation rate and CRP.

In diagnosing septic bursitis, bursa fluid aspirate should be sent for cell count, cytology, crystal analysis, gram stain, and culture. Bursa fluid count greater than 2000 cells/microliter and positive gram stain is consistent with septic bursitis. The absence of crystals rules out crystal-associated bursitis. In terms of imaging, plain radiographs should show soft tissue swelling with bursa fluid accumulation and synovial thickening. Two sets of blood cultures should be sent before initiating antibiotics. One-third of the cases of bursitis are associated with bacteremia, with *S. aureus* being the most common pathogen [5].

Our report features a case of a bricklayer with a history of type 2 diabetes mellitus and recurrent prepatellar bursitis, with prior catheter drainage of hemorrhagic bursa effusion. Bursa fluid aspiration revealed markedly elevated leukocyte count and grew MSSA and *S. mitis*, confirming septic bursitis. One out of two of the first sets of blood cultures grew *S. aureus*. Repeat blood cultures were negative.

Transthoracic echocardiography is recommended in Staphylococcal bacteremia to rule out IE. The Infectious Disease Society of America guidelines for methicillin-resistant *S. aureus* (MRSA) bacteremia recommends echocardiography for all patients with *S. aureus* bacteremia. The TEE is preferred as it is a more sensitive modality. It is important as the presence of concomitants impacts the decision on the duration of antimicrobial therapy [6]. This patient's TTE revealed moderate aortic stenosis due to calcification but no valvular vegetations, however, a negative TTE is insufficient for ruling out IE because of its low sensitivity compared to TEE. We proceeded with performing TEE as the patient's preexisting degenerative aortic valve disease increased his risk of IE [7]. The TEE showed a calcified tri-leaflet aortic valve with a small mass attached to the aortic side of the right coronary cusp. These findings were consistent with endocarditis. Based on our literature review, there have been some case reports of IE complicated with septic bursitis and other rheumatic complications [8-10]. However, there are very few case reports on septic bursitis leading to IE [11]. The temporal event in this patient's case suggests that his septic bursitis was complicated by endocarditis and not vice versa. His history of recurrent prepatellar bursitis related to overuse injury and recent catheter drainage of the hemorrhagic effusion predisposed him to septic bursitis and sepsis, while his underlying degenerative aortic valve disease was his risk factor for IE. The TTE is not indicated for gram-negative bacteremia unless risk factors like a prosthetic heart valve, injection drug abuse, and previous history of IE are present.

Conclusions

Infective endocarditis is rarely secondary to septic prepatellar bursitis, but all patients with gram-positive bacteremia should have at least a TTE to rule out IE and be on the appropriate anti-staphylococcal antibiotics. Patients with underlying risk factors for IE are more predisposed, and appropriate patients with gram-negative bacteremia should also be considered for a work-up of IE. This case highlights the role of TEE in detecting IE in patients with an otherwise normal TTE.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Salvador VB, Chapagain B, Joshi A, Brennessel DJ: Clinical risk factors for infective endocarditis in staphylococcus aureus bacteremia. *Tex Heart Inst J*. 2017, 44:10-15. [10.14503/THIJ-15-5359](#)
- Gendernalik JD, Sechriest VF 2nd: Prepatellar septic bursitis: a case report of skin necrosis associated with open bursectomy. *Mil Med*. 2009, 174:666-669. [10.7205/milmed-d-03-7608](#)
- Baumbach SF, Lobo CM, Badyine I, Mutschler W, Kanz KG: Prepatellar and olecranon bursitis: literature review and development of a treatment algorithm. *Arch Orthop Trauma Surg*. 2014, 134:359-370. [10.1007/s00402-013-1882-7](#)
- Gómez-Rodríguez N, Méndez-García MJ, Ferreiro-Seoane JL, Ibáñez-Ruán J, Penelas-Cortés Bellas Y: Infectious bursitis: study of 40 cases in the pre-patellar and olecranon regions. *Enferm Infecc Microbiol*

- Clin. 1997, 15:237-242. [10.1007/springerreference_38731](https://doi.org/10.1007/springerreference_38731)
5. González-Gay MA, García-Porrúa C: Systemic vasculitis in adults in northwestern Spain, 1988-1997. Clinical and epidemiologic aspects. *Medicine (Baltimore)*. 1999, 78:292-308. [10.1097/00005792-199909000-00002](https://doi.org/10.1097/00005792-199909000-00002)
 6. Rasmussen RV, Høst U, Arpi M, et al.: Prevalence of infective endocarditis in patients with *Staphylococcus aureus* bacteraemia: the value of screening with echocardiography. *Eur J Echocardiogr*. 2011, 12:414-420. [10.1093/ejechocard/jer023](https://doi.org/10.1093/ejechocard/jer023)
 7. Michel PL, Acar J: Native cardiac disease predisposing to infective endocarditis. *Eur Heart J*. 1995, 16:2-6. [10.1093/eurheartj/16.suppl_b.2](https://doi.org/10.1093/eurheartj/16.suppl_b.2)
 8. Soor P, Sharma N, Rao C: Multifocal septic arthritis secondary to infective endocarditis: a rare case report. *J Orthop Case Rep*. 2017, 7:65-68. [10.13107/jocr.2250-0685.691](https://doi.org/10.13107/jocr.2250-0685.691)
 9. Tan WH, Chew LC: Acute infective endocarditis presenting with polyarthritis after turbinoplasty. *Singapore Med J*. 2023, 64:149-151. [10.11622/smedj.2021159](https://doi.org/10.11622/smedj.2021159)
 10. Lin HH, Peng MY, Wu LS, Chang FY: Bilateral olecranon purulent bursitis complicated prosthetic valve endocarditis caused by oxacillin susceptible *Staphylococcus aureus*: a case report. *J Intern Med Taiwan*. 2003, 14:196-200.
 11. Rufael DW, Cohn SE: Native valve endocarditis due to *Corynebacterium striatum*: case report and review. *Clin Infect Dis*. 1994, 19:1054-1061. [10.1093/clinids/19.6.1054](https://doi.org/10.1093/clinids/19.6.1054)