

Ureterovaginal Fistula Post Vaginal Hysterectomy

SaiNidhi G. Reddy¹, Pratap Parihar¹, Rajasbala Dhande¹, Roohi G. Gupta¹, Vadlamudi Nagendra¹

Received 10/10/2022

Review began 10/15/2022

Review ended 10/20/2022

Published 10/26/2022

© Copyright 2022

Reddy et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Radiodiagnosis, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Wardha, IND

Corresponding author: SaiNidhi G. Reddy, nidhireddy51@gmail.com

Abstract

Middle-aged women with ureterovaginal fistula (UVF) after hysterectomy represent a painful condition for the patients in the community. Accurate diagnosis and proper planning before surgery are essential for effective outcomes. CT urography is the modality of choice in diagnosing ureterovaginal fistula. CT urography helps in evaluating the fistula as well the associated renal complications following the condition. Here we present a case of ureterovaginal fistula reported with a history of vaginal hysterectomy for subserosal fibroid in December 2021.

Categories: Radiology, Urology

Keywords: excretory phase, hydronephrosis, fibroid, fistula, ct urography

Introduction

Since a urogenital fistula forms after gynecological conditions or obstetric surgery, treating urinary incontinence is a difficult undertaking. It is a terrible condition, and clinical problems are made worse by social and psychological elements. Vesicovaginal fistula is the most common urogenital fistula. The ureteral fistula often joins the vagina but sporadically may also connect to the uterus or fallopian tube [1]. Endometriosis, obesity, pelvic inflammatory disease, radiation therapy, and malignant illness of the pelvis are some conditions that increase the risk of developing ureteral fistulas [2]. The symptoms of ureteric damage can manifest either suddenly or more gradually, depending on the type of injury, and they are particularly difficult to detect during laparoscopic procedures. The incidence of ureterovaginal fistula (UVF) has been increasing due to the growing use of the laparoscopic surgical technique [3].

Case Presentation

A 42-year-old female presented to the department of obstetrics and gynecology with complaints of leaking urine from the vagina for three weeks. On per-speculum examination, there was evidence of dripping urine from the vagina. The patient has a history of large subserosal fibroid for which she was advised for a total hysterectomy. A large subserosal fibroid was identified arising from the posterior wall of the uterus, and a vaginal hysterectomy was performed, leaving the bilateral adnexa and vagina in situ. The procedure was uneventful, and the patient was discharged in three days. The patient developed urinary incontinence and left-sided flank pain two months after the surgery. The patient was then advised for CT urography for further evaluation. On the excretory phase (3 mins), axial and coronal images are showing normal excretion of contrast in the right renal collecting system filling the bladder but no excretion of contrast in the left renal collecting system. The cavity in the vaginal stump appears distended with fluid-filled collection within, possibly urine (Figure 1). Excretory phase (15 mins) axial, coronal, and sagittal images show dilated left pelvicalyceal system and ureter with contrast excretion. The cavity of the vaginal stump shows contrast within, suggesting an abnormal connection of the left ureter with the vaginal stump (Figure 2). We can also appreciate a contrast-filled tract connecting the left ureter to the vaginal stump, confirming ureterovaginal fistula in the axial and coronal excretory phase (15 mins) (Figure 3).

How to cite this article

Reddy S G, Parihar P, Dhande R, et al. (October 26, 2022) Ureterovaginal Fistula Post Vaginal Hysterectomy. Cureus 14(10): e30694. DOI 10.7759/cureus.30694




FIGURE 1: Axial (A) and coronal (B) CT urography excretory phase (3 mins), showing normal excretion of contrast in the right renal collecting system (black arrow) filling the bladder (red arrow), no excretion of contrast in the left renal collecting system (white arrow). The cavity in the vaginal stump appears distended with fluid-filled collection within, possibly urine (blue arrow).

CT: Computed tomography




FIGURE 2: Coronal (A) and sagittal (B) images of the excretory phase (15 mins) shows dilated left pelvicalyceal system (white arrow) and ureter (black arrow) with contrast excretion into the bladder (red arrow). The cavity of the vaginal stump shows e/o excretion of contrast (blue arrow), within suggesting an abnormal connection of the left ureter with the vaginal stump.




FIGURE 3: Axial (A) and coronal (B) excretory phase (15 mins) images showing a contrast-filled tract (red arrow) connecting the left ureter to the vaginal stump, confirming ureterovaginal fistula.

Discussion

Fistulous tracts between the vagina and ureter are rare, occurring primarily as complications of obstetric and gynaecologic procedures such as low transverse cesarean section or uterine curettage for elective abortion [1]. Pelvic adhesions due to repeated surgery significantly enlarged uterus during abdominal surgery and heavy bleeding that obscures the surgical site during surgery were the main causes of ureteral injury [4]. Clamp, suture ligation, partial or total primary transection, angulation, avulsion, or ischemic necrosis following electrocoagulation are examples of mechanisms that can cause ureter damage during surgical procedures [5]. Endometriosis, obesity, pelvic inflammatory disease, radiation therapy, and pelvic cancer are risk factors for the ureterovaginal fistula [2].

Ureterovaginal fistulas can also be a side effect of pelvic procedures, most commonly hysterectomy. In a 15-year review of genitourinary fistulas at the Mayo Clinic, ureterovaginal fistulas accounted for 31 of 303 cases, with the vast majority resulting from hysterectomy [6]. Due to its potential to result in incontinence, infection, and renal failure, the ureterovaginal fistula is the most dangerous of the urovaginal fistulas [7].

Despite the regular act of micturition, the primary ureterovaginal fistula presentation is an incontinence disease. Incontinence usually develops 1 to 4 weeks after surgery [8]. Affected women will present with urinary outflow from the vagina, and pelvic examination can demonstrate urine leakage at the cervical os [9].

Prevention of renal impairment requires early diagnosis. The diagnosis is made using various imaging techniques, which include excretory urography, intravenous CT urography, and magnetic resonance imaging (MRI). Vesicovaginal fistula frequently coexists. The diagnosis is confirmed by the evidence of contrast extravasation from the ureter into the vagina, which may be seen on a delayed phase CT scan. Patients with renal failure or conditions where CT is contraindicated can use MRI [10]. MRI can be performed using contrast-enhanced T1-weighted image sequences or non-contrast T2-weighted image sequences. In situations involving renal insufficiency, T2-weighted MR urography is employed [11].

The fistula may be able to heal with conservative therapy using a percutaneous nephrostomy tube and/or ureteral stent. Once the ureterovaginal fistula is diagnosed, an immediate ureteral stent or nephrostomy tube placement should be tried. Prompt urine drainage is necessary to maintain renal function even if surgery is subsequently required [12,13].

Conclusions

The morbidity linked to ureteral injury may be severe, leading to a protracted hospital stay, a poor surgical outcome, additional invasive procedures, decreased renal function, and a lower quality of life for the patient. Following a vaginal hysterectomy, the risk of ureteral complications is similar to that of an open laparotomy. The most effective ways to avoid harm are surgical skills, understanding of the anatomy of the pelvic wall, and meticulous identification and mobilization of the ureter. It's essential to get a diagnosis early to avoid needing long-term care. In the vast majority of patients with early diagnosis, endoscopic

procedures are adequate. Our case was managed with a percutaneous nephrostomy tube followed by antegrade stent placement. However, severe ureteral strictures, urogenital fistulas, and renal insufficiency caused by ureteral injuries necessitate more comprehensive reconstructive surgery.

Additional Information

Disclosures

Human subjects: All authors have confirmed that this study did not involve human participants or tissue.

Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Billemeyer BR, Nygaard IE, Kreder KJ: Ureterouterine and vesicoureterovaginal fistulas as a complication of cesarean section. *J Urol*. 2001, 165:1212-3. [10.1016/S0022-5347\(05\)66481-6](https://doi.org/10.1016/S0022-5347(05)66481-6)
2. Murtaza B, Mahmood A, Niaz WA: Ureterovaginal fistula-etiological factors and outcome. *J Pak Med Assoc*. 2012, 62:999-1003.
3. Parpala-Spärman T, Paananen I, Santala M, Ohtonen P, Hellström P: Increasing numbers of ureteric injuries after the introduction of laparoscopic surgery. *Scand J Urol Nephrol*. 2008, 42:422-7. [10.1080/00365590802025857](https://doi.org/10.1080/00365590802025857)
4. Meirou D, Moriel EZ, Zilberman M, Farkas A: Evaluation and treatment of iatrogenic ureteral injuries during obstetric and gynecologic operations for nonmalignant conditions. *J Am Coll Surg*. 1994, 178:144-8.
5. Ratajczak JM, Hladun T, Orchel G: Latrogenic ureterovaginal fistula after laparoscopic hysterectomy: a case report. *Prz Menopauzalny*. 2021, 20:48-51. [10.5114/pm.2021.104337](https://doi.org/10.5114/pm.2021.104337)
6. Lee RA, Symmonds RE, Williams TJ: Current status of genitourinary fistula. *Obstetrics and gynecology*. 1988, 1:313-9. [10.1016/0020-7292\(89\)90762-5](https://doi.org/10.1016/0020-7292(89)90762-5)
7. Labasky RF, Leach GE: Prevention and management of urovaginal fistulas. *Clin Obstet Gynecol*. 1990, 33:382-91.
8. Mandal AK, Sharma SK, Vaidyanathan S, Goswami AK: Ureterovaginal fistula: summary of 18 years' experience. *Br J Urol*. 1990, 65:453-6. [10.1111/j.1464-410x.1990.tb14785.x](https://doi.org/10.1111/j.1464-410x.1990.tb14785.x)
9. Sheen JH, Lin CT: A simple means of making the differential diagnosis of ureterouterine and vesicouterine fistula. *J Urol*. 1998, 1:1420-1.
10. Yu NC, Raman SS, Patel M, Barbaric Z: Fistulas of the genitourinary tract: a radiologic review. *Radiographics*. 2004, 24:1331-52. [10.1148/rg.245035219](https://doi.org/10.1148/rg.245035219)
11. Mamere AE, Coelho RDS: Evaluation of urogenital fistulas by magnetic resonance urography. *Radiol Bras* [online]. 2008, 41:19-23. [10.1590/S0100-39842008000100007](https://doi.org/10.1590/S0100-39842008000100007)
12. Dowling RA, Corriere JN Jr, Sandler CM: Latrogenic ureteral injury. *J Urol*. 1986, 135:912-5.
13. Gerber GS, Schoenberg HW: Female urinary tract fistulas. *J Urol*. 1993, 149:229-36.