

Desensitization of Brentuximab Vedotin in a Patient with Hodgkin Lymphoma

Sidra Khalid ¹, Aariez Khalid ², Bernadette A. Clark ³, Hamed Daw ⁴

1. Internal Medicine Residency, Fairview Hospital, Cleveland Clinic, USA 2. Biomedical Science, University of Guelph, Guelph, CAN 3. Oncology, Cleveland Clinic Fairview Hospital, Cleveland, USA 4. Department of Hematology and Oncology, Fairview Hospital, Cleveland Clinic, USA, Cleveland, USA

✉ **Corresponding author:** Sidra Khalid, drsidrakhalid@hotmail.com

Disclosures can be found in Additional Information at the end of the article

Abstract

Brentuximab vedotin is a monoclonal antibody that targets the CD30 antigen. It is indicated for the treatment of Hodgkin lymphoma. Hypersensitivity reactions have occurred during infusions of brentuximab vedotin, ranging from mild to severe. We report a case of a 46-year-old male with stage IV nodular sclerosis Hodgkin lymphoma who developed a hypersensitivity reaction to brentuximab vedotin. He experienced a generalized rash, facial swelling, and mild airway obstruction. In order to continue treatment with brentuximab vedotin, we implemented a desensitization protocol. He was premedicated and a 12-step process was performed in which brentuximab vedotin was titrated over three hours. The protocol was successful, allowing the patient to receive subsequent infusions without hypersensitivity reactions.

Categories: Internal Medicine, Allergy/Immunology, Oncology

Keywords: brentuximab vedotin, hypersensitivity, desensitization, hodgkin lymphoma

Introduction

Brentuximab vedotin is a monoclonal antibody against the CD30 antigen present on Reed-Sternberg cells [1]. Brentuximab vedotin is indicated for the treatment of Hodgkin lymphoma in previously untreated advanced disease, relapsed or refractory disease and for consolidation after autologous hematopoietic stem cell transplantation. One of the adverse effects of brentuximab vedotin is hypersensitivity reaction during infusion. In clinical trials using brentuximab vedotin as monotherapy, 13% of patients experienced an infusion reaction, out of which 1.2% of patients experienced a grade 3 infusion reaction. When brentuximab vedotin was studied in combination with chemotherapy, 9% of patients experienced an infusion reaction, out of which 0.4% experienced grade 3 reactions. The hypersensitivity reaction can range from a mild infusion reaction to a more severe reaction including anaphylaxis. Although the manufacturer's information on brentuximab vedotin recommends permanently discontinuing brentuximab vedotin if an anaphylactic reaction occurs, several case reports have shown success with a desensitization protocol [2]. In this case, we report a successful desensitization regimen with brentuximab vedotin, with no further hypersensitivity reactions noted during the subsequent infusions.

Case Presentation

A 46-year-old male with stage IVB nodular sclerosis Hodgkin lymphoma received treatment with six cycles of ABVD (doxorubicin, bleomycin, vinblastine, dacarbazine) and then autologous stem cell transplant. Despite these treatments, he had progression of his disease. The positron emission tomography (PET) showed hypermetabolic activity in the anterior

Received 06/14/2018

Review began 07/02/2018

Review ended 07/12/2018

Published 07/13/2018

© Copyright 2018

Khalid et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Khalid S, Khalid A, Clark B A, et al. (July 13, 2018) Desensitization of Brentuximab Vedotin in a Patient with Hodgkin Lymphoma. Cureus 10(7): e2981. DOI 10.7759/cureus.2981

mediastinal mass and left hilar lymph node. He underwent radiation to the mediastinum. A subsequent computed tomography (CT) scan of the chest and liver showed a progression of the anterior mediastinal mass measuring 3.7 x 2.2 cm and a new mass in the left lobe of the liver measuring 2.9 x 2.8 cm (Figures 1-2).



FIGURE 1: Computed tomography (CT) scan of the chest showing the anterior mediastinal mass



FIGURE 2: Computed tomography (CT) scan of the liver showing a mass in the left lobe

A liver biopsy of the mass showed recurrent classical Hodgkin lymphoma. He was started on brentuximab vedotin at a dose of 1.8 mg/kg after administration of oral diphenhydramine 25 mg and oral acetaminophen 650 mg. Three weeks later, the second cycle was administered at the same dose with the same premedications. After receiving 10 mL of the medication, he developed facial flushing, swelling, generalized rash, and a scratchy sensation in his throat. He was given hydrocortisone 100 mg intravenously and his symptoms improved. Prior to the next cycle of treatment, the diphenhydramine 25 mg was changed to the intravenous (iv) route and dexamethasone sodium phosphate 20 mg iv and famotidine 20mg iv were added. During his third brentuximab vedotin infusion, he developed facial flushing and swelling, scratchy throat, and a rash. The infusion was stopped and he was given diphenhydramine 25 mg iv. His symptoms resolved gradually. It was decided to admit him to the hospital for cycle four for brentuximab vedotin desensitization. He was premedicated with methylprednisolone 60 mg iv, acetaminophen 650 mg orally, famotidine 20 mg iv, diphenhydramine 50 mg iv, and montelukast 10 mg orally. The following protocol was used for brentuximab vedotin at 1.8 mg/kg (Table [1](#)).

Brentuximab 1.8 mg/kg = 149 mg total dose		
Elapsed Timed	Administration Rate	Ratio of total dose
Bag 1: 100 fold dilution made as 1.49 mg in 250 mL. Total dose infused from Bag 1 = 9.25 mL x (1.49 mg/250 mL) = 0.05513 mg		
0 minutes	2 mL/hr	1:50,000
15 minutes	5 mL/hr	1:20,000
30 minutes	10 mL/hr	1:10,000
45 minutes	20 mL/hr	1:5,000
Bag 2: 10 fold dilution made as 14.9 mg in 250 mL. Total dose infused from Bag 1 = 18.75 mL x (14.9 mg/250 mL) = 1.1175 mg		
75 minutes	5 mL/hr	1:2,000
90 minutes	10 mL/hr	1:1,000
105 minutes	20 mL/hr	1:500
120 minutes	40 mL/hr	1:250
Bag 3: Remainder of dose made as 147.8 mg in 250 mL		
135 minutes	10 mL/hr	1:100
150 minutes	20 mL/hr	1:50
165 minutes	40 mL/hr	1:25
180 minutes	80 mL/hr	Infuse remainder of dose (137mg) at 80mL/hr

TABLE 1: Twelve-step desensitization protocol for brentuximab vedotin

The patient had no complications during the desensitization process. For cycle five, he was admitted as well, and similarly, had no hypersensitivity reactions. The dose of brentuximab vedotin was reduced to 1.2 mg/kg on cycle six for worsening neuropathy, but the desensitization was continued. After cycle six, the treatment with brentuximab vedotin was discontinued due to worsening neuropathy and progression of the cancer. The PET scan showed several new hypermetabolic foci in the liver with mildly hypermetabolic lung nodules slightly progressed from prior imaging studies. He was started on nivolumab for further management of his Hodgkin lymphoma.

Discussion

Based on the American Cancer Society, in 2018, there are about 8500 new cases of Hodgkin lymphoma. There are about 1050 deaths that occur due to this disease. It is commonly diagnosed in the third decade of life [3]. According to the World Health Organization (WHO), Hodgkin lymphoma is classified as nodular lymphocyte predominant or classical. The classical form is further subdivided into nodular sclerosis, lymphocyte-rich, mixed cellularity,

lymphocyte-depleted [4]. Nodular sclerosis classical Hodgkin lymphoma is the most common subtype accounting for about seven out of every ten cases. The Ann Arbor classification system is used for staging Hodgkin lymphoma. Stage I – involvement of a single lymphatic site to localized involvement of a single extralymphatic organ (except lymph nodes); stage II – involvement of two or more lymph node regions on the same side of the diaphragm, or localized involvement of a single extralymphatic organ site with associated regional lymph nodes; stage III – involvement of lymph node regions on both sides of the diaphragm, with extension to other adjacent lymph nodes and/or spleen; stage IV – diffuse involvement of one or more extralymphatic organs with or without lymph node involvement, or involvement of liver, bone marrow, cerebrospinal fluid, or isolated extralymphatic organ with or without lymph node involvement but involving a distant site. These stages can be further sub classified into A (no symptoms) and B (symptoms of unexplained weight loss >10% in six months, fever above 38°C, night sweats). Clinical staging is performed by CT thoracic, abdomen and pelvis with or without PET scan [1].

Treatment is based on the stage of the disease. For stages IA or IIA, chemotherapy, combined chemotherapy, and radiation therapy, or radiation therapy alone are options. Treatment options could include ABVD for four to six cycles; ABVD for two cycles plus radiation therapy (20 Gy or 30 Gy); or radiation therapy alone. For stage III or IV disease, bulky disease (>10 cm mass or mediastinal disease involving >33% of the transthoracic diameter) or B symptoms; combination chemotherapy with or without radiation therapy is recommended. ABVD for six to eight cycles is the standard of care [1].

In relapsed Hodgkin lymphoma, patients can undergo re-induction with the same chemotherapy or another chemotherapy regimen followed by autologous hematopoietic stem cell transplant. Subsequently after stem cell transplant, if there is a relapse, an involved-field radiation therapy for residual masses could be performed [1]. Another option is brentuximab vedotin, which is a monoclonal antibody which targets the CD30 antigen on the Reed-Sternberg cells [5]. It is FDA (US Food and Drug Administration) approved for patients who have relapsed after autologous stem cell transplantation or have failed at least two lines of therapy. On March 20, 2018, the FDA approved brentuximab vedotin to treat adult patients with previously untreated stage II or IV classical Hodgkin lymphoma in combination with chemotherapy based on results from the ECHELON-1 trial [6]. As brentuximab vedotin is an important treatment option for patients with Hodgkin lymphoma, it is crucial to know about its hypersensitivity reaction and its plausible method of desensitization. Our case describes a hypersensitivity reaction to brentuximab vedotin, which manifested as a rash and mild airway obstruction. Based on case reports about brentuximab vedotin desensitization, we implemented the protocol in our patient (Table 2).

Case Reports	Hypersensitivity Reaction	Premedication Agents Used	Desensitization Process
Arora et al. [7]	<ul style="list-style-type: none">Anaphylaxis	<ul style="list-style-type: none">Methylprednisolone 125 mgDiphenhydramine 50 mgRanitidine 50 mgMontelukast 10 mg	<ul style="list-style-type: none">Three brentuximab vedotin (1.8 mg/kg) dilutions of increasing concentrations administered at increasing rates over a 12-step process

DeVita et al. [8]	<ul style="list-style-type: none">Anaphylaxis	<ul style="list-style-type: none">Methylprednisolone 60 mgDiphenhydramine 50 mgFamotidine 20 mgMontelukast 10 mgAcetaminophen	<ul style="list-style-type: none">Three brentuximab vedotin (1.8 mg/kg) dilutions of increasing concentrations were delivered at four increasing rates, over 13 consecutive steps
Fizesan et al. [9]	<ul style="list-style-type: none">Anaphylaxis	<ul style="list-style-type: none">Aspirin 500 mgCetirizine 10 mgMontelukast 10 mgRanitidine 150 mg	<ul style="list-style-type: none">A 17-step protocol in which four brentuximab vedotin (1.8 mg/kg) dilutions were given at increasing concentrationsThis resulted in a hypersensitivity reaction, methylprednisolone 120 mg and dexachlorpheniramine 5 mg were given and added to future regimens (next protocol with 1 mg/kg, then 1.5 mg/kg, followed by 1.8 mg/kg)
O'Connell et al. [10]	<ul style="list-style-type: none">Anaphylaxis	<ul style="list-style-type: none">Four days of premedication with dexamethasone 4 mg/dayDiphenhydramine 50 mgFamotidine 20 mgCetirizine 10 mgAcetaminophen 1 g	<ul style="list-style-type: none">Three brentuximab vedotin (1.8 mg/kg) dilutions of increasing concentrations administered at increasing rates over a 12-step process
Story et al. [11]	<ul style="list-style-type: none">Hives	<ul style="list-style-type: none">Methylprednisolone 60 mgFexofenadine 180 mg	<ul style="list-style-type: none">A 13-step protocol with 150 mg of brentuximab vedotin (1.8 mg/kg) administered at increasing concentrations

TABLE 2: Case reports of desensitization protocols for brentuximab vedotin

We implemented the desensitization protocol and premedicated our patient with 50 mg iv diphenhydramine, 20 mg iv famotidine, 60 mg iv methylprednisolone, 10 mg oral montelukast, and acetaminophen. We then followed the 12-step desensitization protocol, in which the concentration of the drug was gradually increased. Our patient underwent successful brentuximab vedotin desensitization and was able to undergo further treatment. Routinely,

when there is disease progression with brentuximab vedotin then the next step would be to start treatment with anti-program cell death (PD)1 inhibitor, such as nivolumab.

Conclusions

Through our case, we have highlighted that brentuximab vedotin is an important treatment option for patients with advanced Hodgkin lymphoma. Hypersensitivity reactions can occur with this drug, which can be successfully managed with desensitization.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Adult hodgkin lymphoma treatment. (2018). Accessed: May 2018: <https://www.ncbi.nlm.nih.gov/books/NBK66038/>.
2. Highlights of prescribing information - Adcetris . (2018). Accessed: May 2018: http://www.seattlegenetics.com/application/files/1915/2157/0234/adcetris_USPI.pdf.
3. Key statistics for hodgkin lymphoma. (2018). Accessed: May 2018: <https://www.cancer.org/cancer/hodgkin-lymphoma/about/key-statistics.html>.
4. Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R: The 2016 revision of the world health organization classification of lymphoid neoplasms. *Blood*. 2016, 127:2375-2390. [10.1182/blood-2016-01-643569](https://doi.org/10.1182/blood-2016-01-643569)
5. Domingo-Domenech E, Comai A, Sureda A: Brentuximab vedotin in relapsed/refractory hodgkin's lymphoma. *Eur Oncol Haematol*. 2015, 11:21-24. [10.17925/EOH.2015.11.01.21](https://doi.org/10.17925/EOH.2015.11.01.21)
6. Brentuximab vedotin. (2018). Accessed: May 2018: <https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm601947.htm>.
7. Arora A, Chatt VR, Liewer S, Armitage JO, Bociek RG: Brentuximab vedotin desensitization in a patient with refractory hodgkin's lymphoma. *Eur J Haematol*. 2015, 95:361-364. [10.1111/ejh.12570](https://doi.org/10.1111/ejh.12570)
8. DeVita MD, Evens AM, Rosen ST, Greenberger PA, Petrich AM: Multiple successful desensitizations to brentuximab vedotin: a case report and literature review. *J Natl Compr Canc Netw*. 2014, 12:465-471. [10.6004/jnccn.2014.0052](https://doi.org/10.6004/jnccn.2014.0052)
9. Fizesan M, Boin C, Aujoulat O, Newinger G, Ghergus D, Watier H: Successful rapid desensitization to the antibody-drug conjugate brentuximab vedotin in a patient with refractory hodgkin lymphoma. *J Oncol Pharm Pract*. 2016, 22:188-192. [10.1177/1078155214567164](https://doi.org/10.1177/1078155214567164)
10. O'Connell AE, Lee JP, Yee C, Kesselheim J, Dioun A: Successful desensitization to brentuximab vedotin after anaphylaxis. *Clin Lymphoma Myeloma Leuk*. 2014, 14:73-75. [10.1016/j.clml.2013.11.003](https://doi.org/10.1016/j.clml.2013.11.003)
11. Story SK, Petrov AA, Geskin LJ: Successful desensitization to brentuximab vedotin after hypersensitivity reaction. *J Drugs Dermatol*. 2014, 13:749-751.