

Gangrenous Cystitis Secondary to a Spontaneous Hematoma of the Lesser Pelvis

Review began 08/28/2023

Review ended 09/12/2023

Published 09/18/2023

© Copyright 2023

Angelopoulos et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Panagiotis Angelopoulos¹, Stamatios Katsimperis¹, Ioannis Manolitsis¹, Themistoklis T. Bellos¹, Lazaros Tzelves¹, Marinos Berdempes¹, Andreas Skolarikos¹

¹. Second Department of Urology, National and Kapodistrian University of Athens, Sismanogleio General Hospital, Athens, GRC

Corresponding author: Panagiotis Angelopoulos, angelopoulospanag@gmail.com

Abstract

Gangrenous cystitis is considered a life-threatening but rather rare clinical entity due to the widespread use of antibiotics. We herein report a case of a 78-year-old female with gangrenous cystitis secondary to a spontaneous expanded hematoma of the lesser pelvis who underwent partial cystectomy followed by bilateral ureterostomies with no favorable outcome.

Categories: Urology

Keywords: radical cystectomy, acute abdomen, cystitis, partial cystectomy, gangrenous cystitis

Introduction

Gangrenous cystitis is an extremely uncommon condition. Since 1934, only 204 cases have been described in the literature [1]. In recent years, the use of antibiotics has sharply reduced the number of reported cases [2]. In the majority of instances, conservative interventions supplemented by targeted antibiotic regimens have demonstrated sufficient efficacy. However, in cases marked by advanced clinical presentations, recourse to invasive interventions becomes imperative [2,3]. We hereby present a documented case wherein a patient underwent partial cystectomy to address the condition of gangrenous cystitis caused by microbial pathogens. Regrettably, the patient eventually succumbed. We present a brief review of the literature regarding the etiology, the diagnosis, and the management of this much-overlooked life-threatening infection.

Case Presentation

A 78-year-old patient, a nonsmoker, with a history of coronary artery disease, chronic obstructive pulmonary disease, and arterial hypertension, presented to the emergency department due to dyspnea. The urine culture was negative, as well as the screening for fungi. The patient was admitted to the pulmonary clinic, and having experienced an episode of atrial fibrillation, treatment with low-molecular-weight heparin was initiated. The patient's clinical condition gradually worsened, and ultimately, she was intubated and transferred to the intensive care unit (ICU). During her hospitalization in the ICU, the patient had an episode of macroscopic hematuria, a catheter was placed, and irrigations were performed, but the hematuria persisted, and the patient's clinical condition remained unstable. Therefore, she underwent a computed tomography (CT) scan, which revealed a large extraperitoneal hematoma in the lesser pelvis (Figure 1). A CT cystography followed, revealing contrast extravasation into the prevesical space (Figure 2). The patient consequently underwent an exploratory laparotomy for the removal of the hematoma and the suturing of the bladder rupture, while meticulous hemostasis was achieved without evidence of active bleeding.

How to cite this article

Angelopoulos P, Katsimperis S, Manolitsis I, et al. (September 18, 2023) Gangrenous Cystitis Secondary to a Spontaneous Hematoma of the Lesser Pelvis. Cureus 15(9): e45502. DOI 10.7759/cureus.45502




FIGURE 1: Large extraperitoneal hematoma in the lesser pelvis displacing the urinary bladder to the left (axial)




FIGURE 2: CT cystography showing contrast extravasation into the prevesical space (axial)

CT: computed tomography

The patient on the 10th postoperative day presented with a worsened clinical status and was febrile, and a new CT scan showed a re-expanded hematoma in the lesser pelvis (Figure 3), while an increase in drainage fluids was noticed. A sample was collected and identified as urine.




FIGURE 3: CT scan showing hematoma after the exploratory laparotomy (axial)

CT: computed tomography

Following the deterioration of the patient's clinical condition, a new computed tomography scan revealed the purulence of the hematoma with the presence of air bubbles (Figure 4), possibly due to gas-forming infection. A culture of pus obtained from the wound yielded three kinds of bacteria: *Pseudomonas aeruginosa*, *Enterococcus faecium*, and *Klebsiella pneumoniae*. The antibiotic treatment was upgraded to tigecycline and colistin from the initial regimen of meropenem and linezolid. The laboratory tests showed a noticeable worsening in their values (Table 1).




FIGURE 4: CT scan showing hematoma with air bubbles (axial)

CT: computed tomography

	The 10th postoperative day of exploratory laparotomy	Before the partial cystectomy	Normal range
White blood count	10.420/ μ L	16.700/ μ L	4.000-11.000/ μ L
Neutrophils	84%	91%	40%-75%
Hemoglobin	8.3 g/dL	7.8 g/dL	12-15 g/dL
Hematocrit	26.4%	23.6%	37%-45.6%
Platelets	271.000/ μ L	147.000/ μ L	150.000-400.000/ μ L
Urea	83 mg/dL	97 mg/dL	10-50 mg/dL
Creatinine	0.7 mg/dL	1.3 mg/dL	0.5-1.5 mg/dL
C-reactive protein	55 U/L	205 U/L	<6 U/L

TABLE 1: Laboratory values

The patient underwent a cystoscopy, revealing a significant deficit of the urinary bladder wall. The ureteral orifices were catheterized with double J stents. An exploratory laparotomy followed, necrotic tissue was found, and the area of the bladder triangle was examined without any pathological findings. It was decided to preserve it, and the necrotic area was removed. Partial cystectomy with bilateral ureterostomies was performed. Unfortunately, the patient succumbed after two days in the ICU. The report of the pathological autopsy described extensive lesions of ulcerative cystitis, with areas of interstitial hemorrhage and abscessed necrosis.

Discussion

In the presented case, the most likely cause of bladder necrosis is the infection. The etiological factors might be classified as indirect and direct causes [2-4]. Indirect factors encompass elements that disrupt blood supply, giving rise to vascular complications and bladder ischemia. These factors are internal pressure (caused by prolonged urine retention and subsequent bladder overdistension), external pressure (stemming from conditions such as mispositioned pregnant uterus, extended labor, pelvic tumors, or surgical interventions), and the obstruction of significant arterial and venous pathways (resulting from ligation, embolism, or thrombophlebitis). Direct factors refer to those that inflict direct harm upon the bladder wall, culminating in cellular demise. These encompass a range of substances introduced into the bladder, pelvic radiation exposure, or extensive infection. This infection may arise from systemic sources (such as typhoid or diphtheria) or localized sources [5,6]. The isolated microorganisms span both aerobic and anaerobic varieties. Notably observed in individuals with diabetes, certain gas-producing bacteria can cause emphysematous cystitis, ultimately leading to complete bladder necrosis. Most of the patients resent comorbidities such as neurological conditions or diabetes mellitus [2].

The diagnosis is extremely difficult as the symptoms are not disease-specific. In most cases, the patient presents with dysuria, pyuria, hematuria, and even urinary retention with pain in the lower abdomen [4,7-9]. The condition of acute abdomen manifests as the patient presents in a state of advanced progression of the disease and is commonly concomitant with an unfavorable prognosis.

While computed tomography (CT) is the most useful diagnostic tool, it is typically requested when the patient's clinical presentation indicates a significant severity of symptoms (acute abdomen) [4,6,7]. Cystoscopy in particular is the only examination that permits a correct diagnosis to establish and may allow the debridement of necrotic tissue for culture [5].

The conservative approach, involving appropriate antibiotic therapy and the sealing of the urinary bladder, is often effective [10]. Surgical intervention is required in more advanced cases, where the patient undergoes either a radical or partial cystectomy with the preservation of the trigone [2,11,12]. Fortunately, the trigone usually remains viable because some of its blood supply comes from ureteral and prostatic vessels, and consequently, total cystectomy and urinary diversion are rarely required [5,6]. Gangrenous cystitis is a condition with a poor prognosis and a reported mortality of approximately 35% [2].

Over the recent years, various surgical approaches for replacing a removed bladder have gained prominence. These approaches can be broadly categorized into three main techniques: non-continent urinary diversions, continent non-orthotopic urinary diversions, and orthotopic urinary diversions [13]. Urinary tract infections (UTIs) are highly prevalent and often linked to significant morbidity [14].

Conclusions

The cases of gangrenous cystitis are still observed despite the widespread use of antibiotics. A clinical physician should maintain a high degree of suspicion in most cases to arrive at a diagnosis. Nevertheless, early diagnosis can often prove decisive for patients, as delayed diagnosis increases the mortality rate. Treatment in such cases is surgical.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Devitt AT, Sethia KK: Gangrenous cystitis: case report and review of the literature . J Urol. 1993, 149:1544-5. [10.1016/s0022-5347\(17\)36442-x](https://doi.org/10.1016/s0022-5347(17)36442-x)
- Noulas CN, Markou M, Sotiropoulou R, Diamantidis DE, Karanikas M: Gangrenous cystitis: a rare case report. Cureus. 2023, 15:e36233. [10.7759/cureus.36233](https://doi.org/10.7759/cureus.36233)
- Hinev A, Anakievski D, Krasnaliev I: Gangrenous cystitis: report of a case and review of the literature . Urol Int. 2010, 85:479-81. [10.1159/000321010](https://doi.org/10.1159/000321010)
- Kotoulas K, Georgellis C, Kigitzi E, Liappis A, Spounos M, Tsalkidou V, Patris E: Acute diffuse peritonitis caused by urinary retention: a rare case of gangrenous cystitis. Case Rep Urol. 2018, 2018:4948375. [10.1155/2018/4948375](https://doi.org/10.1155/2018/4948375)
- Ballas K, Rafailidis S, Pavlidis T, Marakis G, Psarras K, Anagnostara E, Sakadamis A: Gangrenous cystitis. Int Urogynecol J Pelvic Floor Dysfunct. 2007, 18:1507-9. [10.1007/s00192-007-0430-9](https://doi.org/10.1007/s00192-007-0430-9)
- Schmidt SS: Gangrene of the bladder due to over distention . J Urol. 1995, 153:409-10. [10.1097/00005392-199502000-00037](https://doi.org/10.1097/00005392-199502000-00037)
- Elalaoui A, Elmoudane A, Mokhtari M, Elfarhaoui H, Motaouakil A, Barki A: Rare case of gangrenous cystitis revealed by acute peritonitis. Urol Case Rep. 2022, 45:102239. [10.1016/j.eucr.2022.102239](https://doi.org/10.1016/j.eucr.2022.102239)
- De Rosa A, Amer T, Waraich N, Bello A, Parkinson R: Gangrenous cystitis in a 42-year-old male . BMJ Case Rep. 2011, 2011:bcr1120103526. [10.1136/bcr.11.2010.3526](https://doi.org/10.1136/bcr.11.2010.3526)
- Daines SL, Hodgson NB: Spontaneous rupture of a necrotic bladder . J Urol. 1969, 102:431-3. [10.1016/s0022-5347\(17\)62166-9](https://doi.org/10.1016/s0022-5347(17)62166-9)
- Boschieter J, Oudshoorn FH, Meuleman EJ, Nieuwenhuijzen JA: Bladder necrosis: 'a man without a bladder'. BMJ Case Rep. 2018, 2018:bcr2016217769. [10.1136/bcr-2016-217769](https://doi.org/10.1136/bcr-2016-217769)
- White MD, Das AK, Kaufman RP Jr: Gangrenous cystitis in the elderly: pathogenesis and management options. Br J Urol. 1998, 82:297-9. [10.1046/j.1464-410x.1998.00741.x](https://doi.org/10.1046/j.1464-410x.1998.00741.x)
- Fujiwara S, Noguchi T, Noguchi T, Emoto A, Tasaki Y: Panperitonitis caused by gangrene of the urinary bladder: report of a successfully treated case. Am Surg. 2008, 74:302-4.
- Symeonidis EN, Falagas ME, Dimitriadis F: Urinary tract infections in patients undergoing radical cystectomy and urinary diversion: challenges and considerations in antibiotic prophylaxis. Transl Androl Urol. 2019, 8:286-9. [10.21037/tau.2019.07.12](https://doi.org/10.21037/tau.2019.07.12)
- Clifford TG, Katebian B, Van Horn CM, et al.: Urinary tract infections following radical cystectomy and urinary diversion: a review of 1133 patients. World J Urol. 2018, 36:775-81. [10.1007/s00345-018-2181-2](https://doi.org/10.1007/s00345-018-2181-2)