

Bracial Artery Injury Resulting From a Dog Bite in a Pediatric Patient: A Case Report

Sarinya Meknarit¹, Adam J. Mann², Faris Azar³, Robert Borrego³

Review began 09/13/2023

Review ended 09/19/2023

Published 09/25/2023

© Copyright 2023

Meknarit et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Surgery, Florida Atlantic University Charles E. Schmidt College of Medicine, Boca Raton, USA 2. General Surgery, Florida Atlantic University, Boca Raton, USA 3. Surgery, St. Mary's Medical Center, West Palm Beach, USA

Corresponding author: Sarinya Meknarit, smeknarit2021@health.fau.edu

Abstract

Pediatric trauma surgeons frequently encounter severe injuries from animal bites, with dog bites being especially prevalent in children, often leading to facial injuries. This paper details the case of a six-year-old male who suffered a dog bite resulting in a rare proximal right brachial artery injury. The bite caused deep lacerations and avulsion injuries, prompting admission to the trauma center, where nonpalpable right radial and ulnar pulses and arm weakness were observed. Surgical intervention, including wound exploration and brachial artery repair using a saphenous vein graft, successfully restored vascular perfusion. This case underscores the urgency of addressing pediatric dog bite injuries through timely exploration, thorough irrigation, and antibiotic prophylaxis, while also highlighting the need for further research on preventive education and clinical guidelines for assessing vascular injuries in such cases.

Categories: Cardiac/Thoracic/Vascular Surgery, General Surgery, Trauma

Keywords: saphenous vein graft (svg), trauma, pediatric, upper extremity ischemia, wound exploration, dog bite wounds, animal bite, traumatic vascular injury

Introduction

Trauma surgery, among other surgical specialties, often bears the brunt of serious injuries from animal bites. The largest subset of these injuries from 1971 to 2018 has been bites from pet canines [1] and is more prevalent in children than adults [2]. In fact, dog bites are among the leading causes of non-fatal emergency room visits in children. Dog bite injury in pediatric patients often results in facial injuries [3], causing both physical and psychological distress [4].

The concerning feature of dog bite injuries is the deep penetration of the puncture wounds, causing injuries to deeper structures including the vasculature. Though vascular injury from a dog bite is uncommon [5], its consequence can be fatal resulting in tissue ischemia if not treated emergently. We present a rare case of a dog bite injury to the proximal right arm in a six-year-old male pediatric patient, resulting in loss of distal perfusion due to an intimal injury and thrombosis of the proximal right brachial artery.

Case Presentation

The patient is a 6-year-old male who presented to the trauma center three hours after he was bitten on his proximal right upper extremity by his family dog, a Belgian Malinois, a breed more commonly used in police and military force. Upon examining the wound, there are deep lacerations and avulsion injuries extending through the muscle layer on the medial and lateral upper right arm and within the axilla. Right radial and ulnar pulses were nonpalpable and not detected using Doppler ultrasonography. He had right arm weakness, but motor and sensory function were intact. Computed tomography angiography (CTA) showed evidence of a proximal right brachial artery thrombosis with no extravasation (Figure 1), and multiple soft tissue avulsion injuries of the upper arm and axilla. The patient was given appropriate antibiotic coverage with ceftriaxone and taken directly to the operating room for right upper extremity wound exploration, repair of brachial artery, and wound washout.

How to cite this article

Meknarit S, Mann A J, Azar F, et al. (September 25, 2023) Bracial Artery Injury Resulting From a Dog Bite in a Pediatric Patient: A Case Report. Cureus 15(9): e45889. DOI 10.7759/cureus.45889



FIGURE 1: Proximal right brachial artery occlusion without evidence extravasation shown on CTA. The occlusion is 2.5cm in length with reconstitution of the distal brachial artery.

In the operating room under general anesthesia, we identified lateral avulsion injuries measuring 6x4cm, 4x3cm, and 4x2cm down to the muscle (Figure 2), and two medial avulsion injuries measuring 7x3cm and 4x2cm down to the muscle layer (Figure 3). We identified a contused brachial artery with intraluminal thrombosis and no perfusion distally, concerning an intimal injury.




FIGURE 2: Right arm lateral avulsion injuries measuring 6x4cm, 4x3cm, and 4x2cm down to muscle layer.



FIGURE 3: Right arm medial avulsion injuries measuring 7x3cm and 4x2cm down to the muscle layer.

We simultaneously turned our attention to the right groin for vein harvest. The right saphenous vein was identified using an ultrasound and harvested. The patient was heparinized with 100 units/kg of unfractionated heparin, and proximal and distal vascular control was obtained (Figure 4). A 2-cm anterior arteriotomy was performed followed by endarterectomy, and balloon thrombectomy proximally and distally, which restored both inflow perfusion and back-bleeding. The intimal flap injury was anchored to the vessel wall with 9-0 prolene, and a saphenous vein venous patch anastomosis was performed with 7-0 prolene in a running fashion (Figure 5). Upon completion of the anastomosis, radial and ulnar pulses were palpable, with triphasic doppler signals to both the radial and ulnar arteries. The wounds were heavily irrigated and partially closed to allow drainage. Patient was extubated and monitored for 24 hours in the pediatric intensive care unit for serial vascular exams.




FIGURE 4: Proximal and distal brachial artery vascular control was obtained in the operating room.




FIGURE 5: Saphenous vein venous patch anastomosis of the brachial artery.

The patient's hospital course was unremarkable with no fevers, chills, wound infection, leukocytosis, or signs of vascular compromise. The patient was taken subsequently three times (hospital day 2, 4, and 6) to the operating room for washout, debridement, and closure of all wounds by tertiary intention. Patient worked extensively with physical and occupational therapy resulting in dramatic improvement in right upper extremity function. The patient completed a one-week course of antibiotic therapy with Amoxicillin-Clavulanate and was discharged on hospital day 9.

Discussion

Over 750,000 dog bite injuries result in emergency department visits, with 1%-2.5% of which requiring hospitalization. Children are at higher risk of dog bites than adults, with the highest incidence occurring in children of five to nine years old [6]. They are commonly single-bite wounds, but multiple wounds are also possible as a result of mauling [7]. A concerning feature of dog bites is the depth of injury resulting in injury to the deeper structures such as arteries and muscles. This is especially true with larger dogs and younger children. As shown in our case, this can result in vascular compromise and ischemia if not treated promptly. Significant complications associated with dog bites are rabies, wound infections, post-traumatic stress disorder, and various manifestations of fear and anxiety [7].

The incidence of rabies is minimal in industrialized countries due to the accessibility of post-exposure rabies prophylaxis and canine rabies vaccination. When encountering a patient with a dog bite, physicians should promptly inquire about the patient's rabies vaccination status and the health status of the involved dog, and assess the patient's need for rabies post-exposure prophylaxis according to the guidelines provided by the CDC [8].

Currently, there is a lack of established clinical guidelines to determine when dog bite wounds should be explored for vascular involvement in patients with normal distal pulses. A retrospective study [9] comparing dog bite injuries with and without vascular injury did not identify any reliable predictive risk factors. This includes factors such as the Glasgow Coma Scale (GCS) upon admission, blood pressure, injury severity score, and the specific type of injury sustained. Based on limited literature, the axillobrachial artery was the most commonly injured followed by the radial artery, with diminished pulses as the most common clinical presentation [4,10]. From our case, we recommend considering surgical wound exploration when there is difficulty visualizing the wound, abnormal Doppler exam, delayed capillary refill, or nonpalpable pulses [11].

Wound infection is the primary complication and sequelae of dog bites. Organisms involved include both aerobic and anaerobic bacteria, and a few fungi [7]. Early and frequent aggressive wound irrigation is recommended to prevent secondary infection, as well as cleaning the area for wound examination [12]. Antibiotics used for prophylaxis should cover a broad spectrum of organisms given the types of microbes found in dog oral flora. Amoxicillin-clavulanate is generally considered a first-line prophylactic treatment for animal bites. We started our patient on a seven-day course of amoxicillin-clavulanate given the severity and depth of the wound. Since hospitalization, the patient had no signs of infection (no fever, chills, purulent drainage from the wound, or leukocytosis).

Wound closure of dog bites is considered contraindicated given the risk of infection. However, there have been studies that showed no significant difference in infection rate between healing by primary and secondary intention [12,13]. However, certain cases may warrant partial closure to promote healing. Thus, we believe that the decision should be based on the severity of the wound. In our case, given the exposure of deeper structures from the bite and the degree of avulsion injury, we partially closed the wound to approximate the skin while also allowing for drainage. The wound was partially closed during each subsequent washout, with all wounds closed by the last washout on hospital day eight. This allowed for the prevention of infection as well as a satisfactory cosmetic outcome.

Most literature studying dog bite injuries is focused on evaluation and management. There is limited research on the effectiveness of strategies used to prevent dog bite injury. A review of two studies evaluating the effectiveness of education as a method of preventing dog bites (in kindergarten and elementary school children) concluded it is still unclear whether it helps reduce rates of dog bite injury [14]. However, this is most likely due to the lack of follow-up to evaluate the incidence of dog bites after education on appropriate behavior around dogs and its consequences was provided. Therefore, more studies in dog bite prevention are needed and future studies should consider evaluating the incidence of dog bites after the intervention. Another important factor in the risk of dog bites is the breed chosen for a house pet. Breeds of dogs are associated with specific temperament, aggressiveness, strength, and activeness. Larger breeds, failure to neuter a dog and selection of male dogs of specific breeds are also associated with more severe wounds and a higher risk of bite injury [3]. Malinois have high prey drive, a trait that can increase the risk of injuries in younger children if not trained properly. This requires tremendous time, patience, and expertise that many households cannot achieve. Therefore, before welcoming a dog to the family, patients should assess breed characteristics, level of responsibility, and safety.

Conclusions

Dog bite wounds can be fatal, especially to smaller children. We presented a case where a six-year-old male sustained a deep avulsion injury to the right shoulder with right brachial artery contusion after being bitten by a Belgian Malinois. This type of injury requires expedient wound irrigation to prevent secondary infection and evaluation of deeper structures. Antibiotic prophylaxis using Amoxicillin-Clavulanate, frequent irrigation, and healing by tertiary intention were highly effective in preventing infection and providing satisfactory cosmetic outcomes. Further research is needed to find effective educational interventions in reducing the incidence of dog bites and to determine the clinical indications for wound exploration to assess for vascular injury.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Acquisition, analysis, or interpretation of data: Sarinya Meknarit

Drafting of the manuscript: Sarinya Meknarit, Adam J. Mann

Concept and design: Adam J. Mann, Faris Azar, Robert Borrego

Critical review of the manuscript for important intellectual content: Adam J. Mann, Faris Azar, Robert Borrego

Supervision: Adam J. Mann, Faris Azar, Robert Borrego

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Israel JS, McCarthy JE, Rose KR, Rao VK: Watch out for wild animals: a systematic review of upper extremity injuries caused by uncommon species. *Plast Reconstr Surg*. 2017, 140:1008-22. [10.1097/PRS.00000000000003754](https://doi.org/10.1097/PRS.00000000000003754)
2. Looke D, Dendle C: Bites (mammalian). *BMJ Clin Evid*. 2010, 2010:0914.
3. Tam B, Matsushima K, Chiba H, et al.: Nationwide analysis of dog bite injuries: different age groups, different injury patterns. *Am Surg*. 2021, 87:1612-5. [10.1177/00031348211024657](https://doi.org/10.1177/00031348211024657)
4. Patterson KN, Horvath KZ, Minneci PC, et al.: Pediatric dog bite injuries in the USA: a systematic review. *World J Pediatr Surg*. 2022, 5:e000281. [10.1136/wjps-2021-000281](https://doi.org/10.1136/wjps-2021-000281)
5. Akingba AG, Robinson EA, Jester AL, et al.: Management of vascular trauma from dog bites. *J Vasc Surg*. 2013, 58:1346-52. [10.1016/j.jvs.2013.05.101](https://doi.org/10.1016/j.jvs.2013.05.101)
6. Greene SE, Fritz SA: Infectious complications of bite injuries. *Infect Dis Clin North Am*. 2021, 35:219-36. [10.1016/j.idc.2020.10.005](https://doi.org/10.1016/j.idc.2020.10.005)
7. Dhillon J, Hoopes J, Epp T: Scoping decades of dog evidence: a scoping review of dog bite-related sequelae. *Can J Public Health*. 2019, 110:364-75. [10.17269/s41997-018-0145-3](https://doi.org/10.17269/s41997-018-0145-3)
8. Manning SE, Rupprecht CE, Fishbein D, et al.: Advisory Committee on Immunization Practices Centers for Disease Control and Prevention (CDC) human rabies prevention-United States, 2008: recommendations of the Advisory Committee on Immunization Practices. *MMWR Recomm Rep*. 2008, 57:1-28.
9. Benfield R, Plurad DS, Lam L, et al.: The epidemiology of dog attacks in an urban environment and the risk of vascular injury. *Am Surg*. 2010, 76:203-5.
10. Cawley OV, Walsh AS, Asghar I, Desmarowitz HU, Antoniou GA: Arterial injury in the upper limb resulting from dog bite. *Ann Vasc Surg*. 2018, 49:314-5. [10.1016/j.avsg.2017.11.053](https://doi.org/10.1016/j.avsg.2017.11.053)
11. Alluri RK, Pannell W, Heckmann N, Sivasundaram L, Stevanovic M, Ghiassi A: Predictive factors of neurovascular and tendon injuries following dog bites to the upper extremity. *Hand (N Y)*. 2016, 11:469-74. [10.1177/1558944715620794](https://doi.org/10.1177/1558944715620794)
12. Hurt JB, Maday KR: Management and treatment of animal bites. *JAAPA*. 2018, 31:27-31. [10.1097/01.JAA.0000531049.59137.cd](https://doi.org/10.1097/01.JAA.0000531049.59137.cd)
13. Paschos NK, Makris EA, Gantsos A, Georgoulis AD: Primary closure versus non-closure of dog bite wounds. A randomised controlled trial. *Injury*. 2014, 45:237-40. [10.1016/j.injury.2013.07.010](https://doi.org/10.1016/j.injury.2013.07.010)
14. Duperrex O, Blackhall K, Burri M, Jeannot E: Education of children and adolescents for the prevention of dog bite injuries. *Cochrane Database Syst Rev*. 2009, CD004726. [10.1002/14651858.CD004726.pub2](https://doi.org/10.1002/14651858.CD004726.pub2)