

Congenital Absence of Patella: A Case Report

Vineeta Pande ¹, Mridu Bahal ¹, Renuka Jadhav ¹, Shailaja Mane ¹, Jasleen Dua ¹

Review began 10/20/2023

Review ended 11/09/2023

Published 11/12/2023

© Copyright 2023

Pande et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Pediatrics, Dr. D.Y. Patil Medical College, Hospital & Research Centre, Pune, IND

Corresponding author: Mridu Bahal, mridubahal@gmail.com

Abstract

Congenital absence of patella is a rare orthopedic condition characterized by an underdeveloped or complete lack of patella. This condition is very rare in isolation and is usually accompanied by other genetic syndromes. The prevalence is difficult to estimate as very few cases of this condition have been reported worldwide. Here, we report a case of congenital bilateral absence of patella in an 18-month-old female child who came with a chief complaint of inability to stand and walk without support, with hyperextension at the knee joint with no other associated abnormalities. The patient was started with active and resisted physiotherapy sessions that alleviated the condition of our patient.

Categories: Pediatrics, Pediatric Surgery, Orthopedics

Keywords: knee joints, nail patella syndrome, gait disorder, congenital abnormalities, patella

Introduction

The patella is the largest sesamoid bone found on the anterior aspect of the distal femur and contributes to forming the knee joint. It plays a critical role in improving the effectiveness of quadriceps extension capacity which is required to maintain normal gait and mobility [1]. Although the exact cause is mostly not clear, genetic predisposition has been attributed as a major contributing factor in many cases [2]. The prevalence rate is not readily available because the patella being completely cartilaginous at birth starts the ossification process between the third and sixth years of life.

It is mostly seen bilaterally and is associated with a clinically diverse and genetically heterogeneous group of disorders. Most commonly associated syndromes include Nail-patella syndrome, Turner-Kieser syndrome, ischiopatellar dysplasia (IPD), Meier-Gorlin syndrome, the RAPADILINO syndrome, the Genito patellar syndrome, and Trisomy 8q syndrome [3]. It is also occasionally associated with various orthopedic malformations, but the most common one is congenital dislocation of the knee or hip and anomalies of femur and fibula, genu recurvatum, genu valgum, cubitus valgus, and pelvic abnormality [4].

Case Presentation

A female child, 18-month-old, came to the OPD with chief complaints of inability to stand and walk without support associated with inability to straighten the knee. The child was born from a non-consanguineous marriage at full term by normal delivery with knees deformed since birth and no history of affected siblings. The child was able to stand unaided for one to two minutes before falling, after which the child was only able to walk with support. All other developmental milestones were age-appropriate. On examination, double knee crease was present, on palpation patella was found to be absent, hyperextension (Figure 1A) and hypermobility were noted at the knee joint, and the patient was also able to do complete flexion (Figure 1B). The power and tone of quadriceps, gluteus maximus, and hamstrings were appropriate for age. Nails were also normal and no other deformity was reported in the patient. Systemic examination was within normal limits. Blood parameters were also normal in range. Ultrasonography of the abdomen and MRI brain were normal. X-rays confirmed the absence of the patella in both knees (Figure 2).

How to cite this article

Pande V, Bahal M, Jadhav R, et al. (November 12, 2023) Congenital Absence of Patella: A Case Report. Cureus 15(11): e48688. DOI 10.7759/cureus.48688



FIGURE 1: Clinical image of the patient showing A) hyperextension at the knee joint and B) complete flexion at the knee joint




FIGURE 2: X-ray image confirming the absence of the patella

Antero-posterior (A) and lateral (B) radiograph of bilateral knee. Arrows indicate the absence of the patella in the anterior aspect of the knee joint.

The child was started on active and resisted physiotherapy and was called for follow-up to assess the improvement in muscle strength, joint stability, and overall mobility. On follow-up, the patient was able to perform passive leg extension and was also able to stand for a longer duration without any support.

Discussion

Bilateral congenital absence of patella is a rare disorder that is associated with gait and mobility-related abnormalities. In some cases, as reported by Duygun et al. (2013), isolated unilateral absence of patella accompanied by contralateral small patella can also be found [3]. In the absence of any other osseous defect, severe lateral dislocation of the extensor mechanism occurs in conjunction with the congenital absence of patella [5]. In such cases or when conservative treatments are ineffective, successful placement of the mechanism in the groove between the femoral condyles, medialization of the tibial tuberosity, and transfer of one or more medial hamstring tendons to the extensor mechanism are the preferred treatments resulting in satisfactory functioning [6]. The tibial tuberosity and the femoral condyles are, in the majority of cases, larger than normal. The well-developed extensor mechanism glides between femoral condyles in the patellar groove. Reports have suggested that strengthening muscles around the knee joint can be pivotal in the management plan [7]. Jerome et al. (2008) also concur that regular physiotherapy and stretching exercises can alleviate the deformity and improve the gait of patients [4]. Considering the previous reports and the satisfactory outcomes, we also started active and resisted physiotherapy. The child showed improvement

and was able to stand for a longer duration without any support.

Conclusions

Bilateral congenital absence of patella causes gait and mobility-related abnormalities and has a severe impact on the patient's quality of life. Treatment is directed toward supportive care to prevent major disability in patients. With early diagnosis and multimodal treatment, including physiotherapy and reconstructive surgical intervention where necessary, patients with this condition can lead a fulfilling life with improved joint function and reduced abnormalities.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Mridu Bahal, Vineeta Pande, Renuka Jadhav, Shailaja Mane

Acquisition, analysis, or interpretation of data: Mridu Bahal, Vineeta Pande, Renuka Jadhav, Jasleen Dua

Drafting of the manuscript: Mridu Bahal, Vineeta Pande

Critical review of the manuscript for important intellectual content: Mridu Bahal, Vineeta Pande, Renuka Jadhav, Shailaja Mane, Jasleen Dua

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Fox AJ, Wanivenhaus F, Rodeo SA: The basic science of the patella: structure, composition, and function . J Knee Surg. 2012, 25:127-41. [10.1055/s-0032-1313741](#)
2. Samuels ME, Campeau PM: Genetics of the patella. Eur J Hum Genet. 2019, 27:671-80. [10.1038/s41431-018-0329-6](#)
3. Duygun F, Sertkaya O, Aldemir C, Doğan A: Isolated unilateral absent patella and contralateral small patella . BMJ Case Rep. 2013, 2013:[10.1136/bcr-2013-200353](#)
4. Jerome JT, Varghese M, Sankaran B, Thomas S: Bilateral congenital absence of patella . Indian J Orthop. 2008, 42:228-30.
5. Yu A, Russo M, Kothakapa D, Huang P, Mantica A: Dislocated extensor mechanism in a child with bilateral absent patella without other congenital abnormalities. J Am Acad Orthop Surg Glob Res Rev. 2023, 7:e23.00117. [10.5435/JAOSGlobal-D-23-00117](#)
6. Iliadis AD, Jaiswal PK, Khan W, Johnstone D: The operative management of patella malalignment . Open Orthop J. 2012, 6:327-39. [10.2174/1874325001206010327](#)
7. Duncan JG, Souter WA: Hereditary onycho-osteodysplasia. J Bone Joint Surg Br. 1963, 45:242-58. [10.1302/0301-620x.45b2.242](#)