

Charting the Unknown: Green Urine After Propofol in Pediatric Trauma

Review began 01/25/2024
Review ended 03/12/2024
Published 03/20/2024

© Copyright 2024

El Farh et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Reda El Farh ^{1,2}, Othman Moueqqit ³, Zakaria Bouayed ², Ilias El Kadiri Boutchich ², Omar Alaoui Mhammedi ², Wael El Fergui ², Ghizlane El Aidouni ², Houssam Bkiyar ², Brahim Housni ²

1. Faculty of Medicine, Mohammed First University, Oujda, MAR 2. Anesthesiology and Intensive Care Unit Department, Mohammed VI University Hospital, Oujda, MAR 3. General Medicine, Faculty of Medicine and Pharmacy, Mohammed First University, Oujda, MAR

Corresponding author: Reda El Farh, md.redaelfarh@gmail.com

Abstract

The phenomenon of green urine discoloration, while rare, represents a captivating clinical puzzle that challenges the distinction between benign and pathological conditions. In this report, we present an intriguing case involving a 15-year-old trauma patient admitted following a motorcycle collision, where the ensuing unconsciousness necessitated propofol induction for intubation and sedation. Remarkably, around 48 hours post-admission, the patient displayed green urine discoloration, which resolved spontaneously within just 12 hours. This case serves as a compelling illustration of the uncommon occurrence of propofol-induced green urine in the context of critical care management, underscoring the imperative need to discern and appreciate medication-related chromatic alterations in urine.

Categories: Anesthesiology, Emergency Medicine, Nephrology

Keywords: critical care management, chromatic alterations, pediatric head trauma, propofol, green urine

Introduction

Green urine discoloration can occur due to various factors, such as medications, dyes, infections, ingested substances, and numerous other etiological agents [1]. Propofol (2,6 diisopropylphenol), a frequently employed anesthetic agent, has, in infrequent instances, been documented as a potential cause of green urine discoloration [1]. This can lead to undue concern for healthcare providers and the necessity for additional laboratory testing when not recognized by the attending clinician [2]. The scientific literature contains limited documentation concerning the phenomenon of urine turning green as a consequence of propofol administration [2]. In this case study, we present an instance of green urine discoloration observed in a pediatric patient who received propofol sedation following a head trauma.

Case Presentation

A 15-year-old patient with no notable medical history was admitted to the hospital for severe trauma following a motorcycle collision with a car. The initial clinical examination during the resuscitation process revealed an unconscious patient with a Glasgow Coma Scale score of 11/15, exhibiting agitation. The patient had equal and reactive pupils, with a blood pressure of 110/50 mmHg, heart rate of 95 beats per minute, respiratory rate of 16 cycles per minute, and oxygen saturation of 97% on room air.

A primary survey revealed a lip laceration and multiple abrasions on the face and both knees. Rapid sequence induction was performed, involving the administration of 100 mg of propofol, 70 mg of rocuronium, and 200 µg of fentanyl. The patient was intubated using a size 4 laryngoscope blade, and an endotracheal tube of size 6.5 was inserted and secured at a depth of 21 cm. Controlled ventilation was initiated with the following parameters: a tidal volume of 420 mL, respiratory rate of 14 breaths per minute, positive end-expiratory pressure set at 3 cmH₂O, fraction of inspired oxygen of 100%, and an inspiratory to expiratory ratio of 1:2.

Nasogastric and urinary catheterizations were performed, resulting in 200 mL of normal-looking urine being drained by the urinary catheter. Subsequently, a whole-body CT scan was conducted, revealing multiple findings. In the cerebral region, a minor left parietal hemorrhagic contusion was noted, along with a hemorrhage in the midbrain. Additionally, an intraventricular hemorrhage was observed (Figure 1). In the thoracic region, bilateral lower lobe parenchymal consolidations were noted, suggestive of contusions (Figure 2). Furthermore, in the osseous window, fractures were identified, including a fracture line of the nasal bones, a fracture line extending from the right condyle of the mandible to the ipsilateral tympanic bone, and a fracture line of the left jugular foramen (Figure 2).

How to cite this article

El Farh R, Moueqqit O, Bouayed Z, et al. (March 20, 2024) Charting the Unknown: Green Urine After Propofol in Pediatric Trauma. Cureus 16(3): e56588. DOI 10.7759/cureus.56588




FIGURE 1: Cerebral region imaging showing a minor hemorrhagic contusion in the left parietal lobe (A), midbrain hemorrhage (B), and intraventricular hemorrhage (C).




FIGURE 2: Thoracic and osseous region findings.

A: Thoracic imaging indicating parenchymal consolidations in the lower lobes. B: Osseous window imaging highlighting a fracture line in the nasal bones along with a fracture line extending from the right condyle of the mandible to the ipsilateral tympanic bone. C: Osseous window imaging showing a fracture line in the left jugular foramen.

Laboratory tests, including renal and hepatic function, showed normal results. Arterial blood gas analysis revealed normal values, with lactate levels measured at 0.68 mmol/L.

A consultation with the neurosurgery team was obtained, and the decision was made to maintain sedation using propofol and midazolam, as there was no indication for surgical intervention. Additionally, the patient was started on levetiracetam 500 mg every 12 hours for seizure prophylaxis. Approximately 48 hours later, a change in urine color to green was noted (Figure 3). However, subsequent laboratory investigations, including urine culture, remained unremarkable.



FIGURE 3: Grass-green-colored urine with Foley catheter in situ.

Approximately 12 hours later, the green discoloration of the urine spontaneously resolved. After 14 days, the patient was successfully extubated and subsequently transferred to the neurosurgery department for further care.

Discussion

Numerous intrinsic and extrinsic elements have the potential to induce a green hue in urine. Medications such as cimetidine, promethazine, amitriptyline, indomethacin, metoclopramide, methocarbamol, and flutamide, along with dyes such as methylene blue, indigo blue, biliverdin, and food dyes, as well as metabolic conditions such as Hartnup disease, *Pseudomonas aeruginosa* infections, and vesical fistulas, constitute additional plausible etiological factors for the manifestation of green urine discoloration [2].

Pedersen et al. [3] documented that the development of discoloration becomes evident when the clearance of propofol surpasses hepatic elimination, subsequently leading to extrahepatic elimination of propofol. Whereas Shioya et al. [4] noted that green urine was associated with enterohepatic circulatory issues due to constipation and peristalsis impairment, as well as enhanced carrier protein utilization from albumin and erythrocyte administration, along with extrahepatic glucuronidation in the kidneys. However, our patient had no prior or current hepatic complications according to the biological evaluations. Hence, while phenolic derivatives of propofol are suspected culprits for the green urine discoloration, its exact origin remains unclear.

The green discoloration resolved on its own within about 12 hours after propofol discontinuation, consistent with previous reports that have noted variations in onset times, ranging from as early as two hours to up to two days after discontinuing propofol [5].

This case highlights the need for a comprehensive medical history and a thorough evaluation of medication usage, as even commonly administered drugs such as propofol can have unexpected effects. It serves as a reminder for healthcare professionals to maintain vigilance in identifying such atypical presentations and encourages further exploration of the underlying mechanisms behind this phenomenon.

Conclusions

This case of green urine following propofol in a pediatric trauma patient, though rare and initially puzzling, ultimately resolved spontaneously without consequence. It underscores the need for careful monitoring, recognition, and management of medication-related urine color changes. This contributes to medical knowledge, emphasizing the importance of sharing such cases within the healthcare community, demonstrating that even uncommon occurrences can be effectively managed with clinical expertise.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Reda El Farh, Othman Moueqqit, Zakaria Bouayed, Ilias El Kadiri Boutchich, Omar Alaoui Mhammedi, Ghizlane El Aidouni, Houssam Bkiyar, Brahim Housni, Wael El Fergui

Acquisition, analysis, or interpretation of data: Reda El Farh, Othman Moueqqit, Zakaria Bouayed, Ilias El Kadiri Boutchich, Omar Alaoui Mhammedi, Ghizlane El Aidouni, Houssam Bkiyar, Brahim Housni, Wael El Fergui

Drafting of the manuscript: Reda El Farh, Othman Moueqqit, Zakaria Bouayed, Ilias El Kadiri Boutchich, Omar Alaoui Mhammedi, Ghizlane El Aidouni, Houssam Bkiyar, Brahim Housni, Wael El Fergui

Critical review of the manuscript for important intellectual content: Reda El Farh, Othman Moueqqit, Zakaria Bouayed, Ilias El Kadiri Boutchich, Omar Alaoui Mhammedi, Ghizlane El Aidouni, Houssam Bkiyar, Brahim Housni, Wael El Fergui

Supervision: Ghizlane El Aidouni, Houssam Bkiyar, Brahim Housni

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Elhamdaoui O, Rhaymi C, Meftah H, et al.: Green urine after propofol infusion . International Society of Pharmacovigilance (ISoP). 2019, 1238.
2. Müge Ç, Çelik M, Sibel T: Green urine: a rare side effect of propofol in a pediatric case . Am J Clin Case Rep. 2021, 2:1046.
3. Pedersen AB, Kobborg TK, Larsen JR: Grass-green urine from propofol infusion . Acta Anaesthesiol Scand. 2015, 59:265-7. [10.1111/aas.12437](https://doi.org/10.1111/aas.12437)
4. Shioya N, Ishibe Y, Shibata S, et al.: Green urine discoloration due to propofol infusion: a case report . Case Rep Emerg Med. 2011, 2011:242514. [10.1155/2011/242514](https://doi.org/10.1155/2011/242514)
5. Rawal G, Yadav S: Green urine due to propofol: a case report with review of literature . J Clin Diagn Res. 2015, 9:OD03-4. [10.7860/JCDR/2015/15681.6706](https://doi.org/10.7860/JCDR/2015/15681.6706)