

# Right Upper Quadrant Pain: A Rare Presentation of Diverticulitis

Received 01/02/2024  
Review began 02/09/2024  
Review ended 03/03/2024  
Published 03/06/2024

© Copyright 2024

Atalay et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Rediet Tefera Atalay<sup>1</sup>, Oluwapelumi Kolawole<sup>2</sup>, Girma M. Ayele<sup>1</sup>, Abay A. Gobeze<sup>1</sup>, Angesom Kibreab<sup>3</sup>, Miriam B. Michael<sup>4, 5</sup>

1. Internal Medicine, Howard University Hospital, Washington DC, USA 2. Internal Medicine, Howard University College of Medicine, Washington DC, USA 3. Gastroenterology, Howard University Hospital, Washington DC, USA 4. Internal Medicine, Howard University Hospital, Washington DC, USA 5. Internal Medicine, University of Maryland, Baltimore, USA

**Corresponding author:** Miriam B. Michael, mmichael@som.umaryland.edu

## Abstract

Acute diverticulitis is a prevalent medical condition with increasing incidence rates. While the sigmoid colon is the most commonly affected part of the large intestine, there have been occurrences of right-sided diverticulitis although uncommon. We present a case report highlighting the atypical presentation of diverticulitis in a 27-year-old female patient. The patient experienced right upper and epigastric pain and was ultimately diagnosed with right-sided diverticulitis, supported by her CT imaging findings. This uncommon presentation underscores the importance of considering diverse clinical manifestations when diagnosing and managing diverticulitis.

**Categories:** Gastroenterology, Internal Medicine

**Keywords:** appendicitis, right-sided diverticulitis, diverticula, diverticulitis, right upper quadrant pain

## Introduction


Diverticular disease is more prevalent in older individuals within Western populations, affecting nearly 65% of the population by the age of 85 [1]. The prevalence of diverticulitis is greater among men under the age of 50, but it becomes increasingly apparent among older women. Hospitalization rates were higher for men under 50, whereas they were elevated for women over 50 in multiple hospital review datasets. Studies indicate that diverticulosis is found in less than 10% of individuals under the age of 40 [2,3]. In most cases, patients remain asymptomatic throughout their lives, with the diagnosis often made incidentally during endoscopy or cross-sectional imaging [4]. Various factors, including advancing age, low dietary fiber intake, reduced physical activity, obesity, smoking, and heavy alcohol consumption, elevate the risk of developing acute diverticulitis [5]. The severity of diverticulitis varies widely, from mild abdominal pain to the more severe development of an acute abdomen due to perforation and peritonitis [6]. In Western countries, right-sided diverticulitis is uncommon, representing only 1.5% of recorded cases [7]. This case report details a rare occurrence of right-sided diverticulitis in a 27-year-old African American female.

## Case Presentation

A 27-year-old African American female with an unremarkable medical and surgical history presented to the Emergency Department with acute abdominal discomfort localized in the epigastric and right upper quadrant regions, originating two days before her visit. She described the pain as severe, with a 10/10 sharp sensation that extended to her back. Concurrently, she reported experiencing symptoms such as nausea, reduced appetite, headache, chills, and night sweats over the same two-day period. Although the patient did not subjectively report fever, her temperature upon presentation was 39.1°C. Other vital signs were stable and on physical examination, there was epigastric and right upper quadrant tenderness. Laboratory results showed leukocytosis of 21,460 with left shift. The urine pregnancy test was negative, and lipase, kidney, and liver function tests were normal. Ultrasound (US) of the abdomen and pelvis showed a benign left ovarian cyst. CT abdomen pelvis w/ contrast showed ascending colon/transverse colon/hepatic flexure diverticulosis with hepatic flexure wall thickening with pericolic fat stranding suggestive of acute diverticulitis. Figures 1 and 2 show the CT scan of the abdomen with axial and coronal view, respectively.


### How to cite this article

Atalay R, Kolawole O, Ayele G M, et al. (March 06, 2024) Right Upper Quadrant Pain: A Rare Presentation of Diverticulitis. Cureus 16(3): e55674. DOI 10.7759/cureus.55674



**FIGURE 1: CT abdomen - axial view**

The arrow points towards the diverticula.



**FIGURE 2: CT abdomen - coronal view**

CT of the abdomen with contrast showing diverticulosis in the transverse colon, hepatic flexure, and thickening of the hepatic flexure wall, accompanied by pericolic fat stranding. The arrow points towards the diverticula.

Our patient initially received IV ceftriaxone and metronidazole, fluids, and analgesics. Antibiotics were later switched to oral ciprofloxacin and metronidazole. The patient has been scheduled for an outpatient colonoscopy in six weeks and advised to follow up with colorectal surgery, as a surgical intervention may be necessary in the long term.

## Discussion

While left-sided diverticulitis is prevalent in Western populations, right-sided diverticulitis is considered uncommon [8]. The Asian population, however, exhibits a higher incidence of right-sided diverticulitis, with genetic factors and diet presumed to influence its onset [1]. The rising occurrence of diverticular disease in other regions of the world may be attributed to the adoption of Western lifestyle and dietary habits [1]. US and CT scans play a crucial role in distinguishing diverticulitis from acute appendicitis, especially since relying solely on history and physical examination can be challenging [9]. Although right-sided diverticulitis tends to be less severe and predominantly occurs in young males compared to its left-sided counterpart, which is associated with worse laboratory findings such as elevated C-reactive protein (CRP), creatinine, and low hemoglobin and albumin [10].

CT scanning is considered the superior imaging modality for diagnosing possible acute right-sided colonic diverticulitis (ARCD) [11]. However, US is a more economical and radiation-free alternative, which is particularly relevant for younger patients with right-sided diverticulitis. US features can offer valuable preoperative diagnostic information, such as diverticular wall thickening, surrounding echogenic fat, and intra-diverticular echogenic material [12,13]. While US demonstrates good diagnostic accuracy for diverticulitis compared to CT scans, it is less effective in assessing the extent of large abscesses and detecting free air. Moreover, its accuracy is compromised in obese patients and those with obstructing gas. However, US remains a preferable option for pregnant women to avoid exposure to ionizing radiation [14]. However, the quality of US results relies on the operator's expertise. In cases where US findings are inconclusive, a contrast-enhanced CT scan can be a useful complementary tool [12,13].

The diagnostic and treatment principles for ARCD closely mirror those for acute left-sided colonic diverticulitis (ALCD). Distinct management guidelines are lacking for ARCD. Non-operative approaches are typically favored, especially in cases without widespread peritonitis, although distinguishing between benign and malignant instances before surgery can pose challenges [15]. The utilization of antibiotic medication for treating uncomplicated acute diverticulitis is deemed crucial, though its use remains a subject of controversy, with selective administration recommended for patients with uncomplicated acute diverticulitis [9]. Surgical intervention is commonly reserved for complicated cases [16].

In a study by Zhin Chen et al. on 111 cases of cecal and ascending colonic diverticulitis, 13 patients underwent resection and repair for right-sided colonic diverticulitis, while no patients had colectomy. Also, 15 patients with right-sided colonic diverticulitis underwent abdominal drainage, and the remainder were managed nonoperatively. The length of hospital stay did not significantly differ among the three treatments for right-sided colonic diverticulitis [17].

A review of observational cohort studies on nonoperative management of right diverticulitis that included 1,584 adult participants from 11 studies to evaluate the primary outcomes of the recurrence rate and morbidity associated with recurrence was done. The pooled recurrence rate over a median follow-up of 34.2 months was 12% (95% CI, 10%-15%). Only 9.9% of patients required urgent surgery at the first recurrence, and there were no reported mortalities [16]. The study concluded that non-operative management in right-side diverticulitis is safe, and the risk of recurrence is rare [16].

A recent meta-analysis encompassing 38 studies and involving 10,129 patients examined the outcomes of non-operative management. The analysis revealed low rates of recurrence and treatment failure for both right-sided and left-sided Crohn's disease (CD), with left-sided CD showing a slightly higher recurrence rate. Interestingly, the recurrence rates were similar regardless of whether patients received antibiotics for uncomplicated CD or not [18].

## Conclusions

Diverticulitis traditionally manifests with left lower quadrant pain and is primarily diagnosed in the elderly population. According to available literature, right-sided diverticulitis is rare and appears to occur more frequently among Asian populations. Its manifestation with symptoms such as abdominal pain in the right lower quadrant, tenderness, nausea, or vomiting may sometimes be mistaken for acute appendicitis, further complicating diagnosis. We emphasize the significance of considering diverticulitis as a potential diagnosis in atypical cases, especially in younger populations. Studies indicate that once the diagnosis is established, there are no distinct management guidelines for right-sided diverticulitis, and the management approach is similar to that of left-sided diverticulitis.

## Additional Information

### Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

**Concept and design:** Miriam B. Michael, Girma M. Ayele, Rediet Tefera Atalay, Oluwapelumi Kolawole,

Angesom Kibreab

**Drafting of the manuscript:** Miriam B. Michael, Girma M. Ayele, Rediet Tefera Atalay, Oluwapelumi Kolawole

**Critical review of the manuscript for important intellectual content:** Miriam B. Michael, Rediet Tefera Atalay, Oluwapelumi Kolawole, Angesom Kibreab, Abay A. Gobezie

**Supervision:** Miriam B. Michael

**Acquisition, analysis, or interpretation of data:** Girma M. Ayele, Rediet Tefera Atalay, Abay A. Gobezie

## Disclosures

**Human subjects:** All authors have confirmed that this study did not involve human participants or tissue.

**Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

## References

1. Kamal MU, Baiomi A, Balar B: Acute diverticulitis: a rare cause of abdominal pain . *Gastroenterology Res.* 2019, 12:203-7. [10.14740/gr1166](#)
2. Painter NS, Burkitt DP: Diverticular disease of the colon: a deficiency disease of Western civilization . *Br Med J.* 1971, 2:450-4. [10.1136/bmj.2.5759.450](#)
3. Turner GA, O'Grady MJ, Purcell RV, Frizelle FA: The epidemiology and etiology of right-sided colonic diverticulosis: a review. *Ann Coloproctol.* 2021, 37:196-203. [10.3393/ac.2021.00192.0027](#)
4. Hajibandeh S, Hajibandeh S, Smart NJ, Maw A: Meta-analysis of the demographic and prognostic significance of right-sided versus left-sided acute diverticulitis. *Colorectal Dis.* 2020, 22:1908-23. [10.1111/codi.15328](#)
5. Kaiser AM, Jiang JK, Lake JP, et al.: The management of complicated diverticulitis and the role of computed tomography. *Am J Gastroenterol.* 2005, 100:910-17. [10.1111/j.1572-0241.2005.41154.x](#)
6. Colonic diverticulosis and diverticular disease: epidemiology, risk factors, and pathogenesis . (2023). Accessed: February 28, 2024: <https://www.uptodate.com/contents/colonic-diverticulosis-and-diverticular-disease-epidemiology-risk-factors-and-patho....>
7. Lee JH, Ahn BK, Lee KH: Conservative treatment of uncomplicated right-sided diverticulitis: a systematic review and meta-analysis. *Int J Colorectal Dis.* 2021, 36:1791-9. [10.1007/s00384-021-03913-x](#)
8. Kahveci S, Tokmak TT, Yildirim A, Mavili E: Acute right-sided colonic diverticulitis mimicking appendicitis: a report of two cases. *J Clin Ultrasound.* 2013, 41:238-41. [10.1002/jcu.21967](#)
9. Huang SS, Sung CW, Wang HP, Lien WC: The outcomes of right-sided and left-sided colonic diverticulitis following non-operative management: a systematic review and meta-analysis. *World J Emerg Surg.* 2022, 17:56. [10.1186/s13017-022-00463-7](#)
10. Espinosa J, Sharma R, Lucerna A, Stranges D: Medical approach to right colon diverticulitis with perforation. *Case Rep Emerg Med.* 2017, 2017:2563218. [10.1155/2017/2563218](#)
11. Kim SH, Byun CG, Cha JW, Choi SH, Kho YT, Seo DY: Comparative study of the clinical features and treatment for right and left colonic diverticulitis. *J Korean Soc Coloproctol.* 2010, 26:407-12. [10.3393/jksc.2010.26.6.407](#)
12. Sartelli M, Weber DG, Kluger Y, et al.: 2020 update of the WSES guidelines for the management of acute colonic diverticulitis in the emergency setting. *World J Emerg Surg.* 2020, 15:32. [10.1186/s13017-020-00313-4](#)
13. Destek S, Gül VO: Effectiveness of conservative approach in right colon diverticulitis . *Ulus Travma Acil Cerrahi Derg.* 2019, 25:396-402.
14. Kim JY, Park SG, Kang HJ, et al.: Prospective randomized clinical trial of uncomplicated right-sided colonic diverticulitis: antibiotics versus no antibiotics. *Int J Colorectal Dis.* 2019, 34:1413-20. [10.1007/s00384-019-03343-w](#)
15. Chen Z, Zhang B, Wu D, Jin Y: Characteristics of predominantly right-sided colonic diverticulitis without need for colectomy. *BMC Surg.* 2020, 20:202. [10.1186/s12893-020-00863-z](#)
16. Lee YF, Tang DD, Patel SH, Battaglia MA, Shanker BA, Cleary RK: Recurrence of acute right colon diverticulitis following nonoperative management: a systematic review and meta-analysis. *Dis Colon Rectum.* 2020, 63:1466-73. [10.1097/DCR.0000000000001787](#)
17. Weizman AV, Nguyen GC: Diverticular disease: epidemiology and management. *Can J Gastroenterol.* 2011, 25:385-9. [10.1155/2011/795241](#)
18. Mazzei MA, Guerrini S, Cioffi Squitieri N, et al.: The role of US examination in the management of acute abdomen. *Crit Ultrasound J.* 2013, 5:S6. [10.1186/2036-7902-5-S1-S6](#)