

Rare Insights: Temporoparietal Necrotizing Fasciitis Stemming From a Dental Source

Chetan Gupta ¹, Nitin Bhola ¹, Parmarth Sonpal ¹, Nidhi Bodiwala ¹

1. Oral & Maxillofacial Surgery, Sharad Pawar Dental College & Hospital, Wardha, Wardha, IND

Corresponding author: Parmarth Sonpal, palmsonpal@gmail.com

Received 02/21/2024

Review began 03/13/2024

Review ended 05/11/2024

Published 05/14/2024

© Copyright 2024

Gupta et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

DOI: 10.7759/cureus.60278

Abstract

Necrotizing fasciitis (NF) of the face is a rare yet serious condition requiring prompt and comprehensive management. This approach typically involves input from various medical specialties such as infectious disease specialists, critical care physicians, and surgeons. The primary goals are early recognition, aggressive surgical debridement, appropriate antibiotic therapy, and supportive care. Prompt diagnosis is crucial, based on symptoms like severe pain, rapidly spreading erythema, and systemic signs of infection. Broad-spectrum antibiotics are initiated empirically, and adjusted based on culture results. Urgent surgical debridement is crucial, removing all necrotic tissue. Careful consideration must be given to preserve vital structures. Close monitoring and intensive care may be necessary, especially for severe cases. Soft tissue reconstruction may follow once the infection is controlled, aiming to restore function and aesthetics. Long-term follow-up is essential to observe for complications and recurrence.

Categories: Other, Geriatrics, Dentistry

Keywords: odontogenic, anesthesia, infratemporal, fascial spaces, necrotizing fasciitis (nf)

Introduction

Necrotizing fasciitis (NF) is a severe soft tissue infection primarily spreading through fascial planes. It is usually accompanied by systemic inflammatory response syndrome (SIRS) and needs prolonged intensive care treatment [1]. Although necrotizing fasciitis can affect various parts of the body, including the extremities, abdomen, and perineum, it can also arise in the oro-pharyngeal and maxillofacial regions [2]. In the head and neck region, odontogenic causes are most prevalent, leading to rapid tissue infiltration, vascular compromise, and potential organ failure or limb loss [3]. The condition is commonly referred to as the "flesh-eating disease" due to its aggressive nature and ability to cause extensive tissue destruction. Early NF diagnosis is challenging due to subtle symptoms unless triggered by toxic shock syndrome or organ failure [3]. Advanced cases manifest with significant tissue loss, vascular complications, or systemic involvement, often leading to fatal outcomes. If not treated at once with appropriate drugs or surgery, it may spread through the bone marrow and cortex, reaching muscle layers, fascial spaces, and thence various vital organs [4]. Aggressive forms are often the result of underestimation of the infection, late diagnosis, erroneous surgical approach, and septic complications. However, NF involving the temporalis muscle remains infrequently reported. In the current study, we present a case of Temporal Necrotizing Fasciitis due to a suspected odontogenic cause in a geriatric patient.

Case Presentation

A 77-year-old female patient presented to our outpatient department with the complaint of severe pain and pus discharge from the right side of the head (Figure 1) with general malaise in the past 10-15 days approximately. The pain was gradual in onset, throbbing, continuous, and localized in nature. While inspecting inflamed, slough tissue with draining pus from the right tempo-parietal and right retromolar region.

How to cite this article

Gupta C, Bhola N, Sonpal P, et al. (May 14, 2024) Rare Insights: Temporoparietal Necrotizing Fasciitis Stemming From a Dental Source. Cureus 16(5): e60278. DOI 10.7759/cureus.60278



FIGURE 1: Pre-operative image showing necrotizing fasciitis over right temporal region

Pre-operative image showing necrotizing fasciitis over right temporal region with sloughed margins and exposed temporal bone.

Upon examination, there was crepitus and fluctuant swelling noted in the right temporo-parietal region. A thorough intra-oral examination was done for the patient upon which a grossly decayed tooth in the lower right mandibular region was found. Contrast Enhanced Computed Tomograph (CECT) of the Head (Figure 2, Figure 3) was done for this patient, which showed evidence of peripherally enhancing collection with internal enhancing septations noted over the right fronto-parieto-temporal regions with multiple air density foci within. The collection approximately measured 9.5 x 2.9 x 12 cm. Inferiorly collection is seen extending into the right infratemporal fossa, masticator, and submandibular spaces.




FIGURE 2: Axial Section of Contrast-Enhanced Computed Tomograph (CECT) Head

There is evidence of peripherally enhancing collection with internal enhancing septations within noted over the right frontoparieto temporal regions with multiple air density foci within. The collection approximately measures 9.5 x 2.9 x 12 cm (Axial Plane Topography with Volume Computerized Tomography Contrast (APTVCC)). Inferiorly collection is seen extending into the right infratemporal fossa, masticator, and submandibular spaces. There is involvement of the medial pterygoid and superficial and deep fibers of the temporalis muscle. Masseter muscle also shows heterogeneous enhancement and is mildly bulky as compared to the opposite side.



FIGURE 3: Coronal Section of Contrast-Enhanced Computed Tomograph (CECT) Head

There is evidence of peripherally enhancing collection with internal enhancing septations within noted over the right frontoparieto temporal regions with multiple air density foci within (coronal section).

The general condition of the patient was poor with severe nutritional cachexia, Body Mass Index of 10.82, reduced air entry in the bilateral lower lobe, and crepitus in the left lower lobe of the lung. Following routine investigations were sent for the patient (Table 1).

Investigation	Observed Value	Biological Reference Range	Method
Haemoglobin (Hb)	6.8 gm%	Male: 13-17 gm% Female: 12-15 gm%	Photometric Measurement
Total WBC Count	23500/cu.mm	Male: 4000-10000 cu.mm Female: 4000-10000 cu.mm	Coulter Principal
Red Blood Cells (RBCs)	2.79 millions/cu.mm	Male: 4.5-5.5 millions/cu.mm Female: 3.8-4.8 millions/cu.mm	Coulter Principal
Total Platelet Count	6.91 Lacs/cu.mm	Male: 1.50-4.10 Lacs/cu.mm Female: 1.50-4.10 Lacs/cu.mm	Coulter Principal
HbA1c	4.0 % A1C NGSP	Non Diabetic- <6.0 % A1C NGSP, Action suggested- >7.0 % A1C NGSP, ADA Target- 6.0-7.0 % A1C NGSP	Enzymatic

TABLE 1: Pre-operative Investigations
NGSP: National Glycohemoglobin Standardization Program; HbA1c/A1C: glycated hemoglobin; ADA: American Diabetes Association

The patient was planned for surgical drainage of space infection under general anesthesia. After pre-anesthetic checkup high risk and poor prognosis fitness was obtained. The patient’s relatives were not willing to get operated on under general anesthesia, hence it was planned under local anesthesia. Before starting the antibiotics, we sent pus from the extraoral temporoparietal and intraoral retromandibular region for the culture and sensitivity test. We initiated empirical antibiotic therapy for the patient until the final culture report was available.

Incision and drainage under local anesthesia were done. Right auriculotemporal nerves, zygomaticotemporal, lesser occipital, posterior auricular nerve and mandibular nerve blocks were given. Incision and drainage of the right superficial temporal, deep temporal, buccal, lateral pharyngeal, and submandibular space were done. The extraction of 2nd and 3rd right mandibular molar teeth with through curettage was done. The dead and decayed soft tissue debridement is done meticulously. The submandibular space was explored and through the same incision lateral pharyngeal space exploration and drainage followed by corrugated rubber drains (Figure 4) were placed in respective areas.



FIGURE 4: Post-operative image showing corrugated rubber drain placement over the right temporal and submandibular region

It is also showing signs of healing over the right temporal region.

The patient started on IV antibiotics according to the blood counts. After 24 hours of incubation, no growth was there in the sent specimen for culture sensitivity. Post-operative blood transfusion and supportive care were given. There was an improvement in the patient's general condition the post-operative routine labs are given in Table 2.

Investigation	Observed Value	Biological Reference Range	Method
Haemoglobin (Hb)	9.2 gm%	Male: 13-17 gm% Female: 12-15 gm%	Photometric Measurement
Total WBC Count	11500/cu.mm	Male: 4000-10000 cu.mm Female: 4000-10000 cu.mm	Coulter Principal
Red Blood Cells (RBCs)	3.55 millions/cu.mm	Male: 4.5-5.5 millions/cu.mm Female: 3.8-4.8 millions/cu.mm	Coulter Principal
Total Platelet Count	6.29 Lacs/cu.mm	Male: 1.50-4.10 Lacs/cu.mm Female: 1.50-4.10 Lacs/cu.mm	Coulter Principal

TABLE 2: Postoperative Investigations

Due to some circumstances, the patient's relatives discontinued the treatment and took discharge against medical advice.

Discussion

The infection of the second or third mandibular molar is the frequent reason for odontogenic necrotizing fasciitis (NF) [5]. This infection may travel cranially up to the skull base or caudally towards the thorax [6]. The mortality rate for NF patients is 9.8%, without diabetes mellitus [7]. NF can result from various bacteria, including *Streptococcal*, *Staphylococcal*, *Fusobacterium*, or *Acinetobacter* species, with polymicrobial infections being common [8]. Bacterial infiltration causes tissue necrosis and gas formation in subcutaneous, fascial, and deep tissues [9]. In the present case the infection spread from an abscess in the right submandibular region to the peripharyngeal fascia and the external pterygoid muscle, and along the ramus to the right zygomatic area, involving the soft tissue of the temporoparietal region. In this case, thorough drainage of affected spaces was performed to prevent further tissue destruction and systemic complications. Additionally, the placement of corrugated rubber drains aids in continuous drainage and prevents re-accumulation of pus. Prompt and aggressive surgical debridement is essential to remove necrotic tissue and control the spread of infection. The use of broad-spectrum intravenous antibiotics is crucial to target the causative bacterial pathogens effectively [10]. Considering the poly-microbial nature of NF, a combination of antibiotics targeting both aerobic and anaerobic bacteria is often recommended. The initiation of antibiotics alongside surgical intervention helps to eradicate bacteria systemically and reduce the risk of systemic complications [11]. Early recognition and aggressive management are essential to mitigate adverse outcomes. Treatment involves aggressive surgical debridement alongside IV antibiotics, with some patients receiving hyperbaric oxygen therapy. Prompt recognition, extensive debridement, and broad-spectrum antibiotics are crucial for preventing severe morbidity and mortality [12].

Conclusions

Necrotizing fasciitis originating from odontogenic sources poses a grave threat, necessitating prompt intervention to avert severe complications. Given the significant mortality and morbidity associated with NFs, it's pertinent to consider the saying, "Surgery is often the most effective antibiotic". Further research is imperative to enhance diagnostic and therapeutic strategies for improved patient outcomes.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Parmarth Sonpal, Chetan Gupta, Nitin Bhola, Nidhi Bodiwala

Acquisition, analysis, or interpretation of data: Parmarth Sonpal, Chetan Gupta, Nitin Bhola, Nidhi Bodiwala

Drafting of the manuscript: Parmarth Sonpal, Chetan Gupta, Nitin Bhola, Nidhi Bodiwala

Critical review of the manuscript for important intellectual content: Parmarth Sonpal, Chetan Gupta, Nitin Bhola, Nidhi Bodiwala

Supervision: Parmarth Sonpal, Chetan Gupta, Nitin Bhola, Nidhi Bodiwala

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Roje Z, Roje Z, Matić D, Librenjak D, Dokuzović S, Varvodić J: Necrotizing fasciitis: literature review of contemporary strategies for diagnosing and management with three case reports: torso, abdominal wall, upper and lower limbs. *World J Emerg Surg.* 2011, 6:46. [10.1186/1749-7922-6-46](https://doi.org/10.1186/1749-7922-6-46)
2. Hohlweg-Majert B, Weyer N, Metzger MC, Schön R: Cervicofacial necrotizing fasciitis. *Diabetes Res Clin Pract.* 2006, 72:206-8. [10.1016/j.diabres.2005.09.012](https://doi.org/10.1016/j.diabres.2005.09.012)
3. Crew JR, Varilla R, Allandale Rocas Iii T, Abdul Rani S, Debabov D: Treatment of acute necrotizing fasciitis using negative pressure wound therapy and adjunctive NeutroPhase irrigation under the foam. *Wounds.* 2013, 25:272-7.
4. Brunworth J, Shibuya TY: Craniocervical necrotizing fasciitis resulting from dentoalveolar infection. *Oral Maxillofac Surg Clin North Am.* 2011, 23:425-32. [10.1016/j.coms.2011.04.007](https://doi.org/10.1016/j.coms.2011.04.007)

5. Shaariyah MM, Marina MB, Mohd Razif MY, Mazita A, Primuharsa Putra SH: Necrotizing fasciitis of the head and neck: surgical outcomes in three cases. *Malays J Med Sci.* 2010, 17:51-5.
6. Whitesides L, Cotto-Cumba C, Myers RA: Cervical necrotizing fasciitis of odontogenic origin: a case report and review of 12 cases. *J Oral Maxillofac Surg.* 2000, 58:144-51; discussion 152. [10.1016/s0278-2391\(00\)90327-6](https://doi.org/10.1016/s0278-2391(00)90327-6)
7. Gore MR: Odontogenic necrotizing fasciitis: a systematic review of the literature . *BMC Ear Nose Throat Disord.* 2018, 18:14. [10.1186/s12901-018-0059-y](https://doi.org/10.1186/s12901-018-0059-y)
8. Quereshy FA, Baskin J, Barbu AM, Zechel MA: Report of a case of cervicothoracic necrotizing fasciitis along with a current review of reported cases. *J Oral Maxillofac Surg.* 2009, 67:419-23. [10.1016/j.joms.2008.07.017](https://doi.org/10.1016/j.joms.2008.07.017)
9. Tung-Yiu W, Jehn-Shyun H, Ching-Hung C, Hung-An C: Cervical necrotizing fasciitis of odontogenic origin: a report of 11 cases. *J Oral Maxillofac Surg.* 2000, 58:1347-52; discussion 1353. [10.1053/joms.2000.18259](https://doi.org/10.1053/joms.2000.18259)
10. Agarwal L, Yasin A: Necrotising fasciitis: a fatal case of sepsis and a diagnostic challenge - case report and review of literature. *Int J Emerg Med.* 2018, 11:23. [10.1186/s12245-018-0183-x](https://doi.org/10.1186/s12245-018-0183-x)
11. Nazir H, Chieng CY, Rogers SN, Nekrasius R, Dodd M, Shah N: Outcomes of necrotizing fasciitis in the head and neck region in the United Kingdom-a case series and literature review. *Adv Oral Maxillofac Surg.* 2022, 6:100254. [10.1016/j.adoms.2022.100254](https://doi.org/10.1016/j.adoms.2022.100254)
12. Stevens DL, Bryant AE: Necrotizing soft-tissue infections. *N Engl J Med.* 2017, 377:2253-65. [10.1056/NEJMra1600673](https://doi.org/10.1056/NEJMra1600673)