

Effective “Pop-Out” Nucleus Technique in Cataract Surgery With Pseudoexfoliation: A Case Report

Review began 06/25/2026

Review ended 06/30/2026

Published 07/03/2026

© Copyright 2026

Sharaf. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

DOI: 10.7759/cureus.111999

Talal Sharaf¹

1. Ophthalmology, Emirates Hospital, Dubai, ARE

Corresponding author: Talal Sharaf, dr.talalsharaf@gmail.com

Abstract

Pseudoexfoliation syndrome (PXF) presents significant challenges in cataract surgery due to zonular weakness and poor pupillary dilation. We report a case of a 72-year-old woman with PXF who underwent cataract surgery using a “pop-out” nucleus technique facilitated by hydrodissection. The technique allowed the safe prolapse of the nucleus into the anterior chamber despite zonular instability. Postoperative visual outcomes were favorable, improving from 0.1 preoperatively to 0.9 one week after surgery. This case highlights the safety and effectiveness of the “pop-out” technique in managing complex cataract cases associated with pseudoexfoliation.

Categories: General Surgery, Internal Medicine, Ophthalmology

Keywords: capsular tension ring, cataract surgery, complex cataract, hydrodissection, nucleus prolapse technique, phacoemulsification, pseudoexfoliation syndrome, zonular weakness

Introduction

Pseudoexfoliation syndrome (PXF) is a systemic condition characterized by the accumulation of fibrillar extracellular material in ocular tissues, particularly on the lens capsule and zonules [1,2]. It is commonly associated with poor pupillary dilation, zonular weakness, and an increased risk of complications during cataract surgery [3].

Managing cataract in eyes with pseudoexfoliation presents significant surgical challenges, including zonular instability and a higher risk of capsular rupture [2,4]. Various surgical techniques have been described to improve safety and outcomes in such cases.

We present a case demonstrating the use of a “pop-out” nucleus technique facilitated by hydrodissection in a patient with pseudoexfoliation syndrome, highlighting its effectiveness and safety in the presence of zonular weakness.

Case Presentation

A 72-year-old woman presented with decreased vision in both eyes, more pronounced in the right eye. Preoperative best-corrected visual acuity (BCVA) in the right eye was approximately 0.1. Slit-lamp examination revealed features consistent with pseudoexfoliation syndrome, including the deposition of pseudoexfoliative material on the anterior lens capsule and poor pupillary dilation (Figure 1).

How to cite this article

Sharaf T (July 03, 2026) Effective “Pop-Out” Nucleus Technique in Cataract Surgery With Pseudoexfoliation: A Case Report. Cureus 18(7): e111999. DOI 10.7759/cureus.111999




FIGURE 1: Slit-lamp examination revealed pseudoexfoliative material on the anterior lens capsule.

Cataract surgery was planned for the right eye. Intraoperatively, zonular weakness with areas of zonular dialysis was noted. After successful capsulorhexis, hydrodissection was carefully performed. Due to zonular instability, a “pop-out” nucleus technique was utilized, whereby the nucleus was gently prolapsed into the anterior chamber using controlled hydrodissection while preserving the integrity of the posterior capsule.

The nucleus was then emulsified using phacoemulsification, followed by the removal of the epinucleus and cortical material via irrigation and aspiration. A capsular tension ring (CTR) was implanted to stabilize the capsular bag, and a posterior chamber intraocular lens (IOL) was successfully placed.

On the first postoperative day, visual acuity improved to 0.3 with mild corneal edema. At one-week follow-up, the patient achieved a BCVA of 0.9 with a stable intraocular lens and no complications.

Discussion

Pseudoexfoliation syndrome (PXF) is a well-recognized age-related systemic condition characterized by the progressive accumulation of fibrillar extracellular material in ocular tissues, particularly on the anterior lens capsule, zonules, and pupillary margin [1]. It is frequently associated with increased surgical complexity during cataract extraction due to zonular weakness, poor pupillary dilation, and a higher risk of intraoperative complications [2].

In the present case, the presence of pseudoexfoliative material on the anterior lens capsule was clearly identified on slit-lamp examination, consistent with classical findings described in the literature [1]. Such cases require careful preoperative planning and intraoperative modifications to ensure surgical safety and optimal outcomes.

Previous studies have consistently reported a higher incidence of intraoperative complications in eyes with pseudoexfoliation syndrome, particularly zonular dialysis, capsular rupture, and vitreous loss [2,4]. Shingleton et al. emphasized that zonular instability remains one of the most significant challenges in managing these cases, often necessitating modified surgical strategies [2].

The pop-out nucleus technique represents an effective alternative approach in managing cataracts associated with pseudoexfoliation. This method minimizes zonular stress by facilitating controlled nucleus prolapse into the anterior chamber, thereby reducing manipulation within the capsular bag. Compared to conventional phacoemulsification techniques, which rely heavily on in-the-bag nucleus disassembly, this approach may decrease mechanical stress on weakened zonules and lower the risk of intraoperative complications [3,5].

Furthermore, Shingleton et al. have highlighted the importance of adapting surgical techniques in eyes with compromised zonular support, recommending strategies that limit zonular traction and enhance intraoperative control [3]. In this context, the pop-out nucleus technique aligns well with these principles

and offers a safer surgical alternative in selected cases.

Additionally, the use of adjunctive measures such as capsular tension rings, careful hydrodissection, and viscoelastic support has been shown to improve surgical outcomes in pseudoexfoliation cases [5]. Surgeons should remain vigilant for signs of zonular weakness and be prepared to modify their surgical approach accordingly.

This case underscores the importance of recognizing pseudoexfoliation preoperatively and tailoring the surgical technique to minimize complications. The successful use of the pop-out nucleus technique in this setting demonstrates its value as a practical and effective strategy in managing complex cataract cases associated with pseudoexfoliation syndrome.

Conclusions

Pseudoexfoliation syndrome remains a significant risk factor for complications during cataract surgery due to zonular instability and poor pupillary dilation. Early recognition through careful slit-lamp examination is essential for appropriate surgical planning. The pop-out nucleus technique offers a safe and effective approach in such cases by minimizing zonular stress and reducing intraoperative risks. This case highlights the importance of adapting surgical strategies to the underlying pathology to achieve optimal outcomes.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Talal Sharaf

Acquisition, analysis, or interpretation of data: Talal Sharaf

Drafting of the manuscript: Talal Sharaf

Critical review of the manuscript for important intellectual content: Talal Sharaf

Supervision: Talal Sharaf

Disclosures

Human subjects: Informed consent for treatment and open access publication was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Ritch R, Schlötzer-Schrehardt U: Exfoliation syndrome. *Surv Ophthalmol*. 2001, 45:265-315. [10.1016/S0039-6257\(00\)00196-X](https://doi.org/10.1016/S0039-6257(00)00196-X)
2. Shingleton BJ, Crandall AS, Ahmed II: Pseudoexfoliation and the cataract surgeon: preoperative, intraoperative, and postoperative issues related to intraocular pressure, cataract, and intraocular lenses. *J Cataract Refract Surg*. 2009, 35:1101-20. [10.1016/j.jcrs.2009.03.011](https://doi.org/10.1016/j.jcrs.2009.03.011)
3. Shingleton BJ, Heltzer J, O'Donoghue MW: Outcomes of phacoemulsification in patients with and without pseudoexfoliation syndrome. *J Cataract Refract Surg*. 2003, 29:1080-6. [10.1016/S0886-3350\(02\)01993-4](https://doi.org/10.1016/S0886-3350(02)01993-4)
4. Schlötzer-Schrehardt U, Naumann GO: Ocular and systemic pseudoexfoliation syndrome. *Am J Ophthalmol*. 2006, 141:921-37. [10.1016/j.ajo.2006.01.047](https://doi.org/10.1016/j.ajo.2006.01.047)
5. Belovay GW, Varma DK, Ahmed II: Cataract surgery in pseudoexfoliation syndrome. *Curr Opin Ophthalmol*. 2010, 21:25-34. [10.1097/ICU.0b013e328332f814](https://doi.org/10.1097/ICU.0b013e328332f814)