

Evaluation of Salivary Alkaline Phosphatase Levels in Passive Smokers of Different Age Groups

Sayem A. Mulla¹, Aarti S. Bedia¹, Haritha K. Nimmagadda², Sumit Bedia³, Amit H. Patil⁴

Review began 06/25/2023

Review ended 06/30/2023

Published 07/03/2023

© Copyright 2023

Mulla et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Oral Medicine and Radiology, Bharati Vidyapeeth (Deemed to be University) Dental College and Hospital, Navi Mumbai, IND 2. Anatomy, Bharati Vidyapeeth (Deemed to be University) Dental College and Hospital, Navi Mumbai, IND 3. Prosthodontics, Bharati Vidyapeeth (Deemed to be University) Dental College and Hospital, Navi Mumbai, IND 4. Conservative Dentistry and Endodontics, Bharati Vidyapeeth (Deemed to be University) Dental College and Hospital, Navi Mumbai, IND

Corresponding author: Sayem A. Mulla, sayemmullaa@gmail.com

Abstract

Background

The smoke inhaled by a nonsmoker from the smoldering end of a cigarette is referred to as passive smoke. The nicotine present in smoke is known to cause tissue damage and alter the enzymatic composition of the body. Alkaline phosphatase (ALP) is a group of intracellular hydrolytic enzymes known to partake in cellular metabolism. ALP levels are affected by smoking as well as passive smoking (PS) with a change in the pH of the oral cavity. The association of salivary alkaline phosphatase (S-ALP) levels in different age groups, gender, and times of exposure is not thoroughly explored yet, which was the primary aim of this study.

Material and methods

A total of 64 samples were collected from passive smokers and non-smokers. Unstimulated saliva (2-2.5 mL) was collected from each subject after obtaining their consent, followed by centrifuging and mixing with ALP reagent in a semi-autoanalyzer to obtain the S-ALP levels.

Results

Higher S-ALP levels were seen in passive smokers (34.70 IU/L) compared to healthy individuals (12 IU/L), which came to be statistically significant ($p < 0.01$). S-ALP levels, when compared to different age groups and gender, were statistically insignificant ($p > 0.05$). However, higher levels were seen in association with time of exposure in passive smokers where the data was statistically significant ($p < 0.01$), suggesting tissue damage possibly due to oxidative stresses and tissue inflammation on continuous exposure for a minimum of 30-60 minutes daily as per our study.

Conclusion

Significantly high levels of S-ALP were found in passive smokers in comparison to non-smokers. This suggests that passive smoking has negative effects on the body tissues. Age, gender, and time of exposure of a non-smoker to tobacco smoke can lead to alterations in S-ALP levels. High levels of S-ALP were seen in individuals with prolonged exposure to tobacco smoke on a daily basis. Salivaomics can thus be used as a non-invasive, economical, and accurate alternative in tissue damage diagnosis.

Categories: Dentistry, Oral Medicine

Keywords: salivaomics, salivary alkaline phosphatase, tobacco smoke, secondhand smoke, passive smoking

Introduction

Passive smoke, also known as secondhand smoke or involuntary smoke, is considered to be an indoor pollutant. It is notorious for being a lethal cocktail of over 4,000 chemicals, 250 of which are said to be harmful with 25 of them being linked to cancers. It is the smoke inhaled by a non-smoker from the smoldering end of a cigarette. It is composed of aged exhaled mainstream smoke and diluted sidestream smoke [1]. It is known to contain a variety of carcinogenic chemical compounds such as carbon monoxide and nicotine.

Passive smoking (PS) is ill-reputed to cause a number of health hazards in both adults and children. It is infamous for primarily causing lung cancer in adults. Other forms of cancers such as cancer of the larynx and nasopharynx and lymphoma are also a major cause of concern. In recent times, it has been linked to respiratory disease symptoms. Children are more vulnerable to the effects of passive smoking due to the presence of more red blood cells and faster breathing rates. They are usually exposed to PS by adults (parents or others), which can cause lung infections (bronchitis and pneumonia), frequent ear infections, increased frequency of sickness, cough, wheezing, shortness of breath, and other symptoms. Other diseases or

How to cite this article

Mulla S A, Bedia A S, Nimmagadda H K, et al. (July 03, 2023) Evaluation of Salivary Alkaline Phosphatase Levels in Passive Smokers of Different Age Groups. Cureus 15(7): e41336. DOI 10.7759/cureus.41336

discomforts caused by passive smoking include odor annoyance, nasal irritation, and allergies [2].

Salivary biomarkers, which can be measured in both healthy and diseased people, are sentinel molecules that could be used to monitor health and disease surveillance [3]. Tobacco products contain a high concentration of toxic chemicals, carcinogens, and free radicals in both stable and unstable forms. This can cause local irritation and tissue damage, which can result in the production of biomarkers [4]. The concentrations and levels of these biomarkers can be determined using specialized, designated reagents run in autoanalyzer or semi-autoanalyzer machines to interpret their concentrations and levels in a specific sample of that population. Such processes can aid in the diagnosis of various diseases and disorders, as well as their consequences for human health.

Alkaline phosphatase (ALP) is an intracellular hydrolase enzyme that is involved in cell metabolism. This isoenzyme group is found on the cell membrane's outer layer. They assist in catalyzing the hydrolysis of organic phosphate esters found in extracellular space [5]. It has a crucial role in intracellular destructive processes and cellular damage. It has bicarbonate and phosphate ions, which help in buffering against acids [6]. It is usually measured in blood, urine, and saliva samples under the unit "IU/L," i.e., "international units per liter." A change in the levels of salivary alkaline phosphatase (S-ALP) can indicate a number of health issues. A rise in S-ALP levels denotes inflammation and healthy tissue destruction [7].

Although urine or serum samples are the most commonly used methods for ALP diagnosis, the use of saliva has some compelling arguments. When compared to serum or urine sample collection, salivary diagnosis is less invasive and faster. It is inexpensive and can produce results in as little as an hour. It includes no patient discomfort and easier sample handling for the healthcare professional [6].

S-ALP estimation can be used to identify tissue injury as an early biomarker. Identification of its levels in different gender and age groups will provide information about preventive measures that can be taken in passive smokers to avoid the effect of tobacco on tissue damage. The purpose of the present study was to compare S-ALP levels in passive smokers of various age groups, levels of exposure, and gender and compare it with that of healthy non-smokers.

Materials And Methods

Study design and setup

The present study was conducted in the outpatient department (OPD) of a private dental college and hospital situated in Navi Mumbai. Ethical clearance was obtained prior to the commencement of the study.

Sampling technique

The sample size was calculated using the formula for the finite population. After assuming a confidence level of 95%, margin of error of 5%, and a population proportion of 50%, we obtained 53 as the baseline to generalize the sample size. Therefore, a lower limit of 53 samples had to be selected for the study, and after rounding off as well as considering equal distribution between all the groups, 64 samples were included in the study.

Inclusion criteria

Patients with routine dental problems who came for treatment in dental OPD giving consent for participation in the study were included. Patients with passive smoking exposure for at least a year for the test group and participants who were not exposed to passive smoking ever were taken as the control group. Patients of all gender were included in the study.

Exclusion criteria

Patients with oral potentially malignant disorders and systemic illnesses such as diabetes, hypertension, thyroid or any known endocrine disorders, osteoarthritis, osteosarcoma, and renal failure were excluded. Patients who were diagnosed with periodontitis clinically based on probing depth, bleeding gums, and radiographic bone loss were also excluded from the study. Patients consuming alcohol or with any liver diseases were deferred. Patients with Paget's disease, hyperparathyroidism, osteomalacia, metastatic bone disease, and a recent fracture were barred. Patients with any of these conditions were exempted from the study because these conditions are associated with altered alkaline phosphatase.

Choice of subjects and control

There were 64 samples included. Patients who visited the dental OPD for routine dental checkups and treatment for various dental ailments were considered, and their consent was obtained for inclusion in the study. A brief history and demographic data were gathered. To assess the exposure to passive smoking, a pre-study questionnaire/patient assessment sheet (PAS) was used (Appendices).

Grouping of age

Group 1 consisted of participants below the age of 14 years (childhood). Group 2 has participants aged between 15 and 25 years (young age). Group 3 had participants aged between 26 and 60 years (adulthood). Group 4 consisted of participants above the age of 60 years (old age). After inclusion in the study, the subjects were separated into test groups (passive smokers) and control groups (non-passive smokers).

Procedure

Subjects were informed about the purpose of the study followed by obtaining informed consent. They were instructed to fast for two hours prior to sample collection as it reduces alterations in salivary enzyme levels. The subjects were asked to collect the saliva on the floor of the oral cavity and spit it in the sample collection tube. A minimum of 2-2.5 mL of unstimulated saliva was collected from each subject. The samples were then centrifuged at 3,000 rotations per minute (rpm) for 10 minutes. The resultant 20 µL of supernatant was then mixed with 1,000 µL of ERBA Mannheim ALP reagent (Mannheim, Germany) and run in a semi-autoanalyzer. This was based on the kinetic method recommended by the International Federation of Clinical Chemistry. The reading obtained on the screen (in IU/L) was then noted and subjected to statistical tests.

Statistical tests used

Data obtained were compiled on a Microsoft Office Excel sheet version 2019 (Microsoft Corp., Redmond, WA, USA). Data were subjected to statistical analysis using the Statistical Package for the Social Sciences version 26.0 (IBM SPSS Statistics, Armonk, NY, USA). Frequencies and percentages were calculated for age and gender distribution. An intergroup comparison (between test and control) (two groups) was done using Student’s t-test. An inter-subgroup comparison was done using one-way analysis of variance (ANOVA). For all the statistical tests, p<0.05 was considered to be statistically significant, keeping α error at 5% and β error at 20%, thus giving a study power of 80%.

Results

Demographic data

A total of 64 saliva samples were collected. Out of this, 32 were test (passive smokers) and 32 were control (healthy) samples. A total of 26 (13 test and 13 control) female participants’ saliva samples were included, and 38 (19 test and 19 control) male participants’ saliva samples were collected.

Comparison of S-ALP values in the test group and control group

An intergroup comparison of S-ALP values between the two groups (test and control) was done. There was a statistically highly significant difference seen between the S-ALP values between the test and control group (p<0.01) with higher values in the test group (Table 1). An inter-subgroup comparison was done to compare S-ALP values with age groups, gender, and time of exposure.

S-ALP value	Number	Mean (IU/L)	Standard deviation (IU/L)	Standard error mean	t value	p value of t-test
Test	32	34.7213	11.80244	2.08640	9.931	0.000**
Control	32	12.0456	5.24742	0.92762		

TABLE 1: Intergroup comparison of S-ALP values

**Statistically significant (p<0.01)

S-ALP: salivary alkaline phosphatase

Comparison of S-ALP values with age in the test group

There was a statistically non-significant difference seen for the S-ALP values in comparison with age groups in the test group (p>0.05) (Table 2).

Age groups	Number	Mean (IU/L)	Standard deviation (IU/L)	Standard error	95% confidence interval for mean		Minimum	Maximum	F value	p value of one-way ANOVA
					Lower bound	Upper bound				
Below 14 years of age	8	34.2125	11.64189	4.11603	24.4796	43.9454	25.61	58.65	0.053	0.984#
15-25 years of age	8	34.4538	14.34105	5.07033	22.4643	46.4432	22.00	55.10		
26-60 years of age	8	36.2100	11.08853	3.92039	26.9398	45.4802	22.60	54.00		
Above 60 years of age	8	34.0088	12.21654	4.31920	23.7955	44.2220	21.60	52.98		
Total	32	34.7213	11.80244	2.08640	30.4660	38.9765	21.60	58.65		

TABLE 2: Comparison of S-ALP values with age groups in the test group

#Statistically non-significant (p>0.05)

S-ALP: salivary alkaline phosphatase, ANOVA: analysis of variance

Comparison of S-ALP values with gender in the test group

While comparing S-ALP values of different gender in the test group, no statistically significant difference was found (p>0.05) (Table 3).

	Gender	Number	Mean (IU/L)	Standard deviation (IU/L)	Standard error	t value	p value of t-test
S-ALP value	Male	19	32.9053	11.47846	2.63334	-1.054	0.300#
	Female	13	37.3754	12.22156	3.38965		

TABLE 3: Comparison of S-ALP values with gender in the test group

#Statistically non-significant (p>0.05)

S-ALP: salivary alkaline phosphatase

Comparison of S-ALP values with time of exposure in the test group

On comparing the S-ALP values with time of exposure in the test group, we found a statistically highly significant difference (p<0.001) with higher values in individuals who were exposed to smoke for at least 30-60 minutes daily (Table 4).

	Time of exposure	Number	Mean (IU/L)	Standard deviation (IU/L)	Standard error	t value	p value of t-test
S-ALP value	<30 minutes	27	31.8515	10.34790	1.99145	-3.840	0.001**
	30-60 minutes	05	50.2180	5.26937	2.35653		

TABLE 4: Comparison of S-ALP values with time of exposure in the test group

**Statistically significant (p<0.01)

S-ALP: salivary alkaline phosphatase

Discussion

Saliva has been extensively researched as a potential diagnostic tool over the last decade due to its ease of use and non-invasive accessibility, as well as its abundance of biomarkers such as genetic material and proteins [8]. It has the potential to enable chairside diagnosis of a variety of oral and systemic diseases [9]. It

contains a diverse range of hormones, antibodies, growth factors, enzymes, microbes, and their products [10,11], allowing it to function as a promising and accurate diagnostic tool. Many of these constituents enter saliva via blood through passive diffusion, active transport, or extracellular ultrafiltration [10,12], making it a reflection of bodily physiological functioning [13]. It is a non-invasive technique that usually has better patient cooperation and is cost-effective and nearly as accurate as blood investigations.

As stated earlier, alkaline phosphatase is a group of intracellular hydrolytic enzymes participating in cellular metabolism [5]. It is mainly found in the liver and bones with trace amounts present in the kidneys, intestines, placenta, and leukocytes. The enzymatic group of ALP creates an alkaline environment that catalyzes the hydrolysis of phosphate esters. It also aids bone calcification metabolism as well as intestinal lipid transport, although its exact metabolic role is yet to be established [6,14]. The source of this enzyme in the oral cavity includes neutrophils, bacteria, and oral epithelial cells [15,16]. An increase in ALP levels is frequently linked to a number of diseases and conditions, including dental caries, periodontitis, malignant and potentially malignant disorders, smoking, and some bone diseases such as Paget's disease. Patients with untreated celiac disease also show elevated levels. Low ALP levels are seen in patients with hypophosphatasia, postmenopausal women receiving estrogen therapy for osteoporosis, men who have recently had heart surgery, malnutrition, magnesium deficiency, hypothyroidism, severe anemia, children with achondroplasias, and cretinism [17].

The present study was conducted to find out S-ALP values in passive smokers and correlate them to different age groups, gender, and times of exposure. The mean values of S-ALP in the control group, i.e., healthy individuals, were found to be around 12 IU/L, which was similar to that obtained by Sushmitha et al. [18]. The mean S-ALP values obtained in the test group, i.e., passive smokers, were found to be 34.70 IU/L.

Studies done by Ibraheem et al. [19] and Hassan [20] showed that there was a rise in S-ALP levels in passive smokers in comparison to healthy individuals; however, the difference was statistically insignificant. In contrast to this, the present study's results found that S-ALP levels were high in passive smokers when compared to healthy individuals and were statistically significant ($p < 0.01$).

The present study's objective was to evaluate the difference in S-ALP values in regard to age, gender, and time of exposure in individuals. We found no difference in S-ALP values in different age groups or gender. However, there was a statistically significant ($p < 0.01$) difference seen in regard to time of exposure. S-ALP values were high in individuals who were exposed to smoke daily for around 30-60 minutes. Continuous exposure to smoke causes deleterious effects and leads to alteration in membrane-bound ALP levels. The effect of smoking or passive smoking on ALP may be due to an altered imbalance between the free oxygen radicals and the imbalance in the antioxidant levels [6].

Howard et al. [21], Howard et al. [22], and Chen et al. [23] have highlighted that passive smoking and active smoking have systemic effects in the same direction, although the magnitude may not be the same.

The use of saliva to detect the effect of passive smoking on health can help expand the literature on the utility of salivary alkaline phosphatase as a biomarker in assessing passive smoking effects. The current study's results can indicate the effect of passive smoking in different age groups and gender, although the insignificant results may indicate the need for a larger sample size and more time to compare the results. Further studies with a larger sample size can help us understand the impact of passive smoking on tissue health more clearly.

Future recommendations

Future studies with larger sample sizes should be done to better understand the effects of passive smoking on tissue health. There is literature stating that passive smoking can lead to an increased risk of low-birth-weight pregnancy outcomes, and there is an increase in ALP levels in pregnant females exposed to smoke [24,25]. Studies can be done concentrating on S-ALP levels in such individuals, which would positively and effectively contribute to the literature as well as public health.

Conclusions

The present study pointed out that there is a significant difference ($p < 0.01$) between the S-ALP values of passive smokers and healthy individuals. This suggests that passive smoking has negative effects on body tissues. However, there exists no traceable difference ($p > 0.05$) in the S-ALP values of different age groups and gender of passive smokers, but, to highlight, there exists a significant difference ($p < 0.01$) in the S-ALP values in passive smokers based on their time of exposure to the smoke. This shows that continuous exposure to smoke may have harmful effects on healthy tissues, leading to an alteration in ALP levels. The study also showed that the S-ALP enzyme can act as a major non-invasive, accurate, and economical alternative in salivaomics.

Appendices

Figure 1 shows the pre-study questionnaire/patient assessment sheet (PAS) used in the present study.

<ol style="list-style-type: none"> 1. E-Mail I.D. _____ 2. Name _____ 3. Date of Birth _____ 4. Which age group do you fall in? <ul style="list-style-type: none"> <input type="radio"/> Below 14 years <input type="radio"/> 15-25 years <input type="radio"/> 26-40 years <input type="radio"/> Above 60 years 5. Gender. <ul style="list-style-type: none"> <input type="radio"/> Male <input type="radio"/> Female <input type="radio"/> Prefer not to say 6. Address _____ 7. Where is your locality based in geographically? <ul style="list-style-type: none"> <input type="radio"/> Urban <input type="radio"/> Semi urban <input type="radio"/> Rural <input type="radio"/> Semi rural 8. What is the highest level of education you have completed? <ul style="list-style-type: none"> <input type="radio"/> Less than higher school <input type="radio"/> Higher secondary <input type="radio"/> Diploma <input type="radio"/> Undergraduation <input type="radio"/> Postgraduation <input type="radio"/> Not educated 9. Do other people smoke cigarette in your house? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No 10. How many individuals in your surrounding smoke on an average? <ul style="list-style-type: none"> <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> More than 5 11. How many times are the cigarettes smoked in your surroundings per day? <ul style="list-style-type: none"> <input type="radio"/> >20 cigarettes per day <input type="radio"/> 10-20 cigarettes per day <input type="radio"/> 6-5 cigarettes per day 	<ul style="list-style-type: none"> <input type="radio"/> 1-5 cigarettes per day <input type="radio"/> None <ol style="list-style-type: none"> 12. What is the extent of time you/your child is exposed to passive smoking on an average per day? <ul style="list-style-type: none"> <input type="radio"/> Below 30 min/day <input type="radio"/> 30-60 min/day <input type="radio"/> 60-120 min/day <input type="radio"/> More than 120 min/day 13. Are you aware of the passive smoking concept? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe 14. Do you agree that passive smoking is hazardous to your child as well as the people around you? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe 15. Do you think your child/you have been exposed to passive smoking? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe 16. Is smoking allowed/practiced in your household/close surroundings? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe 17. Is it possible that women around you/yourself had been exposed to passive smoking during pregnancy? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe <input type="radio"/> Not applicable 18. Have you ever offered help to someone to stop smoking for other individuals? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe 19. Would you be interested in helping individuals for stopping smoking? <ul style="list-style-type: none"> <input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Maybe
--	--

FIGURE 1: Patient assessment sheet

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. The Biomedical Ethics Committee of Bharati Vidyapeeth (Deemed to be University) Dental College and Hospital, Navi Mumbai, issued approval IEC343072022 version number 001 dated 4.07.2022. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** The present study was selected and approved by the Indian Council of Medical Research (ICMR) for its annual Short Term Studentship 2022 program (STS 2022). Thus, the project was funded under the ICMR-STs 2022 program (2022-05239). **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

We would like to thank Mrs. Swati R. Dhande, Head, Department of Pharmacology, Bharati Vidyapeeth's College of Pharmacy, Navi Mumbai, for helping us in the processing of the samples.

References

1. Davey Smith G: Effect of passive smoking on health . BMJ. 2003, 326:1048-9. [10.1136/bmj.326.7398.1048](https://doi.org/10.1136/bmj.326.7398.1048)
2. Saulyte J, Regueira C, Montes-Martínez A, Khudyakov P, Takkouche B: Active or passive exposure to tobacco smoking and allergic rhinitis, allergic dermatitis, and food allergy in adults and children: a systematic review and meta-analysis. PLoS Med. 2014, 11:e1001611. [10.1371/journal.pmed.1001611](https://doi.org/10.1371/journal.pmed.1001611)
3. Zhang L, Xiao H, Wong DT: Salivary biomarkers for clinical applications . Mol Diagn Ther. 2009, 13:245-59. [10.1007/BF03256330](https://doi.org/10.1007/BF03256330)
4. Sivapathasundharam B: Textbook of oral embryology and histology . Jaypee Brothers Medical Publishers, New Delhi, India; 2018.
5. Lowe D, Sanvictores T, Zubair M, John S: Alkaline phosphatase. StatPearls Publishing, Treasure Island, FL; 2023.
6. Prakash AR, Nahar P, Ashtekar M, Natarajan S, Singh R, Kulkarni G: Detection of salivary alkaline phosphatase levels in smokers, diabetic patients, potentially malignant diseases and oral malignant tumours. J Pharm Bioallied Sci. 2020, 12:S430-5. [10.4103/jpbs.JPBS.129.20](https://doi.org/10.4103/jpbs.JPBS.129.20)
7. Prakash AR, Indupuru K, Sreenath G, Kanth MR, Reddy AV, Indira Y: Salivary alkaline phosphatase levels speak about association of smoking, diabetes and potentially malignant diseases???. J Oral Maxillofac Pathol. 2016, 20:66-70. [10.4103/0973-029X.180934](https://doi.org/10.4103/0973-029X.180934)
8. Javaid MA, Ahmed AS, Durand R, Tran SD: Saliva as a diagnostic tool for oral and systemic diseases . J Oral Biol Craniofac Res. 2016, 6:66-75. [10.1016/j.jobcr.2015.08.006](https://doi.org/10.1016/j.jobcr.2015.08.006)
9. Wong DT: Salivaomics. J Am Dent Assoc. 2012, 143:19S-24S. [10.14219/jada.archive.2012.0339](https://doi.org/10.14219/jada.archive.2012.0339)
10. Pfaffe T, Cooper-White J, Beyerlein P, Kostner K, Punyadeera C: Diagnostic potential of saliva: current state

- and future applications. *Clin Chem*. 2011, 57:675-87. [10.1373/clinchem.2010.153767](https://doi.org/10.1373/clinchem.2010.153767)
11. Gröschl M: The physiological role of hormones in saliva . *Bioessays*. 2009, 31:843-52. [10.1002/bies.200900013](https://doi.org/10.1002/bies.200900013)
 12. Haeckel R, Hänecke P: The application of saliva, sweat and tear fluid for diagnostic purposes . *Ann Biol Clin (Paris)*. 1993, 51:903-10.
 13. Lima DP, Diniz DG, Moimaz SA, Sumida DH, Okamoto AC: Saliva: reflection of the body . *Int J Infect Dis*. 2010, 14:e184-8. [10.1016/j.ijid.2009.04.022](https://doi.org/10.1016/j.ijid.2009.04.022)
 14. Yoga S, Rangdhol V, Maithilikarpagaselvi N, Vandana S, Paulose S: Estimation of salivary sialic acid levels as a prognostic marker in patients with oral leukoplakia—a case control study. *J Adv Clin Res Insights*. 2017, 4:96-9. [10.15713/ins.jcri.168](https://doi.org/10.15713/ins.jcri.168)
 15. Merza KS, Alaaraji SB, Abdullah BH: Comparative study on lactate dehydrogenase, alkaline phosphatase and immunoglobulins in serum and saliva of acute leukemia and oral squamous cell carcinoma patients. *Iraqi J Sci*. 2010, 51:262-70.
 16. Sanikop S, Patil S, Agrawal P: Gingival crevicular fluid alkaline phosphatase as a potential diagnostic marker of periodontal disease. *J Indian Soc Periodontol*. 2012, 16:513-8. [10.4103/0972-124X.106889](https://doi.org/10.4103/0972-124X.106889)
 17. Prakash AR, Nahar P, Ashtekar M: Serum and salivary alkaline phosphatase levels in oral health and disease: a brief review. *IP Int J Maxillofac Imaging*. 2020, 6:53-5. [10.18231/j.ijmi.2020.016](https://doi.org/10.18231/j.ijmi.2020.016)
 18. Sushmitha M, Preethi M, Sivapathasundaram B: Estimation of salivary alkaline phosphatase levels in smokers and tobacco chewers associated with diabetes. *Eur J Mol Clini Med*. 2020, 7:119-28. [10.31838/ejmcm.07.05.12](https://doi.org/10.31838/ejmcm.07.05.12)
 19. Ibraheem L, Karem B, Dhafer AM: Effect of passive smoking on the periodontal health status and salivary enzymes level: a comparative study. *Adv Res J Multidiscip Discov*. 2018, 32:47-51.
 20. Hassan BK: The effect of smoking and passive smoking on periodontal health status and salivary enzymes level. University of Baghdad, Baghdad, Iraq; 2013.
 21. Howard G, Wagenknecht LE, Burke GL, et al.: Cigarette smoking and progression of atherosclerosis: the Atherosclerosis Risk in Communities (ARIC) study. *JAMA*. 1998, 279:119-24. [10.1001/jama.279.2.119](https://doi.org/10.1001/jama.279.2.119)
 22. Howard G, Wagenknecht LE, Cai J, Cooper L, Kraut MA, Toole JF: Cigarette smoking and other risk factors for silent cerebral infarction in the general population. *Stroke*. 1998, 29:913-7. [10.1161/01.str.29.5.913](https://doi.org/10.1161/01.str.29.5.913)
 23. Chen LC, Quan C, Hwang JS, et al.: Atherosclerosis lesion progression during inhalation exposure to environmental tobacco smoke: a comparison to concentrated ambient air fine particles exposure. *Inhal Toxicol*. 2010, 22:449-59. [10.3109/08958370903373845](https://doi.org/10.3109/08958370903373845)
 24. Hofhuis W, Merkus PJ, de Jongste JC: [Negative effects of passive smoking on the (unborn) child] . *Ned Tijdschr Geneesk*. 2002, 146:356-9.
 25. Banihosseini SZ, Baheiraei A, Shirzad N, Heshmat R, Mohsenifar A: The effect of cigarette smoke exposure on vitamin D level and biochemical parameters of mothers and neonates. *J Diabetes Metab Disord*. 2013, 12:19. [10.1186/2251-6581-12-19](https://doi.org/10.1186/2251-6581-12-19)