

Awareness and Practice of Najran University Students Towards Common Problems Related to Ear, Nose, and Throat

Review began 11/21/2023

Review ended 12/17/2023

Published 12/27/2023

© Copyright 2023

Assiri et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abdullah M. Assiri ¹, Asaiel AL Hadi ², Saleh Y. Al Hareth ², Meshal Alwabel ³, Enas M. Al Hadi ², Khalid M. Alkhalifah ³

1. Surgery, College of Medicine, Najran University, Najran, SAU 2. College of Medicine, Najran University, Najran, SAU 3. Unaizah College of Medicine and Medical Sciences, Qassim University, Unaizah, SAU

Corresponding author: Abdullah M. Assiri, al-hayani@hotmail.com

Abstract

Objective: The objective of this study was to evaluate the awareness and practice of Najran University students toward common problems related to the ear, nose, and throat.

Methods: A cross-sectional study was utilized in this research, employing data from a sample of 429 students at Najran University. The participants completed a self-administered questionnaire and ensured anonymity. The questionnaire used in this study had been previously validated.

Results: The sample for the current study primarily consisted of students aged more than 20 years (84.1%; n = 361), with a predominance of females (69.0%; n = 296). The majority of them were in health colleges (45.2%; n=194). The study results show that 37.8% (n = 162) had a good knowledge level, while 62.2% (n = 267) had poor knowledge about problems related to ENT. The vast majority, 87.2% (n = 374), believed that going to the hospital was the appropriate action to take in cases of acute ENT problems. The results established a statistically significant association between age, gender, health college, college, and department with p-values <0.005 (0.002*, 0.001*, 0.003*, and 0.005*), respectively, and the level of knowledge about problems related to ENT. There was no statistically significant association between nationality, clinical history of the participants, and the level of knowledge about problems related to ENT (p > 0.005).

Conclusion: The study revealed that 37.8% of the participants had good knowledge about problems related to the ears, nose, and throat. The participants older than 20 years had better knowledge of common ENT problems than those younger than 20 years. Female participants showed a higher level of knowledge and awareness of problems related to the ears, nose, and throat. The study noted that the participants in Health College, the faculty of medicine, and the third academic level had good knowledge about problems related to ENT. The study established that going to the hospital was the appropriate action to be taken in case of a sudden ENT problem. Therefore, we recommend concerted efforts be made among the medical community to increase knowledge about common ENT problems.

Categories: Epidemiology/Public Health, Otolaryngology

Keywords: university student, knowledge, public health and safety, orl-hns, ent

Introduction

Otorhinolaryngology (ORL), also known as ear, nose, and throat (ENT) medicine, is a specialized branch of medicine focused on the diagnosis and treatment of disorders related to the ear, nose, and throat, as well as the head and neck. The importance of ORL knowledge and awareness among the general population cannot be overstated, as many common ENT conditions can significantly impact an individual's quality of life and may lead to more severe complications if not addressed on time [1,2]. However, diseases of the ENT are of extreme importance due to the morbidities that can result in physiological function impairment. The aforementioned difficulties encompass impairments in flavor, olfaction, speech, respiration, swallowing, phonation, protection of the lower respiratory tract, hearing, and secretion clearance [3,4].

Among the leading reasons individuals consult primary health care physicians around the world are symptoms associated with ENT [5,6]. Based on the WHO, approximately 278 million people globally suffer from bilateral hearing loss ranging from moderate to severe [7]. It was reported that 16.1% of adults in the United States (US) had encountered hearing loss during the period from 2003 to 2004 [8]. In 2004, an estimation placed the prevalence of rhinosinusitis among nasal diseases among the adult population of the United States at 16% [9].

In Saudi Arabia, there is a growing concern about the level of knowledge and awareness of ORL-related issues among the younger population, specifically university students. This concern arises from the

How to cite this article

Assiri A M, AL Hadi A, Al Hareth S Y, et al. (December 27, 2023) Awareness and Practice of Najran University Students Towards Common Problems Related to Ear, Nose, and Throat. Cureus 15(12): e51178. DOI 10.7759/cureus.51178

increasing prevalence of ENT problems in the country, with sinusitis, otitis media, and tonsillitis frequently reported among the population [10]. Early recognition and appropriate management of these conditions are essential to prevent complications and improve the overall health of affected individuals [11]. The reports of local knowledge and attitudes about ENT-related issues were insufficient. In a previous study, it was determined that 2.3% of individuals residing in the city of Riyadh showed exceptional knowledge, while 18.4% demonstrated a high level of knowledge. The majority of individuals (79.4%) have limited knowledge of ENT-related issues [12]. Another study was among school and university students in Makkah City, which showed a significant proportion of participants (42.22%) showed a moderate comprehension of ENT issues, while 41.48% showed a low level of understanding. On the other hand, only 16.30% of the respondents displayed a good degree of knowledge in this domain [10].

Najran City, located in the south of Saudi Arabia, has a substantial number of university students who may be at risk for ENT-related issues due to various factors, such as environmental conditions, lifestyle habits, and limited access to specialized healthcare services in the region [13]. However, to date, there is limited information available about the knowledge and awareness of common ORL-related issues among Najran or university students.

To address this knowledge gap, this cross-sectional study aims to assess the level of knowledge and awareness of common ORL-related issues among university students in Najran City, Saudi Arabia. This study's findings will help identify gaps in ORL knowledge and awareness, inform the development of targeted educational interventions, and contribute to the improvement of ENT health outcomes in the region.

Materials And Methods

Study design and population

This study employs a cross-sectional design to evaluate the knowledge and awareness of common otorhinolaryngology-related issues among students at Najran University in Saudi Arabia. The target population consists of university students from different faculties and academic levels.

Sample size and sampling technique

After determining the required sample size using a standard formula for cross-sectional studies, it was 376. And we employ a convenience sampling technique to select participants based on their accessibility and willingness to participate from various faculties and academic levels at Najran University. Although this method is easy to implement, it may result in a biased sample, limiting the generalizability of the findings.

Data collection tool

The study uses a structured, self-administered questionnaire to collect data from the participants. After permission, we used the questionnaire used by Jaladdin et al. (2023) [10]. The questionnaire was divided into four sections. The first section was to explain the goal of our study and the consent of the participant. The second section was for demographic information about the participants. The third section was for the clinical history, with four questions. The fourth section contains a series of multiple-choice questions with three answer options, except for the last two: "The appropriate action in case of hearing loss" had four answers and "Usually get information and knowledge about ENT issues" had six answers. The questionnaire was developed by board-certified otorhinolaryngologists. However, before conducting this survey, we performed content validity testing to ensure that the questions covered all aspects of the intended construct or topic. After validity, a reliability test was done using Cronbach's alpha. The result indicated very good internal consistency and reliably measured the practice and level of knowledge of ear, nose, and throat-related problems.

Data analysis

After collecting the questionnaires, researchers enter the data into a statistical software program for analysis. They use descriptive statistics, such as frequencies, percentages, means, and standard deviations, to summarize the demographic information and responses related to knowledge and awareness of otorhinolaryngology issues. They also employ inferential statistics, including chi-square tests and t-tests, to examine associations between demographic variables and awareness levels.

Ethical considerations

The study adheres to ethical considerations, such as obtaining ethical approval from the Committee of Scientific Research and Conferences at Najran University, which issued approval No. 4/3/2023, obtaining informed consent from participants, ensuring confidentiality and anonymity, and respecting participants' rights to withdraw from the study at any time.

Results

Table 1 shows that a total of 429 participants completed the questionnaire. The vast majority (84.1%, (n=361) of participants were more than 20 years old, with more than half (69.0%, (n=296) being female. 45.2% (n=194) of the students who participated in the survey studied at Health College, with 22.1% (n=95) in the faculty of medicine and 8.6% (n=37) in the faculty of nursing. The majority (16.8%; (n=72) of the participants were in the third academic level, with the vast majority (98.8%; (n=424) being Saudi nationals.

Demographic information	Category	Frequency and Proportion n (%)
Age	Less than 20 years	68 (15.9%)
	More than 20 years	361 (84.1%)
Gender	Male	133 (31.0%)
	Female	296 (69.0%)
Health college	Yes	194 (45.2%)
	No	235 (54.8%)
College and department	Faculty of medicine	95 (22.1%)
	College of nursing	37 (8.6%)
	College of applied medical science	29 (6.8%)
	Faculty of administration science	44 (10.3%)
	Faculty of science and literature	37 (8.6%)
	Others	187 (43.6%)
Academic level	Level one	55 (12.8%)
	Second Level	17 (4.0%)
	The third level	72 (16.8%)
	Fourth level	44 (10.3%)
	Level five	51 (11.9%)
	Sixth level	34 (7.9%)
	Seventh level	41 (9.6%)
	Eighth level	27(6.3%)
	Ninth	18 (4.2%)
	Tenth level	17 (4.0%)
	Eleventh level	11 (2.6%)
	Twelfth level	12 (2.8%)
	Thirteenth level	9 (2.1%)
	Fourteenth level	21 (4.9%)
Nationality	Saudi	424 (98.8%)
	Non-Saudi	5 (1.2%)

TABLE 1: Demographic information of the participants (N=429)

Demographic information is presented in frequencies (n) and proportions (%)

Table 2 (below) depicts the clinical history of the participants. The findings revealed that 22.1% (n=95) reported a previous diagnosis of ear problems, 25.4% (n=109) reported a previous diagnosis of nose problems, and 22.1% (n=95) reported a previous diagnosis of throat problems. Moreover, 14.2% (n=61) reported having undergone ear, nose, and throat surgery.

Clinical history	Categories	Frequency and Proportion n (%)
Previously diagnosed with ear problems	Yes	95 (22.1%)
	No	334 (77.9%)
Previously diagnosed with nose problems	Yes	109 (25.4%)
	No	320 (74.6%)
Previously diagnosed with throat problems	Yes	95 (22.1%)
	No	334 (77.9%)
Previous surgery concerning ear, nose, throat	Yes	61 (14.2%)
	No	368 (87.8%)

TABLE 2: Clinical history of the participants.

The clinical history of participants is presented in frequencies (n) and proportions (%)

Table 3 (below) depicts the knowledge about problems related to the ear, nose, and throat of participants. The vast majority, 61.1% (n=262), of the participants provided correct answers regarding the safety of using cotton swabs for ear cleaning. 61.8% (n=265) were aware that it is recommended to get flu vaccines annually as a preventive measure. More than half (62.2%, n=267) of the participants provided the correct answers on the importance of using vitamin C to treat and prevent influenza, and 77.6% (n=333) provided the correct definition of vertigo. Vast majority 80.4% (n=345) of the participants were aware that hearing loss can affect their social lives, with more than half (n=253) of the participants aware that constant exposure to loud and boisterous noises or sounds may cause hearing loss. 75.0% (n=324) of the participants had the knowledge that some infections of the inner ear and middle ear cause dizziness. About 43.6% (n=187) of the participants provided the correct answer to the question about handling nose bleeding. However, fewer than 50% of the respondents provided the correct answer to the question about the risk of developing obesity and immunodeficiency after tonsillectomy (29.1%, 42.2%). The vast majority (79%; n=339) of the participants were aware that constant misuse of voice causes vocal cord disorders.

Knowledge about problems related to ENT	Categories	Frequency and Proportion n (%)
Cotton swabs are an unsafe way to clean the ears?	Yes	262 (61.1%)
	No	92(21.4%)
	I don't know	75(17.4%)
Antibiotics are not considered to treat upper respiratory infections (such as colds or coughs).	Yes	151(35.1%)
	No	116(27%)
	I don't know	162 (37.8%)
It is recommended to get the flu vaccine annually as a preventive measure	Yes	265 (61.8%)
	No	74(17.2%)
	I don't know	90 (21.0%)
It is recommended for patients with diabetes and high blood pressure to take the seasonal flu vaccine.	Yes	215 (50.1%)
	No	36 (8.4%)
	I don't know	178(41.5%)
	Yes	267(62.2%)
	No	

Taking vitamin C treats and prevents influenza	No	67(15.6%)
	I don't know	95(22.1%)
Vertigo is another term for dizziness.	Yes	333 (77.6%)
	No	42 (9.8%)
	I don't know	54(12.6%)
	Yes	345(80.4%)
Hearing loss can affect the social life of the individual.	No	44 (10.3%)
	I don't know	40 (9.3%)
	Yes	253(59.0%)
	No	75 (17.4%)
Constant exposure to loud and annoying noises or sounds may cause hearing loss.	I don't know	101 (23.5%)
	Yes	324 (75.5%)
Some infections of the inner ear and middle ear cause dizziness.	No	25(5.8%)
	I don't know	80 (18.6%)
	Yes	138(32.1%)
	No	187(43.6%)
The correct way to deal with a nosebleed is to put the head back.	I don't know	104 (24.2%)
	Yes	90 (21.0%)
The use of nasal congestion drops is considered safe for long-term use	No	135 (31.5%)
	I don't know	204 (47.6%)
	Yes	181 (42.2%)
	No	90 (21.0%)
Tonsillectomy may cause immunodeficiency	I don't know	158 (36.8%)
	Yes	339 (79%)
Does constant screaming cause vocal cord disorders?	No	20 (4.7%)
	I don't know	70 (16.3%)

TABLE 3: Questions on knowledge about problems related to the ear, nose, and throat.

Questions on knowledge about problems related to ENT are presented in frequencies (n) and proportions (%).

Figure 1 shows the action that respondents should take in case they develop sudden hearing loss or earache. It is evident that most participants would visit an ENT specialist (87.2%), while a significantly smaller proportion would consult a general practitioner (7.2%).




Figure 2 below shows the participants' sources of information about ENT issues. It is evident that most respondents acquired the information from doctors (49.2%) and social media (18.4%).




Table 4 (below) depicts the relationship between participants' demographics, such as age, gender, health college, college department, nationality, previous clinical history, and level of knowledge about problems related to ENT. The results established a statistically significant association between age, gender, health college, college, and department with p-values<0.005 (0.002*, 0.001*, 0.003, and 0.005*), respectively, and the level of knowledge about problems related to ENT. There was no statistically significant association between nationality, clinical history of the participants, and the level of knowledge about problems related to ENT (p>0.005). The study results show that 37.8% (n=162) had a good knowledge level, while 62.2% (n=267) had poor knowledge about problems related to ENT.

Variables	Level of knowledge			
	Category	Poor	Good	p-value
Age	More than 20 years	281 (65.6%)	148 (34.4%)	0.002*
	More than 20 years	275 (64.1%)	154 (35.9%)	
Gender	Male	287 (67.0%)	142 (33.0%)	0.001*
	Female	283 (65.9%)	146 (34.1%)	
Health college	Yes	208 (48.6%)	221 (51.4%)	0.003*
	No	306 (71.4%)	123 (28.6%)	
College and department	Faculty of medicine	121 (28.2%)	308 (71.8%)	0.005*
	College of nursing	160 (37.3%)	296 (62.7%)	
	College of applied medical science	169 (39.5%)	260 (60.5%)	
	Faculty of administration science	288 (67.1%)	141 (32.9%)	
	Faculty of science and literature	284 (66.1%)	145 (33.9%)	
Nationality	Others	323 (75.2%)	106 (24.8%)	0.341
	Saudi	312 (72.8%)	117 (27.2%)	
	Non-Saudi	332 (77.4%)	97 (22.6%)	
Previously diagnosed with ear problems	Yes	256 (59.7%)	173 (40.3%)	0.155
	No	288 (67.1%)	141 (32.9%)	
Previously diagnosed with nose problems	Yes	284 (66.3%)	145 (33.7%)	0.148
	No	279 (65.1%)	150 (34.9%)	
Previously diagnosed with throat problems	Yes	283 (66.0%)	146 (34.0%)	0.371
	No	282 (65.9%)	147 (34.1%)	
Previous surgery concerning ear, nose and throat	Yes	291 (67.9%)	138 (32.1%)	0.537
	No	279 (65.1%)	150 (34.9%)	

TABLE 4: The association between age, gender, health college, college department, nationality, previous clinical history, and level of knowledge about problems related to ENT.

Association between participants' demographics and knowledge of problems related to ENT

* Significant at the $p < 0.05$ level.

Most of the respondents (61%) showed good knowledge of common ENT problems. About 18% of the respondents had moderate knowledge, while 21% demonstrated poor knowledge, as shown in Figure 3.




FIGURE 3: Pie chart showing the knowledge score.

Discussion

The study aims to assess the awareness and practice of Najran University students toward common problems related to the ear, nose, and throat. The sample for the current study primarily consisted of students aged more than 20 years, with a predominance of females. The majority of them were in health colleges, faculties of medicine, and third level, with the majority of them being Saudi nationals.

The findings revealed that only 37.8% (n = 162) of the participants had good knowledge about problems related to ENT. The study showed that participants older than 20 years had better knowledge of common ENT problems than those younger than 20 years. Additionally, female participants showed a higher level of knowledge and awareness. The findings are consistent with the findings of the study conducted by Alassaf et al. (2023), which noted that women had a higher level of knowledge and awareness of problems related to the ears, nose, and throat, considering that locally, they are caretakers of family health issues rather than males in general [11]. Most of the participants showed good knowledge regarding the safety of using cotton swabs for ear cleaning. 61.1% (n = 262) and 61.8% (n = 265) of the participants had good knowledge about vaccinations recommended for the prevention of flu. More than half (62.2%, n = 267) of the participants had good knowledge about the use of vitamin C to treat and prevent influenza, and 77.6% (n = 333) had good knowledge regarding the definition of vertigo. The current findings mirror the findings of a study conducted by Alkholaiwi et al. (2020) in Saudi Arabia, which established good public awareness regarding the potential harms of using cotton swabs for ear cleaning [14]. The findings were also consistent with the findings of a study conducted by Di Berardino et al. (2013) in Milan, Italy, which showed that most of the participants in the study had good knowledge about the damage caused by cotton earbuds [9].

The findings revealed that the majority of the participants had good knowledge that some infections of the inner ear and middle ear cause dizziness and the appropriate measures for handling nose bleeding. The study noted that the majority of the participants, 87.2% (n = 374), had good knowledge with respect to the appropriate action to be taken in case of a sudden hearing problem or earache-going to the hospital-and that the information and knowledge about ENT issues were acquired by the participants from doctors. The findings concur with those of Alshehri et al. (2019), who reported a good level of awareness of going to the hospital as the most appropriate action to be taken in case of a sudden ear, nose, and throat problem [15].

The significant constraint and limitation in this investigation was the employment of a cross-sectional study design, which can only establish relations between factors but not causalities. As this was a survey-based study, recollection bias might be a limitation that also needs further investigation. Also, the study findings cannot be generalized to the entire Saudi Arabian student population, considering that they were conducted at only one institution.

Conclusions

The study revealed that 37.8% of the participants had good knowledge about problems related to the ears, nose, and throat. The participants older than 20 years had better knowledge of common ENT problems than those younger than 20 years. Female participants showed a higher level of knowledge and awareness of problems related to the ears, nose, and throat. The study noted that the participants in Health College, the faculty of medicine, and the third academic level had good knowledge about problems related to ENT. The study established that going to the hospital was the appropriate action to be taken in case of a sudden ENT

pathology. Therefore, we recommend concerted efforts be made among the medical community to increase knowledge about common ENT problems.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Acquisition, analysis, or interpretation of data: Khalid M. Alkhalifah, Meshal Alwabel, Saleh Y. Al Hareth

Drafting of the manuscript: Khalid M. Alkhalifah, Enas M. Al Hadi, Meshal Alwabel, Saleh Y. Al Hareth, Asaiel AL Hadi, Abdullah M. Assiri

Critical review of the manuscript for important intellectual content: Khalid M. Alkhalifah

Concept and design: Enas M. Al Hadi, Asaiel AL Hadi, Abdullah M. Assiri

Supervision: Abdullah M. Assiri

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. Committee of Scientific Research & Conferences in Najran University issued approval No. 4\3\2023. We are pleased to inform you that the ethical committee have reviewed the study protocol and ethically approved the study application to be conducted which is a matched cross sectional study under title of (Awareness and Practice of Najran University Students towards common problems related to Ear, Nose and Throat, Njran, KSA). **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Roux B, Sirois C, Simard M, Gagnon ME, Laroche ML: Potentially inappropriate medications in older adults: a population-based cohort study. *Fam Pract.* 2020, 37:173-9. [10.1093/fampra/cmz060](https://doi.org/10.1093/fampra/cmz060)
2. Ibekwe TS, Nwaorgu OG, Onakoya PA, Ibekwe PU: Spectrum of otorhinolaryngology emergencies in the elderly in Ibadan, Nigeria. *Niger J Med.* 2005, 14:411-4. [10.4314/njm.v14i4.37199](https://doi.org/10.4314/njm.v14i4.37199)
3. Fasunla AJ, Samdi M, Nwaorgu OG: An audit of ear, nose and throat diseases in a tertiary health institution in South-western Nigeria. *Pan Afr Med J.* 2013, 14:1. [10.11604/pamj.2013.14.1.1092](https://doi.org/10.11604/pamj.2013.14.1.1092)
4. Kirkwood CR, Clure HR, Brodsky R, Gould GH, Knaak R, Metcalf M: The diagnostic content of family practice: 50 most common diagnoses recorded in the WAMI community practices. *J Fam Pract.* 1982, 15:485-92.
5. McCormick A, Fleming D, Charlton J: Morbidity Statistics from General Practice 4th National Study 1992-1993. 1995.
6. Finley CR, Chan DS, Garrison S, Korownyk C, Kolber MR, Campbell S: What are the most common conditions in primary care?. Systematic review *Can Fam Physician.* 2018, 64:832-40.
7. WHO: Hearing care. (2006). http://www.who.int/pbd/deafness/activities/hearing_care/en/index.
8. Agrawal Y, Platz EA, Niparko JK: Prevalence of hearing loss and differences by demographic characteristics among US adults: data from the National Health and Nutrition Examination Survey, 1999-2004. *Arch Intern Med.* 2008, 168:1522-30. [10.1001/archinte.168.14.1522](https://doi.org/10.1001/archinte.168.14.1522)
9. Di Berardino F, Forti S, Iacona E, Orlandi GP, Ambrosetti U, Cesarani A: Public awareness of ear and hearing management as measured using a specific questionnaire. *Eur Arch Otorhinolaryngol.* 2013, 270:449-53. [10.1007/s00405-012-1961-3](https://doi.org/10.1007/s00405-012-1961-3)
10. Jalaladdin MS, Sindi EE, Kabli AF, et al.: An assessment of the knowledge and awareness of common otorhinolaryngology-related issues among school and university students in Makkah City, Saudi Arabia: a cross-sectional study. *Cureus.* 2023, 15:e37444. [10.7759/cureus.37444](https://doi.org/10.7759/cureus.37444)
11. Bhattacharyya N: The prevalence of pediatric voice and swallowing problems in the United States. *Laryngoscope.* 2015, 125:746-50. [10.1002/lary.24931](https://doi.org/10.1002/lary.24931)
12. Alkholaiwi F, Alharbi MM, Aldayel AY, Shadid AM, Almutairi FE, Dahmash AB: Public perception in the management and prevention of common ear, nose, and throat diseases in Saudi Arabia. *Jr Nat Sci Med.* 2020, 3:126-35. [10.4103/JNSM.JNSM_50_19](https://doi.org/10.4103/JNSM.JNSM_50_19)
13. Alqahtani AS: Knowledge and attitude toward Middle East respiratory syndrome coronavirus among health colleges' students in Najran, Saudi Arabia.
14. Alassaf HI, AlTurki AY, Almansour NM: Basic ear, nose, and throat knowledge among medical students: A multi-institutional study reflecting the current stage of ear, nose, and throat curriculum in undergraduate

- medical education. *Jr Nat Sci Med.* 2023;1, 6:59-65. [10.4103/jnsm.jnsm_169_20](https://doi.org/10.4103/jnsm.jnsm_169_20)
15. Alshehri KA, Alqulayti WM, Yaghmoor BE, Alem H: Public awareness of ear health and hearing loss in Jeddah, Saudi Arabia. *S Afr J Commun Disord.* 2019, 66:e1-6. [10.4102/sajcd.v66i1.633](https://doi.org/10.4102/sajcd.v66i1.633)