

Review began 02/12/2024  
Review ended 02/23/2024  
Published 02/27/2024

© Copyright 2024  
I et al. This is an open access article  
distributed under the terms of the Creative  
Commons Attribution License CC-BY 4.0.,  
which permits unrestricted use, distribution,  
and reproduction in any medium, provided  
the original author and source are credited.

# Identification and Evaluation of Cancer Stem Cells in Oral Squamous Cell Carcinoma and Oral Epithelial Dysplasia Using NANOG: An Immunohistochemical Study

Arya I<sup>1</sup>, Varun B. Raghavan Pillai<sup>2, 1</sup>, Anna P. Joseph<sup>1</sup>, Pratibha Ramani<sup>2</sup>, Jayanthi P<sup>3</sup>, Karthikeyan Ramalingam<sup>2</sup>

1. Oral and Maxillofacial Pathology, PMS College of Dental Science and Research, Trivandrum, IND 2. Oral Pathology and Microbiology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, IND 3. Oral and Maxillofacial Pathology, Azeezia College of Dental Sciences and Research, Kollam, IND

**Corresponding author:** Pratibha Ramani, pratibaramani@saveetha.com

---

## Abstract

**Background:** Squamous cell carcinoma of the oral cavity may show precursor lesions, termed as potentially malignant disorders, of which leukoplakia is the most frequent one. Oral leukoplakia is a clinical diagnosis for which the histological diagnosis may be either hyperplasia or oral epithelial dysplasia (OED) and sometimes even oral squamous cell carcinoma (OSCC). Cancer stem cells (CSCs), identified in various tumors, are a specific group of cells that exhibit the properties of self-renewal and differentiation. Among the various biomarkers that identify CSCs, the transcription factor NANOG is considered to be a significant one.

**Aim:** In this study, we intend to identify and compare the immunohistochemical expression of NANOG in OSCC, OED, and normal oral mucosa.

**Methodology:** Tissue blocks of OSCC (n=28), OED (n=28), and normal oral mucosa (n=28) were used in this study. Specimens were immunohistochemically analyzed for NANOG expression. The results were statistically analyzed using one-way ANOVA, Games-Howell post hoc, and Student t-test. Statistical Product and Service Solutions (SPSS, version 21; IBM SPSS Statistics for Windows, Armonk, NY) software was used for performing the statistical analysis, and the level of significance was set as 0.05.

**Observations:** NANOG expression was higher in OSCC when compared to oral dysplasias and normal oral mucosa, in decreasing order. A significantly higher histo-score and labeling index score were observed in OSCC and oral dysplasias compared to normal oral mucosa ( $p < 0.001$ ).

**Conclusion:** The expression levels of NANOG were positively correlated with disease progression in OSCC, implicating that NANOG can be used as a surrogate marker of oral oncogenesis and prognosis. Therefore, decoding the molecular mechanisms of NANOG regulation in the progression of cancer helps in developing new therapeutic strategies for oral cancer.

---

**Categories:** Pathology, Dentistry, Oncology

**Keywords:** expression, progression, potentially malignant, prognosis, marker, oral leukoplakia, immunohistochemistry, oscc, oral epithelial dysplasias, cancer stem cells

## Introduction

Oral squamous cell carcinoma (OSCC) forms the major type of cancer in the head and neck region and poses a significant threat to the well-being of the patient [1,2]. The cumulative incidence of oral cancer is four per 100,000 persons with approximately 350,000 cases diagnosed each year. Different nations around the globe exhibit varied prevalence of OSCC and in developing countries, such as India, around 30% of oral cancer cases are diagnosed every year [3,4]. A staggering 22.9% of cancer deaths in India are attributed to OSCC [5,6].

In about 15%-45% of the cases, oral cancer might show visible precursor lesions, of which oral leukoplakia (OL) is the most frequently occurring lesion of the mouth. Oral leukoplakia is predominantly a non-scrappable white patch clinically diagnosed after excluding other possible causes of white lesions in the mouth [7]. Histopathologically, OL may exhibit features of either hyperplasia or oral epithelial dysplasia (OED) and sometimes even frank carcinoma. Even though OL has multifactorial etiologies, tobacco is considered to be the most common causative agent. The prevalence of leukoplakia ranges between 1.5% and 4% throughout the world, whereas, in India, this range is wider with an estimated value of 0.2%-4.8% [8,9].

### How to cite this article

I A, Raghavan Pillai V B, P. Joseph A, et al. (February 27, 2024) Identification and Evaluation of Cancer Stem Cells in Oral Squamous Cell Carcinoma and Oral Epithelial Dysplasia Using NANOG: An Immunohistochemical Study. Cureus 16(2): e55111. DOI 10.7759/cureus.55111

Cancer stem cells (CSCs) have been identified in tumors and exhibit self-renewal and differentiation properties. It has been demonstrated that, when these cells are transplanted into an animal, they have the capacity for tumor formation [10]. In the year 1994, John Dick and his co-workers first identified cancer stem cells in neoplasms of hematolymphoid origin [11]. In later years, these cells were found in various primary malignant tumors and in OSCC cell lines. The presence of cancer stem cells was associated with poor prognosis as the result of their involvement in lymph node metastasis and local recurrence. CSCs in OSCC express proteins such as NANOG, OCT4, and SOX2, which regulate embryonic stem cells [12,13].

NANOG is a stem cell transcription factor involved in human embryogenesis, cell fate determination, proliferation, and apoptosis. The expression of NANOG is very low or absent after birth and remains so throughout the lifespan. However, NANOG expression is detectable in a small proportion of cancer cells that show stem cell-like properties. Higher expression of NANOG has been observed in various solid tumors of the breast, colon, lung, liver, and cervix. Elevated values of NANOG especially in cancer cells were described by previous research, which means that the NANOG gene is a critical member of cancer stem cells [14]. This study was done to evaluate the cancer stem cells in OSCC and OED using immunoeexpression of NANOG.

## Materials And Methods

Sample size calculation was performed using the reported data [14,15] due to a lack of relevant literature. The tissue sections of oral leukoplakia with OED (n=28), OSCC (n=28), and normal oral mucosa (n=28) were taken from the archival blocks stored in the Department of Oral Pathology Department at PMS Dental College, Trivandrum, India.

Ethical clearance was obtained from the institutional ethical committee before conducting this study (PMS/IEC/2020-21/01). Clinically recorded cases of leukoplakia, which are histopathologically graded as epithelial dysplasia, and histopathologically confirmed cases of OSCC were included in the study. Biopsy specimens from both genders falling under the age group of 20-70 years were included. The tissue sections without epithelium were not considered for this particular study. Tissue specimens of normal oral mucosa harvested during surgical removal of impacted third molars constituted the control group. Those individuals with any systemic illness or any known inflammatory conditions about the third molar were excluded.

Four micron-thick tissue sections were made and transferred onto the aminopropyltriethoxysilan (APES)-coated slide. Seminoma was kept as positive immunohistochemistry (IHC) control. Deparaffinization was done using xylene for 10 minutes, followed by rehydration, with the help of descending grades of alcohol (100%, 90%, and 70%). The slides were then placed in a jar containing Tris EDTA buffer (pH-9), and antigen retrieval was performed in a 5 L pressure cooker at 100°C. After blocking the endogenous peroxidase, the tissue sections were incubated overnight with rabbit anti-NANOG monoclonal antibody (D73G4, Scientific solutions, 1:800 (diluent-signal stain antibody diluent #8112)). After washing with an immuno-wash buffer, the slides were incubated with polyExcel HRP secondary antibody for 30 minutes and with diaminobenzidine (DAB) for 30 minutes. Finally, the sections were counterstained with hematoxylin and observed under light microscopy.

Two independent pathologists performed the analysis of NANOG expression through light microscopy at a magnification of 40x and evaluated the number of positive and negative cells. The manual cell counting was done in five consecutive high-power fields for each slide, and cell counts were averaged. The areas with strong NANOG expression were selected for cell counting. In OSCC, NANOG-positive cells were evaluated mainly in the tumor-invading front, whereas in OED and normal mucosa, positive cells were evaluated within the epithelium. Testicular seminoma was taken as the positive control to compare the expressions of staining intensities.

For analysis of the staining results in OSCC, OED, and normal mucosa, a semi-quantitative approach using histo-score (H-score) was used [15]. The NANOG expression was assessed based on the nuclear and cytoplasmic staining of tumor cells, and the percentage of tumor cells was counted. The intensity score was considered negative for cells that did not show any staining (score 0), weak staining less than that of positive controls (score 1), moderate staining equivalent to positive control (score 2), and intense staining stronger than positive control (score 3). The following formula was used to assign the percentage of cells at each staining intensity level:  $\{[1 \times \% \text{ cells } 1+] + [2 \times \% \text{ cells } 2+] + [3 \times \% \text{ cells } 3+]\}$ . The percentage of positive cells was calculated as the labeling index (LI), which was derived by adding up the total number of NANOG-expressing tumor cells and dividing that value by the total number of tumor cells [16].


Statistical Product and Service Solutions (SPSS, version 21; IBM SPSS Statistics for Windows, Armonk, NY) software was used to perform all the statistical analyses. Comparison between the groups was done with one-way ANOVA and Games-Howell post hoc analysis. Comparison between the various grades of OED and OSCC was done with Student's t-test, and a p-value less than 0.05 was considered to be statistically significant.

## Results


Retrospective analysis of our archival records revealed that, among the 28 samples of OSCC, 20 were males, eight were females, and they belonged to the age group between 40 and 85 years. Among the 28 samples of OED, 19 were males, nine were females, and they belonged to the age group between 20 and 90 years. Among the 28 samples of normal mucosa, 18 were males, 10 were females, and they belonged to the age group between 20 and 50 years. Chewing tobacco was the most prevalent deleterious habit (87%) noted in both OSCC and OED groups. Smoking tobacco was noted in the remaining 13% of both OSCC and OED groups.

The biopsy specimens of OSCC and OED were taken from different anatomic sites within the oral cavity. Out of the 28 samples of OSCC, nine were located on the buccal mucosa, 12 on the tongue, four on the alveolar mucosa, and three on the gingiva. Out of the 28 samples of OED, 10 were located on the buccal mucosa, 11 on the tongue, five on the alveolar mucosa, and two on the gingiva.


All sampled categories, such as OSCC, OED, and normal oral mucosa, were compared for NANOG expression using the immunohistochemical scoring analysis, H-score, and labeling index (LI). Photomicrographs of the study groups and control show the NANOG expression (Figures 1-4).



**FIGURE 1: Photomicrograph of diffuse NANOG immunohistochemical expression in squamous cell carcinoma (IHC, 40x)**




**FIGURE 2: Photomicrograph of focal NANOG immunohistochemical expression in epithelial dysplasia (IHC, 10x)**




**FIGURE 3: Photomicrograph showing low NANOG immunohistochemical expression in normal epithelium (IHC, 20x)**






**FIGURE 4: Photomicrograph of NANOG immunohistochemical expression in the positive control of testicular seminoma (IHC, 10x)**

When comparing the mean value of the H-score and LI score between the three groups, the mean H-scores of OSCC, OED, and normal mucosa were 182, 138, and 5.7, respectively (Figure 5). A significantly higher H-score was observed in OSCC when comparing all three study groups ( $p<0.001$ ).



**FIGURE 5: Graph showing the comparison of H-score of NANOG among the study groups**  
SCC - Squamous Cell Carcinoma

The mean LI scores of OSCC, OED, and normal oral mucosa were 74, 48, and 2.5, respectively. The LI was significantly higher in OSCC than in OED and normal mucosa ( $p<0.001$ ) (Figure 6).



**FIGURE 6: Graph showing the comparison of the LI scores of NANOG among the study groups**

SCC - Squamous Cell Carcinoma

When comparing the H-score and LI scores among well-differentiated (n=12) and moderately differentiated (n=16) OSCCs, the following results were obtained. In OSCC, the expression was mainly noted in the invasive tumor front and within the tumor islands in the connective tissue. In OED, the expression of NANOG was seen throughout the epithelium, whereas in normal mucosa, it was restricted to the basal and para-basal layers of the epithelium. The mean H-score and LI were higher in moderately differentiated tumors, as compared to well-differentiated tumors. However, these differences were not statistically significant (Table 1).

Student's t-test			
Category	Groups	Mean (SD)	p-value
H-score	Well-differentiated OSCC	173.2 (80)	0.514
	Moderately differentiated OSCC	190.8 (67.8)	
Labeling Index (LI)	Well-differentiated OSCC	74.63 (24.6)	0.889
	Moderately differentiated OSCC	73.55 (16.2)	

**TABLE 1: Comparison of NANOG expression using the H score and labeling index among different grades of OSCC with Student's t-test**

OSCC - Oral Squamous Cell Carcinoma

The mean value of the H-score was higher in high-risk OED (n=11) than in low-risk OED (n=17), and the difference was statistically significant. The statistical analysis between OSCC and OED was done using the Games-Howell post-hoc test. The H-score and LI score of NANOG were significantly higher in OSCC compared to those in OED (p=0.02). Though the mean value of the LI score was higher in high-risk OED than in low-risk OED, a statistical significance could not be obtained (p=0.071) (Table 2).

Student's t-test			
Category	Groups	Mean (SD)	p-value
H-score	Low-risk dysplasia	88.08 (36)	0.02*
	High-risk dysplasia	188 (63.2)	
Labeling Index (LI)	Low-risk dysplasia	35.15 (14.3)	0.07
	High-risk dysplasia	60.8 (25)	

**TABLE 2: Comparison of NANOG expression using the H score and labeling index among different grades of dysplasia by Student's t-test**

\* - statistically significant value: p<0.05

Discussion

Squamous cell carcinoma of the oral cavity forms the predominant group of malignant neoplasms in the mouth [17]. Most OSCCs arise from precursor lesions, known collectively as potentially oral malignant disorders, of which OL is the most common one with a malignant transformation rate of around 1.1%-40.8% [18]. Early identification and prompt diagnosis of leukoplakia might be essential to control their transformation into malignancy. Microscopic evaluation and grading of dysplasia is still considered to be the best practical method for predicting malignant conversion. However, the major disadvantage of the currently used grading system is the intraobserver and inter-observer variability. Therefore, continuous search and research to identify accurate biomarkers that can predict the malignant transformation of oral leukoplakia are necessary.

Cancer stem cells are involved not only in the initiation and progression of tumors but are associated with lymph node metastasis and recurrence. Studies have shown that the transcription factor NANOG is a key regulator of CSC properties in carcinomas of the head and neck region. NANOG activates various signaling pathways to promote angiogenesis and also blocks apoptosis. In addition, NANOG was shown to reduce the expression of E-cadherin levels, which might lead to metastasis [19].

Research studies in various cancers have shown over-expression of NANOG. Meng et al. studied the expression of NANOG in colorectal carcinoma and showed that a low survival rate is directly correlated with higher levels of NANOG [20]. Nagata et al. found that NANOG expression levels were favorable prognostic factors for triple-negative breast cancer patients [21]. The combined expression of NANOG with other CSC markers such as OCT-4, and SOX-2, and proliferation markers, such as KI-67 and PCNA, were better predictors of relapse in patients with gastric cancer, as demonstrated by Li et al. [22]. The positive expression of NANOG in serous carcinoma of the ovary was correlated with poor prognosis by Lee et al. [23]. Zhao et al. linked the high levels of NANOG protein to high-grade tumors and metastatic lymph nodes in intestinal-type gastric cancer [24]. All these studies have shown that NANOG may be considered a potential prognostic marker in most primary tumors. However, only a few studies are found in the literature that compare NANOG expression in OSCC and epithelial dysplasia. Hence, in this present study, we have addressed this lacuna.

In our study, NANOG expression was noticed in 71.4% of dysplasia cases, and all these samples showed cytoplasmic staining of cells. When comparing the NANOG expression in different grades of OED, the expression was found to be higher in high-risk dysplasias than in low-risk dysplasias. This is consistent with the previous studies of de Vicente et al., where increased NANOG expression was observed in higher grades of dysplasia. In their study, 55 cases of OED were analyzed for immunohistochemical expression of NANOG protein, and the results showed that 3.6% and 16.4% of cases exhibited nuclear and cytoplasmic staining, respectively. Additionally, after a mean follow-up of 185 months, patients who harbored NANOG-positive cells showed significantly higher chances for malignant progression to oral cancer [15].

In the current study, NANOG protein expression was positive in 92.8% (n=26/28) of OSCC cases, all of which were primarily observed in the cytoplasm of tumor cells, whereas in germ cell tumors, a nuclear staining pattern was seen. Our results are similar to the study conducted by Grubelnik et al. who assessed both the protein and mRNA expressions of NANOG in 30 samples of OSCC and found that 27 cases exhibited positive staining mainly in the cytoplasm of tumor cells [14].

Our study also demonstrates that the expression of NANOG is increased in higher grades of OSCC and high-risk OED similar to previous studies [25-28]. The reason for the two different staining patterns remains unclear. However, Gu et al. conducted a study on cervical cancer and showed that cellular localization of

NANOG was related to the stage of the tumor and the type of cell involved. They suggested that the stromal cytoplasmic NANOG staining was associated with the progression of cervical cancer [28]. Therefore, our study uncovers the potential application of NANOG expression as an early surrogate marker for predicting malignant conversion in leukoplakia and a potential prognostic indicator in OSCC.

The limitations of our study were the retrospective nature of samples and a single-center evaluation of NANOG expression. In our study, the NANOG protein level was higher in OSCC compared to OED and normal mucosa using IHC. Even though there was an overall increase in the immunoeexpression of NANOG in OSCC, few specimens showed lower expressions when compared to OED and normal mucosa. We also did not specifically analyze other histopathological parameters due to differences in the composition of study groups. The number of samples within each grade of OSCC was less, and, hence, a statistically significant difference in NANOG expression between moderately differentiated and well-differentiated OSCC could not be achieved. Further, our study samples did not include poorly differentiated OSCC, and the expression of NANOG in high-grade tumors could not be ascertained in this study. The increased expression of NANOG in high-risk dysplasias compared to low-risk dysplasias uncovers the potential of NANOG as a surrogate marker for malignant transformation into OSCC. However, longitudinal follow-up studies on a larger sample size would help us assess the exact role of NANOG in the malignant transformation and prognosis of OSCC and OED. Multiple institutions across the globe should be involved to assess the clinico-pathological correlation of NANOG expression with clinical presentation, regional invasion, distant metastasis, recurrence, disease-free survival, and mortality rate. NANOG expression in various potentially malignant disorders, along with continuous monitoring, will shed light on its prognostic ability as a biomarker. In the future, longitudinal follow-up studies on a larger sample size would help us assess the exact role of NANOG in the malignant transformation and prognosis of OSCC and OED.

## Conclusions

This study identifies the presence of cancer stem cells using immunohistochemical expression of NANOG in OSCC, compared with OED and normal mucosa. In conclusion, NANOG expression levels seem to be positively correlated with disease progression in OSCC, implicating that NANOG can be used as a biomarker of oral oncogenesis and prognosis. NANOG can also be used as a biomarker to predict the progression of the disease from dysplasia to OSCC. Therefore, decoding the molecular mechanisms of NANOG regulation can help in developing new therapeutic strategies for oral cancer.

## Additional Information

### Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

**Acquisition, analysis, or interpretation of data:** Karthikeyan Ramalingam, Arya I, Varun B. Raghavan Pillai, Jayanthi P

**Drafting of the manuscript:** Karthikeyan Ramalingam, Varun B. Raghavan Pillai, Anna P. Joseph, Jayanthi P

**Critical review of the manuscript for important intellectual content:** Karthikeyan Ramalingam, Arya I, Varun B. Raghavan Pillai, Anna P. Joseph, Pratibha Ramani, Jayanthi P

**Concept and design:** Varun B. Raghavan Pillai, Anna P. Joseph, Pratibha Ramani, Jayanthi P

### Disclosures

**Human subjects:** Consent was obtained or waived by all participants in this study. PMS Dental College Institutional Ethical Committee issued approval PMS/IEC/2020-21/01. Ethical clearance was obtained from the institutional ethical committee before conducting this study. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

## References

1. Warnakulasuriya S: Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol.* 2009, 45:309-16. [10.1016/j.oraloncology.2008.06.002](https://doi.org/10.1016/j.oraloncology.2008.06.002)
2. Lin NC, Hsien SI, Hsu JT, Chen MY: Impact on patients with oral squamous cell carcinoma in different anatomical subsites: a single-center study in Taiwan. *Sci Rep.* 2021, 11:15446. [10.1038/s41598-021-95007-5](https://doi.org/10.1038/s41598-021-95007-5)



3. Joshi P, Dutta S, Chaturvedi P, Nair S: Head and neck cancers in developing countries . *Rambam Maimonides Med J*. 2014, 5:e0009. [10.5041/RMMJ.10143](#)
4. Sankaranarayanan R, Ramadas K, Thomas G, Muwonge R, Thara S, Mathew B: Effect of screening on oral cancer mortality in Kerala, India: a cluster-randomised controlled trial. *Lancet*. 2005, 365:1927-33. [10.1016/S0140-6736\(05\)66658-5](#)
5. Dikshit R, Gupta PC, Ramasundarahettige C: Cancer mortality in India: a nationally representative survey . *Lancet*. 2012, 379:1807-16. [10.1016/S0140-6736\(12\)60358-4](#)
6. Chen SH, Hsiao SY, Chang KY, Chang JY: New insights into oral squamous cell carcinoma: from clinical aspects to molecular tumorigenesis. *Int J Mol Sci*. 2021, 22:2252. [10.3390/ijms22052252](#)
7. Hsue SS, Wang WC, Chen CH, Lin CC, Chen YK, Lin LM: Malignant transformation in 1458 patients with potentially malignant oral mucosal disorders: a follow-up study based in a Taiwanese hospital. *J Oral Pathol Med*. 2007, 36:25-9. [10.1111/j.1600-0714.2006.00491.x](#)
8. Brouns ER, Evren I, Wils LJ, Poell JB, Brakenhoff RH, Bloemena E, de Visscher JG: Oral leukoplakia classification and staging system with incorporation of differentiated dysplasia. *Oral Dis*. 2023, 29:2667-76. [10.1111/odi.14295](#)
9. van der Waal I: Oral leukoplakia, the ongoing discussion on definition and terminology . *Med Oral Patol Oral Cir Bucal*. 2015, 20:e685-92. [10.4317/medoral.21007](#)
10. Yu Z, Pestell TG, Lisanti MP, Pestell RG: Cancer stem cells. *Int J Biochem Cell Biol*. 2012, 44:2144-51. [10.1016/j.biocel.2012.08.022](#)
11. Lapidot T, Sirard C, Vormoor J, et al.: A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature*. 1994, 367:645-8. [10.1038/367645a0](#)
12. Walcher L, Kistenmacher AK, Suo H, et al.: Cancer stem cells—origins and biomarkers: perspectives for targeted personalized therapies. *Front Immunol*. 2020, 11:1280. [10.3389/fimmu.2020.01280](#)
13. Shin KH, Kim RH: An updated review of oral cancer stem cells and their stemness regulation . *Crit Rev Oncog*. 2018, 23:189-200. [10.1615/CritRevOncog.2018027501](#)
14. Grubelnik G, Boštjančič E, Grošelj A, Zidar N: Expression of NANOG and its regulation in oral squamous cell carcinoma. *Biomed Res Int*. 2020, 2020:10.1155/2020/8573793
15. de Vicente JC, Rodríguez-Santamarta T, Rodrigo JP, et al.: The emerging role of NANOG as an early cancer risk biomarker in patients with oral potentially malignant disorders. *J Clin Med*. 2019, 8:1376. [10.3390/jcm8091376](#)
16. Fedchenko N, Reifenrath J: Different approaches for interpretation and reporting of immunohistochemistry analysis results in the bone tissue - a review. *Diagn Pathol*. 2014, 9:221. [10.1186/s13000-014-0221-9](#)
17. Pires FR, Ramos AB, Oliveira JB, Tavares AS, Luz PS, Santos TC: Oral squamous cell carcinoma: clinicopathological features from 346 cases from a single oral pathology service during an 8-year period. *J Appl Oral Sci*. 2013, 21:460-7.
18. Aguirre-Urizar JM, Lafuente-Ibáñez de Mendoza I, Warnakulasuriya S: Malignant transformation of oral leukoplakia: systematic review and meta-analysis of the last 5 years. *Oral Dis*. 2021, 27:1881-95. [10.1111/odi.13810](#)
19. Najafzadeh B, Asadzadeh Z, Motafakker Azad R, et al.: The oncogenic potential of NANOG: an important cancer induction mediator. *J Cell Physiol*. 2021, 236:2443-58. [10.1002/jcp.30063](#)
20. Meng HM, Zheng P, Wang XY, et al.: Over-expression of Nanog predicts tumor progression and poor prognosis in colorectal cancer. *Cancer Biol Ther*. 2010, 9:295-302. [10.4161/cbt.9.4.10666](#)
21. Nagata T, Shimada Y, Sekine S, et al.: KLF4 and NANOG are prognostic biomarkers for triple-negative breast cancer. *Breast Cancer*. 2017, 24:326-35. [10.1007/s12282-016-0708-1](#)
22. Li N, Deng W, Ma J, et al.: Prognostic evaluation of Nanog, Oct4, Sox2, PCNA, Ki67 and E-cadherin expression in gastric cancer. *Med Oncol*. 2015, 32:433. [10.1007/s12032-014-0433-6](#)
23. Lee M, Nam EJ, Kim SW, Kim S, Kim JH, Kim YT: Prognostic impact of the cancer stem cell-related marker NANOG in ovarian serous carcinoma. *Int J Gynecol Cancer*. 2012, 22:1489-96. [10.1097/IGJ.0b013e3182738307](#)
24. Zhao L, Liu J, Chen S, Fang C, Zhang X, Luo Z: Prognostic significance of NANOG expression in solid tumors: a meta-analysis. *Onco Targets Ther*. 2018, 11:5515-26. [10.2147/OTT.S169593](#)
25. Lee YS, Kim HK, Chung S, Kim KS, Dutta A: Depletion of human micro-RNA miR-125b reveals that it is critical for the proliferation of differentiated cells but not for the down-regulation of putative targets during differentiation. *J Biol Chem*. 2005, 280:16635-41. [10.1074/jbc.M412247200](#)
26. Watanabe M, Ohnishi Y, Inoue H, Wato M, Tanaka A, Kakudo K, Nozaki M: NANOG expression correlates with differentiation, metastasis and resistance to preoperative adjuvant therapy in oral squamous cell carcinoma. *Oncol Lett*. 2014, 7:35-40. [10.3892/ol.2013.1690](#)
27. Kim HM, Kang YH, Byun JH, Jang SJ, Rho GJ, Lee JS, Park BW: Midkine and NANOG have similar immunohistochemical expression patterns and contribute equally to an adverse prognosis of oral squamous cell carcinoma. *Int J Mol Sci*. 2017, 18:2339. [10.3390/ijms18112339](#)
28. Gu TT, Liu SY, Zheng PS: Cytoplasmic NANOG-positive stromal cells promote human cervical cancer progression. *Am J Pathol*. 2012, 181:652-61. [10.1016/j.ajpath.2012.04.008](#)