

Trustworthiness, Value, Danger, and Readability of ChatGPT-Generated Responses to Health Questions Related to Pulmonary Arterial Hypertension

Review began 10/02/2024
Review ended 10/10/2024
Published 10/14/2024

© Copyright 2024
Kerkütlüoğlu et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

DOI: 10.7759/cureus.71472

Murat Kerkütlüoğlu¹, Erhan Kaya², Rasim Gökmen²

1. Cardiology, Kahramanmaraş Sütçü İmam University, Kahramanmaraş, TUR 2. Public Health, Kahramanmaraş Sütçü İmam University, Kahramanmaraş, TUR

Corresponding author: Rasim Gökmen, drrasimgokmen@gmail.com

Abstract

Aim: To enhance outcomes for patients with pulmonary arterial hypertension (PAH), comprehensive and individualized therapy is needed. A large language model called Generative Pre-trained Transformer (ChatGPT) has the ability to provide expert yet patient-friendly care. We wanted to determine how well ChatGPT could accurately and consistently respond to inquiries on knowledge and management for PAH.

Materials and methods: When 20 PAH patients were diagnosed, they were asked what concerns they had about PAH and what they had researched online. In the evaluation, it was determined that patients frequently searched the Internet for answers to eight queries. These eight queries were posed to ChatGPT, and their responses were recorded. Ten experts in the field of PAH assessed the trustworthiness, value, and hazard of the answers generated by the ChatGPT.

Results: According to evaluations conducted by experts, the ChatGPT-generated responses were deemed trustworthy with an average score of 8.4 (7.7-9.2) and valuable with an average score of 7.9 (7.4-8.2). Based on the statistical analysis, it can be inferred that most professionals believed that the utilization of prompts provided by ChatGPT did not present a substantial risk, with a mean of 2.1 (1.7-2.5). The answers were assessed for readability using two different indicators, namely the Flesch-Kincaid Grade Level (FKGL) and the Simple Measure of Gobbledygook (SMOG). The average FKGL value was determined to be 13.52 ± 2.40 , indicating a "difficult" level of readability.

Conclusion: ChatGPT provides reliable PAH-related information, but it is important to seek professional medical advice before making any decisions regarding PAH. ChatGPT can only provide general information and support, but a qualified healthcare provider can offer tailored recommendations.

Categories: Public Health, Pulmonology, Healthcare Technology

Keywords: artificial intelligence, chatbot, language models, patient information, pulmonary arterial hypertension, readability, safety

Introduction

This article was previously presented as a meeting abstract at the 3rd International Congress on Innovative Approaches in Medical and Health Sciences on 08 October 2023.

The utilization of the internet for the purpose of obtaining health-related information is a prevalent occurrence. Patients frequently resort to seeking information online due to a lack of communication from their healthcare provider, either due to forgetfulness, inadequate explanation, or a failure to comprehend the information conveyed [1]. Frequently, patients engage in the act of researching and presenting information to their healthcare provider to verify or challenge the decisions made regarding their treatment [2]. The ChatGPT, which is a language model founded on the third generation of the Generative Pre-trained Transformer (GPT-3) architecture, was introduced in November 2022. In a matter of weeks, the recently developed artificial intelligence (AI) system garnered a user base of 100 million and received frequent coverage in popular media outlets. ChatGPT is a robust natural language processing model with the ability to comprehend and produce textual content. The abundance of information available through this resource renders it highly valuable for diverse purposes, including but not limited to language interpretation, condensation of text, and generation of written content. A commonly expressed critique of ChatGPT pertains to the potential inaccuracy of the generated texts, which may occasionally produce erroneous or fictitious content. Within the field of AI, a hallucination refers to a response generated by an AI system that exhibits a high degree of confidence, despite lacking sufficient justification from its training data [3]. The language model generates a text that lacks a basis in reality or empirical evidence.

How to cite this article

Kerkütlüoğlu M, Kaya E, Gökmen R (October 14, 2024) Trustworthiness, Value, Danger, and Readability of ChatGPT-Generated Responses to Health Questions Related to Pulmonary Arterial Hypertension. Cureus 16(10): e71472. DOI 10.7759/cureus.71472

ChatGPT is a communication tool that is commonly utilized by healthcare professionals and consumers [4]. Large language models, such as ChatGPT, are poised to significantly alter the manner in which patients seek information pertaining to their health status [5]. A recent review conducted an investigation into the efficacy of ChatGPT in the domains of healthcare education, research, and practice. The review has also brought to attention certain limitations that may be encountered [6]. One of the constraints identified in the study relates to the potential for erroneous information and hallucinatory effects. The review suggests that further research is necessary to assess the substance of language models as well as their potential to promote academic and scientific progress, with a specific emphasis on healthcare environments.

Patients diagnosed with pulmonary arterial hypertension (PAH) exhibit a positive attitude toward utilizing internet-based technologies [6]. Patients with PAH and their caregivers have the option to seek information pertaining to their condition on ChatGPT. However, the reliability, caliber, and readability of the information available on ChatGPT with regard to PAH remain uncertain. Despite the fact that ChatGPT is readily accessible, free, and simple to comprehend, these characteristics may not always apply to us. Can ChatGPT be considered a credible source of information pertaining to PAH?

Materials And Methods

When 20 PAH patients were diagnosed, they were asked what concerns they had about PAH and what they had researched online. In the evaluation, it was determined that patients frequently searched the internet for answers to eight queries. Table 1 contains the eight queries posed to ChatGPT. The ChatGPT-generated responses have been logged (Supplementary material 1).

Questions	Percent
1. What is pulmonary arterial hypertension?	100% of patients
2. What causes pulmonary arterial hypertension?	100% of patients
3. Is pulmonary hypertension the same as hypertension as we know it?	60% of patients
4. What are the symptoms of pulmonary arterial hypertension?	100% of patients
5. Who has pulmonary arterial hypertension?	80% of patients
6. Pregnancy in a patient with pulmonary arterial hypertension?	60% of patients
7. How is pulmonary arterial hypertension diagnosed?	100% of patients
8. How is pulmonary arterial hypertension treated?	100% of patients

TABLE 1: ChatGPT was asked eight questions about PAH from virtual patients

PAH: pulmonary arterial hypertension

A questionnaire was created exclusively for the purpose of this research (Supplementary material 2). A set of questionnaires was administered to all 10 medical experts who are members of the Turkish Cardiology Association's PAH study group and are cardiologists actively involved in clinical research and studies in this field. Ten cardiologists were selected using the random sampling method from among the volunteer members of this study group. The ChatGPT responses were evaluated by these professionals based on trustworthiness, value, and danger, utilizing a numerical rating system ranging from 1 to 10. The greater the numerical value assigned to a response, the greater its perceived level of trustworthiness, value, or danger. This methodology has been used as explained in a previous study addressing a similar topic [1].

The readability of websites was assessed using the Flesch-Kincaid (FK) grade and the Simple Measure of Gobbledygook (SMOG). The scores from these two formulas were obtained using an online tool (<https://www.webfx.com/tools/read-able/>). Indicators such as FK and SMOG are formulated based on the word in the sentence and the number of syllables in the word. High scores indicate low readability levels of websites. The readability of the text was categorized as "easy" (grade 6), "average" (grades 6-10), or "difficult" (grade >10) based on the Flesch-Kincaid Grade Level (FKGL) indicator (Table 2) [7,8].

Parameters	Formulas
FKGL	$(11.8 \times (\text{total syllables} / \text{total words})) + (0.39 \times (\text{total words} / \text{total sentences})) - 15.59$
SMOG	$3.1291 + 1.0430((\text{complex words})(30 / \text{sentences}))$
FKGL: Flesch-Kincaid Grade Level; SMOG: Simple Measure of Gobbledygook	

TABLE 2: Readability formulas provide indications of text readability

Means and quartiles are reported for descriptive statistics. A line graph is used to graphically represent data.

This study received approval from the Ethics Committee of Kahramanmaraş Sütçü İmam University for Medical Research, under approval number 2023/15-02.

Results

In the examination of ChatGPT's responses in terms of trustworthiness, value, and danger, responses from the experts' questionnaire determined that the responses produced by ChatGPT were trustworthy, with a mean score of 8.4 (7.7-9.2), and valuable, with a mean score of 7.9 (7.4-8.2). According to the statistical analysis, with a mean of 2.1 (1.7-2.5), the majority of professionals held the view that utilizing the prompts supplied by ChatGPT did not pose a significant risk (Figure 1).




FIGURE 1: PAH experts assessment result on the reliability, value, and danger of using ChatGPT-generated responses

PAH: pulmonary arterial hypertension

According to a panel of medical expert cardiologists in the field, the findings generally show that ChatGPT has the capacity to produce trustworthy and significant data regarding PAH.

In their explanations, the experts expressed their appreciation for ChatGPT's nuanced and comprehensive responses, as well as their recommendation to consult with healthcare professionals for further direction.

The readability of the obtained ChatGPT responses was evaluated using the scores derived from the Flesch-Kincaid (FK) grade and the Simple Measure of Gobbledygook (SMOG) formulas, with average values found to be 13.52 for FKGL and 12.49 for SMOG (Table 3).

	Mean (SD)	Median (min-max)
FKGL	13.52 (2.40)	13.10 (10.30-17.50)
SMOG	12.49 (2.27)	11.95 (9.70-16.10)
FKGL: Flesch-Kincaid Grade Level; SMOG: Simple Measure of Gobbledygook		

TABLE 3: Information quality scores and readability indicators assess the grade level of the content

Discussion

PAH is a chronic and severe cardiopulmonary disorder that is characterized by the proliferation of cells and fibrosis in the small pulmonary arteries. This leads to a gradual increase in pulmonary vascular resistance [9]. The development of PAH initiates in the pulmonary circulation; however, it is the primary cause of morbidity and mortality due to right heart failure [10]. Contemporary therapeutic approaches for PAH entail the intervention of aberrant signaling pathways in the pulmonary vasculature, with the objective of mitigating the afterload on the right ventricle and forestalling the advancement toward right-sided heart failure and mortality [9]. In the last two decades, there have been significant changes in the epidemiology and treatment of PAH. PAH is a disease that is not commonly found, with a projected occurrence of 15-50 cases per million people [11]. Individuals afflicted with uncommon diseases, such as PAH, exhibit a strong desire to obtain knowledge pertaining to their condition and pursue solutions to their inquiries regarding the prognosis of particular circumstances and therapeutic procedures through online resources.

Consumers may now easily access a wealth of highly technical information thanks to the development of the internet. Because of this, a significant number of individuals are now turning to Google rather than a qualified medical professional for the medical advice they need. While it doesn't appear that this practice will disappear any time soon, ChatGPT and other types of AI could displace it. The emergence of ChatGPT is attributed to the progress made in large language models, signifying a new era of AI technologies [12]. According to a report, ChatGPT reached 100 million users within 64 days of its launch on November 30, 2022 [13]. The platform is renowned for its proficiency in generating text that closely resembles human quality and covers a diverse array of subjects [13].

The term "artificial intelligence" refers to a cross-disciplinary field that combines computer science and linguistics with the goal of developing machines that possess the ability to execute tasks that typically necessitate human intelligence [14]. The aforementioned tasks encompass a range of cognitive skills, including but not limited to learning, adaptation, rationalization, comprehension of abstract concepts, and responsiveness to intricate human attributes such as attention, emotion, and creativity [15]. The primary purpose of the system was not to cater to healthcare needs, and its potential in addressing patient queries remains unexplored [16]. The reception of ChatGPT within the scientific community and academic circles has been varied, indicating the longstanding debate surrounding the advantages and drawbacks of sophisticated AI technologies [17-19]. ChatGPT has the potential to enhance conversational and writing abilities, thereby improving the effectiveness and precision of the desired outcome [6]. However, there have been concerns raised regarding potential bias stemming from the datasets utilized in the training of ChatGPT. This may restrict its abilities and lead to factual errors, which are disconcertingly scientifically feasible, a phenomenon known as hallucination [20].

Nonetheless, the comprehensive positive evaluation of dependability, as evidenced in our investigation assessing the responses proffered by ChatGPT concerning PAH, has similarly been corroborated in prior research endeavors exploring the precision of the responses produced by ChatGPT. Antaki et al. conducted a study to examine the precision of ChatGPT in the field of ophthalmology. The study employed a set of multiple-choice questions sourced from the Ophthalmic Knowledge Assessment Program [21]. The academic performance of ChatGPT was evaluated, and it was determined that the scores obtained on the exams were comparable to those of a typical first-year resident. The recorded percentages of 55.8% and 42.7% show that the results were noteworthy and promising [21]. Thus, this research has recognized the significance of patients' feedback. A further investigation was conducted to assess the precision and consistency of answers pertaining to knowledge, management, and emotional support regarding cirrhosis and hepatocellular carcinoma [22]. The study conducted by Duong and Solomon aimed to evaluate the efficacy of ChatGPT in addressing queries related to genetics in comparison to human participants [23]. The researchers arrived at the conclusion that ChatGPT offers prompt and precise answers to a diverse set of inquiries pertaining to genetics. They further noted that ChatGPT has the potential to facilitate the accessibility of information for laypersons [23]. In their research, Ziebland et al. claimed that severe illness frequently undermines people's perceptions of themselves as capable members of society [24]. Patients diagnosed with PAH can utilize AI technology to acquire knowledge and skills to showcase their proficiency in managing their condition amidst the challenges posed by the illness. Our study conducted on ChatGPT's responses and recommendations by

experts in the field of PAH concluded that they were deemed secure, beneficial, and devoid of any potential hazards.

Patients with a low educational level may struggle to understand data that has a high readability score. In addition to the quality of the information, health-related content should also aim to present easily understandable material [8]. In our study, when examining the response texts using two different readability scores, we found that the average FKGL value was 13.52 and the average SMOG value was 12.49. In a study examining the readability of websites for ankylosing spondylitis, the average FK and SMOG scores were 8.59 and 7.33, respectively. In another study evaluating the readability of websites for plantar fasciitis and calcaneal spur, the average scores were 7.27 for FKGL and 6.89 for SMOG [7,8]. This situation may make it more challenging for the general population to obtain information from ChatGPT compared to websites and could limit the information they receive.

This study has some limitations. First and foremost, in the stage of determining the questions, starting from the problems that the patients are curious about, although it enables the identification of a patient-oriented problem, it causes the identification to be subjective. Different research questions could have been based on different groups. The second is related to the fact that AI applications respond differently to questions at different times. Since this appearance may potentially jeopardize the reproducibility of the information, we kept the answer samples as images. Moreover, there are differences between the evaluators who check the accuracy of the information. In our methodology, there is no defined grading rubric or systematic approach for the evaluation process of ChatGPT responses; the ratings are based on the subjective assessments of medical experts actively working in this field. For this reason, we included a large number of observers' opinions.

Conclusions

In conclusion, ChatGPT demonstrates extensive expertise in the realm of healthcare, attributable to the utilization of expansive datasets during its training process. The veracity of the data furnished by ChatGPT is contingent upon an accurate comprehension and response to the inquiries posed. The reliability of health-related information furnished by ChatGPT is widely acknowledged by professionals in the field. Nonetheless, a precise diagnosis or treatment suggestion for any health issue necessitates an evaluation of the particular circumstances and a personalized methodology. Hence, it is not reasonable to anticipate that the data furnished by ChatGPT would serve as a substitute for the information dispensed by a medical practitioner or an expert in the field.

Appendices

Supplementary material 1



What is pulmonary arterial hypertension?



Pulmonary arterial hypertension (PAH) is a type of high blood pressure that affects the arteries in the lungs and the right side of the heart. In PAH, the blood vessels in the lungs become narrowed, stiff, or blocked, making it harder for blood to flow through them. This leads to an increase in blood pressure in the pulmonary artery, the main artery that carries blood from the heart to the lungs. As the pressure builds up, the right side of the heart must work harder to pump blood, which can eventually lead to right heart failure.

PAH can be caused by a variety of underlying conditions, such as connective tissue diseases, congenital heart defects, HIV infection, liver disease, and certain medications. In some cases, however, the cause is unknown. PAH is a serious condition that can cause symptoms such as shortness of breath, fatigue, chest pain, and swelling in the legs and ankles. Treatment options include medications to dilate blood vessels, reduce inflammation, and prevent blood clots, as well as oxygen therapy and lifestyle changes. In severe cases, a lung or heart-lung transplant may be necessary.

FIGURE 2: Question 1

PA What causes pulmonary arterial hypertension?



There are many possible causes of pulmonary arterial hypertension (PAH), and in many cases, the exact cause is not known. Some common underlying conditions that can lead to PAH include:

1. **Connective tissue diseases:** Certain autoimmune diseases, such as systemic lupus erythematosus (SLE) and scleroderma, can cause inflammation and damage to the blood vessels in the lungs.
2. **Congenital heart disease:** People born with certain types of heart defects may be more likely to develop PAH later in life.
3. **HIV infection:** HIV can cause inflammation and damage to the blood vessels in the lungs, leading to PAH.
4. **Liver disease:** People with liver disease, such as cirrhosis, may develop PAH due to increased blood flow to the lungs.
5. **Certain medications:** Some drugs, such as appetite suppressants and certain antidepressants, have been linked to PAH.
6. **Genetics:** In some cases, PAH may be caused by mutations in genes that control the growth and function of blood vessels in the lungs.
7. **Idiopathic:** In many cases, the cause of PAH is unknown and is referred to as idiopathic PAH.

It's important to note that while these conditions can increase the risk of developing PAH, not everyone with these conditions will develop PAH. Additionally, there may be other factors involved in the development of PAH, such as environmental factors or a combination of genetic and environmental factors.

FIGURE 3: Question 2

PA Is pulmonary hypertension the same as hypertension as we know it?



No, pulmonary hypertension (PH) is not the same as hypertension (also known as high blood pressure) that affects the systemic circulation. Pulmonary hypertension specifically refers to high blood pressure in the pulmonary artery and the smaller blood vessels in the lungs, while hypertension typically refers to high blood pressure in the arteries throughout the body.

Pulmonary hypertension can be classified into five different categories based on the underlying cause, including pulmonary arterial hypertension (PAH), which specifically refers to PH caused by narrowing or constriction of the pulmonary arteries. Other types of PH may be caused by lung diseases, heart diseases, blood clots, or other medical conditions.

While there may be some similarities in terms of symptoms and treatments, the underlying mechanisms and causes of pulmonary hypertension and hypertension are different. Pulmonary hypertension can be a serious and potentially life-threatening condition if left untreated, and early diagnosis and treatment are important for improving outcomes.

FIGURE 4: Question 3



What are the symptoms of pulmonary arterial hypertension?



The symptoms of pulmonary arterial hypertension (PAH) can be vague and nonspecific, which can make the condition difficult to diagnose in its early stages. Some common symptoms of PAH include:

1. Shortness of breath: This is often the first symptom of PAH and can occur during physical activity or even at rest.
2. Fatigue: People with PAH may feel tired or weak, even after minimal exertion.
3. Chest pain: This may feel like pressure or tightness in the chest, and can occur during physical activity or at rest.
4. Fainting or dizziness: This can occur due to decreased blood flow to the brain.
5. Swelling in the legs and ankles: This is often caused by fluid buildup (edema) due to increased pressure in the blood vessels.
6. Bluish lips or skin (cyanosis): This can occur due to decreased oxygen levels in the blood.
7. Racing heartbeat: This can occur due to the heart working harder to pump blood through the narrowed or stiff pulmonary arteries.

It's important to note that these symptoms can be caused by a variety of other conditions as well, and not everyone with PAH will experience all of these symptoms. If you are experiencing any of these symptoms or are concerned about your risk for PAH, it's important to talk to your healthcare provider for evaluation and diagnosis.

FIGURE 5: Question 4

PA Who has pulmonary arterial hypertension?



Pulmonary arterial hypertension (PAH) can affect people of all ages, genders, and races, but it is more commonly diagnosed in women than in men. According to the Pulmonary Hypertension Association, it is estimated that around 15 to 50 people per million have PAH, and it is more common in people over the age of 60. PAH can occur in people with certain medical conditions or risk factors, including:

1. Connective tissue diseases: PAH is commonly associated with connective tissue diseases such as scleroderma, lupus, and rheumatoid arthritis.
2. Congenital heart disease: People born with certain types of heart defects may be more likely to develop PAH later in life.
3. Liver disease: People with liver disease, such as cirrhosis, may develop PAH due to increased blood flow to the lungs.
4. HIV infection: HIV can cause inflammation and damage to the blood vessels in the lungs, leading to PAH.
5. Family history: In some cases, PAH may be inherited due to genetic mutations.
6. Obesity: People who are overweight or obese may be at an increased risk of developing PAH.
7. Certain medications: Some drugs, such as appetite suppressants and certain antidepressants, have been linked to PAH.

It's important to note that while these conditions can increase the risk of developing PAH, not everyone with these conditions will develop PAH. Additionally, there may be other factors involved in the development of PAH, such as environmental factors or a combination of genetic and environmental factors.

FIGURE 6: Question 5

PA Pregnancy in a patient with pulmonary arterial hypertension



Pregnancy can be a high-risk situation for women with pulmonary arterial hypertension (PAH). PAH can make it more difficult for the heart to pump blood to the lungs, and pregnancy places an additional strain on the heart and circulatory system. This can lead to complications such as heart failure, pulmonary edema, and even death. Additionally, some of the medications used to treat PAH may not be safe for use during pregnancy.

For these reasons, women with PAH are often advised against becoming pregnant or are counseled about the risks involved. Women with PAH who become pregnant should be closely monitored by a multidisciplinary team of healthcare providers, including specialists in cardiology, obstetrics, and anesthesia. They may also need to be treated with specialized medications and may require hospitalization for close monitoring during and after delivery.

In some cases, women with severe PAH may be advised against pregnancy altogether. The decision to become pregnant should be made in consultation with a healthcare provider who can assess the individual risk and provide appropriate guidance and treatment.

FIGURE 7: Question 6



How is pulmonary arterial hypertension diagnosed?



The diagnosis of pulmonary arterial hypertension (PAH) typically involves a combination of medical history, physical examination, and diagnostic tests. Some common diagnostic tests used to diagnose PAH include:

1. Echocardiogram: This test uses sound waves to create images of the heart and can help identify signs of PAH, such as an enlarged right ventricle.
2. Pulmonary function tests: These tests measure lung function and can help identify underlying lung diseases that may be contributing to PAH.
3. Chest X-ray: This imaging test can help identify any abnormalities in the lungs or heart.
4. CT scan or MRI: These imaging tests can provide more detailed images of the heart and lungs, helping to identify structural abnormalities and assess the severity of PAH.
5. Right heart catheterization: This invasive test involves inserting a catheter into a vein in the neck or groin and threading it into the pulmonary artery to measure blood pressure and assess the severity of PAH.
6. Blood tests: These tests can help identify underlying medical conditions that may be contributing to PAH, such as autoimmune diseases or liver disease.

It's important to note that PAH is a rare and complex condition that can be difficult to diagnose, and it may require referral to a specialized center for evaluation and treatment. A thorough diagnostic evaluation and early diagnosis are important for improving outcomes and quality of life for people with PAH.

FIGURE 8: Question 7

PA How is pulmonary arterial hypertension treated?



The treatment of pulmonary arterial hypertension (PAH) typically involves a combination of medications, lifestyle changes, and close monitoring. The goal of treatment is to improve symptoms, slow disease progression, and improve quality of life. Some common treatment options for PAH include:

1. **Medications:** There are several classes of medications used to treat PAH, including vasodilators, endothelin receptor antagonists, and phosphodiesterase-5 inhibitors. These medications work by relaxing the blood vessels in the lungs and improving blood flow, reducing symptoms and improving exercise capacity. The specific medications used may depend on the underlying cause of PAH and the severity of the condition.
2. **Oxygen therapy:** In some cases, supplemental oxygen may be needed to improve oxygen levels in the blood and reduce symptoms.
3. **Lifestyle changes:** Lifestyle changes such as maintaining a healthy weight, staying active, and avoiding smoking can help improve symptoms and slow disease progression.
4. **Surgery:** In rare cases, surgery may be recommended to improve blood flow and reduce pressure in the pulmonary arteries. Procedures such as pulmonary thromboendarterectomy (PTE) and lung transplant may be considered for certain individuals with PAH.
5. **Close monitoring:** Regular monitoring by a healthcare provider with experience in treating PAH is important for assessing disease progression and adjusting treatment as needed.

It's important to note that the treatment of PAH can be complex and may require referral to a specialized center for evaluation and management. The specific treatment approach may depend on several factors, including the underlying cause of PAH, the severity of the condition, and the individual's overall health status.

FIGURE 9: Question 8

Supplementary material 2

Survey questionnaire

How trustworthy is this information about pulmonary arterial hypertension provided by ChatGPT?



How valuable is the information provided by ChatGPT about pulmonary arterial hypertension for patients?



To what extent can the information on pulmonary arterial hypertension provided by ChatGPT be dangerous for patients?



FIGURE 10: Survey questionnaire

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Acquisition, analysis, or interpretation of data: Rasim Gökmen, Erhan Kaya, Murat Kerkütlüoğlu

Critical review of the manuscript for important intellectual content: Rasim Gökmen, Erhan Kaya, Murat Kerkütlüoğlu

Supervision: Erhan Kaya

Concept and design: Murat Kerkütlüoğlu

Drafting of the manuscript: Murat Kerkütlüoğlu

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. Kahramanmaraş Sütçü

İmam University Medical Research Ethics Committee issued approval 2023/15-02. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Van Bulck L, Moons P: What if your patient switches from Dr. Google to Dr. ChatGPT? A vignette-based survey of the trustworthiness, value, and danger of ChatGPT-generated responses to health questions. *Eur J Cardiovasc Nurs*. 2024, 23:95-8. [10.1093/eurjcn/zvad038](https://doi.org/10.1093/eurjcn/zvad038)
2. Lam-Po-Tang J, McKay D: Dr Google, MD: a survey of mental health-related internet use in a private practice sample. *Australas Psychiatry*. 2010, 18:130-3. [10.3109/10398560903473645](https://doi.org/10.3109/10398560903473645)
3. Ji Z, Lee N, Frieske R, et al.: Survey of hallucination in natural language generation. *ACM Comput Surv*. 2023, 55:1-38. [10.1145/3571730](https://doi.org/10.1145/3571730)
4. Moons P, Van Bulck L: ChatGPT: can artificial intelligence language models be of value for cardiovascular nurses and allied health professionals. *Eur J Cardiovasc Nurs*. 2023, 22:e55-9. [10.1093/eurjcn/zvad022](https://doi.org/10.1093/eurjcn/zvad022)
5. Hopkins AM, Logan JM, Kichenadasse G, Sorich MJ: Artificial intelligence chatbots will revolutionize how cancer patients access information: ChatGPT represents a paradigm-shift. *JNCI Cancer Spectr*. 2023, 7:pkad010. [10.1093/jncics/pkad010](https://doi.org/10.1093/jncics/pkad010)
6. Sallam M: ChatGPT utility in healthcare education, research, and practice: systematic review on the promising perspectives and valid concerns. *Healthcare (Basel)*. 2023, 11:[10.3390/healthcare11060887](https://doi.org/10.3390/healthcare11060887)
7. Kocyigit BF, Koca TT, Akaltun MS: Quality and readability of online information on ankylosing spondylitis. *Clin Rheumatol*. 2019, 38:3269-74. [10.1007/s10067-019-04706-y](https://doi.org/10.1007/s10067-019-04706-y)
8. Kaya E, Görmez S: Quality and readability of online information on plantar fasciitis and calcaneal spur. *Rheumatol Int*. 2022, 42:1965-72. [10.1007/s00296-022-05165-6](https://doi.org/10.1007/s00296-022-05165-6)
9. Humbert M, Lau EM, Montani D, Jais X, Sitbon O, Simonneau G: Advances in therapeutic interventions for patients with pulmonary arterial hypertension. *Circulation*. 2014, 130:2189-208. [10.1161/CIRCULATIONAHA.114.006974](https://doi.org/10.1161/CIRCULATIONAHA.114.006974)
10. Tonelli AR, Arelli V, Minai OA, Newman J, Bair N, Heresi GA, Dweik RA: Causes and circumstances of death in pulmonary arterial hypertension. *Am J Respir Crit Care Med*. 2013, 188:365-9. [10.1164/rccm.201209-1640OC](https://doi.org/10.1164/rccm.201209-1640OC)
11. Ling Y, Johnson MK, Kiely DG, et al.: Changing demographics, epidemiology, and survival of incident pulmonary arterial hypertension: results from the pulmonary hypertension registry of the United Kingdom and Ireland. *Am J Respir Crit Care Med*. 2012, 186:790-6. [10.1164/rccm.201205-0383OC](https://doi.org/10.1164/rccm.201205-0383OC)
12. Docs get clever with ChatGPT [Internet]. (2023). <https://www.medscape.com/viewarticle/987526?form=fpf>.
13. ChatGPT sets record for fastest-growing user base - analyst note [Internet]. (2023). <https://www.reuters.com/technology/chatgpt-sets-record-fastest-growing-user-base-analyst-note-2023-02-01/>.
14. Sarker IH: AI-based modeling: techniques, applications and research issues towards automation, intelligent and smart systems. *SN Comput Sci*. 2022, 3:158. [10.1007/s42979-022-01043-x](https://doi.org/10.1007/s42979-022-01043-x)
15. Korteling JE, van de Boer-Visschedijk GC, Blankendaal RA, Boonekamp RC, Eikelboom AR: Human versus artificial intelligence. *Front Artif Intell*. 2021, 4:622364. [10.3389/frai.2021.622364](https://doi.org/10.3389/frai.2021.622364)
16. Devlin J, Chang M-W, Lee K, Toutanova K: Bert: pre-training of deep bidirectional transformers for language understanding. *NAACL*. 2018, [10.48550/arXiv.1810.04805](https://arxiv.org/abs/1810.04805)
17. Wogu IAP, Olu-Owolabi FE, Assingbong PA, et al.: Artificial intelligence, alienation and ontological problems of other minds: a critical investigation into the future of man and machines. *ICCNI*. 2017, 1-10. [10.1109/ICCNI.2017.8123792](https://doi.org/10.1109/ICCNI.2017.8123792)
18. Howard J: Artificial intelligence: implications for the future of work. *Am J Ind Med*. 2019, 62:917-26. [10.1002/ajim.23037](https://doi.org/10.1002/ajim.23037)
19. Tai MC: The impact of artificial intelligence on human society and bioethics. *Tzu Chi Med J*. 2020, 32:339-43. [10.4103/tcmj.tcmj_71_20](https://doi.org/10.4103/tcmj.tcmj_71_20)
20. Deng J, Lin Y: The benefits and challenges of ChatGPT: an overview. *Front Comput Intell Syst*. 2022, 2:81-3. [10.54097/fcis.v2i2.4465](https://doi.org/10.54097/fcis.v2i2.4465)
21. Antaki F, Touma S, Milad D, El-Khoury J, Duval R: Evaluating the performance of ChatGPT in ophthalmology: an analysis of its successes and shortcomings. *Ophthalmol Sci*. 2023, 3:100324. [10.1016/j.xops.2023.100324](https://doi.org/10.1016/j.xops.2023.100324)
22. Yeo YH, Samaan JS, Ng WH, et al.: Assessing the performance of ChatGPT in answering questions regarding cirrhosis and hepatocellular carcinoma. *medRxiv*. 2023, 2023.02. [10.3350/cmh.2023.0089](https://doi.org/10.3350/cmh.2023.0089)
23. Duong D, Solomon BD: Analysis of large-language model versus human performance for genetics questions. *Eur J Hum Genet*. 2024, 32:466-8. [10.1038/s41431-023-01396-8](https://doi.org/10.1038/s41431-023-01396-8)
24. Ziebland S, Chapple A, Dumelow C, Evans J, Prinjha S, Rozmovits L: How the internet affects patients' experience of cancer: a qualitative study. *BMJ*. 2004, 328:564. [10.1136/bmj.328.7439.564](https://doi.org/10.1136/bmj.328.7439.564)