

Comparison of Decompression Alone Versus Decompression with Fusion for Stenotic Lumbar Spine: A Systematic Review and Meta-analysis

Syed Ijlal Ahmed ¹, Gohar Javed ², Syeda Beenish Bareeqa ³, Ali Shah ⁴, Maha Zubair ⁵, Rabbia Faisal. Avedia ⁶, Noor Rahman ⁶, Syeda Sana Samar ⁷, Kashif Aziz ⁸

1. Neurology, Liaquat National Hospital and Medical College, Karachi, PAK 2. Neurosurgery, The Aga Khan University, Karachi, PAK 3. Oncology, Jinnah Medical and Dental College, Karachi, PAK 4. Surgery, Dow University of Health Sciences, Karachi, PAK 5. Miscellaneous, Ziauddin Medical College, Karachi, PAK 6. Miscellaneous, Ziauddin Medical University, Karachi, PAK 7. Internal Medicine, Jinnah Sindh Medical University, Karachi, PAK 8. Internal Medicine, Icahn School of Medicine at Mount Sinai/Queens Hospital Center, New York City, USA

✉ **Corresponding author:** Kashif Aziz, kashifaziz927@gmail.com

Disclosures can be found in Additional Information at the end of the article

Abstract

The first line of treatment for lumbar spinal stenosis (with or without lumbar degenerative spondylolisthesis) involves conservative options such as anti-inflammatory drugs and analgesics. Approximately, 10%-15% of patients require surgery. Surgical treatment aims to decompress the spinal canal and dural sac from degenerative bony and ligamentous overgrowth. Different studies have given conflicting results. The aim of our study is to clear the confusion by comparing two surgical techniques. This meta-analysis was conducted in accordance with the preferred reporting items for systematic reviews and meta-analysis (PRISMA) guidelines. A literature search was conducted of the Ovid Embase, Scopus, Pubmed, Ovid Medline, Google Scholar, and Cochrane library databases. A quality and risk of bias assessment was also done. The analysis was done using Revman software (The Nordic Cochrane Centre, The Cochrane Collaboration, 2014, Copenhagen, Denmark).

A total of 76 studies were extracted from the literature search and 29 studies with relevant information were shortlisted. Nine studies were included in the meta-analysis after a quality assessment and eligibility. Fusion with decompression surgery was found to be a better technique when compared to decompression alone for spinal stenosis in terms of the Oswestry Disability index and the visual analog pain scale for back and leg pain. On the basis of the meta-analysis of the recent medical literature, the authors concluded that decompression with fusion is a 3.5-times better surgical technique than decompression alone for spinal stenosis.

Received 08/08/2018

Review began 08/10/2018

Review ended 08/10/2018

Published 08/13/2018

© Copyright 2018

Ahmed et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Categories: Neurology, Neurosurgery

Keywords: spinal stenosis, decompression surgery, fusion surgery, oswestry disability index

Introduction And Background

Lumbar spinal stenosis (LSS) is defined as the narrowing of the lumbar spinal canal and nerve root canals due to hypertrophic lesions of the facet joints, ligamentum flavum, and inter-vertebral discs, which leads to a debilitating compression of the spinal nerves and blood vessels [1-2]. It is one of the most prevalent spinal disorders in the elderly. Its incidence appears to be

How to cite this article

Ahmed S, Javed G, Bareeqa S, et al. (August 13, 2018) Comparison of Decompression Alone Versus Decompression with Fusion for Stenotic Lumbar Spine: A Systematic Review and Meta-analysis. Cureus 10(8): e3135. DOI 10.7759/cureus.3135

rising and it is estimated that by the year 2025, about 64-million elderly will be affected by it [3]. LSS often occurs in combination with lumbar degenerative spondylolisthesis (LDS), which, in turn, is defined as the slipping forward of one lumbar vertebra on another with an intact neural arch [4]. The first line of treatment for LSS (with or without LDS) involves conservative options, such as anti-inflammatory drugs and analgesics. Approximately 10%-15% of patients develop an incapacitating back and/or leg pain (BP and/or LP), which requires surgery [5]. Surgical treatment aims to decompress the spinal canal and dural sac from degenerative bony and ligamentous overgrowth [6]. Conventional surgical options include decompression (D) and decompression plus fusion (D+F) [7]. The aim of lumbar decompression is to decompress the neural elements while preserving stability [8]. If the lumbar spine is unstable, an instrumented fusion is performed in addition. The immobilization of one or more motion segments by fusion techniques involves joining two or more vertebrae permanently into one solid bone with no space between them [9].

Different studies have given conflicting results with some finding that decompression alone was associated with more favorable outcomes in spinal stenosis measure (SSM) scores and others demonstrating better outcomes with the added effect of fusion [6,10]. Evidence indicates that patients with lumbar spinal stenosis, without deformity or instability, treated with wide decompression, may suffer from iatrogenic lumbar instability [7].

A recent meta-analysis published by Chen et al compared D and D+F in patients with LDS. The study concluded that D+F did not yield better clinical outcomes than D alone [11]. Our study aims to assess the additive effect of fusion on decompression surgery and to ascertain if D+F is more effective than D alone in patients with LSS (with/without LDS).

Review

Materials and methods

Literature Search Strategy

We conducted this meta-analysis in accordance with the preferred reporting items for systematic reviews and meta-analysis (PRISMA) guidelines. A detailed literature search was conducted by two independent authors using the keywords “spinal fusion surgery,” “spinal decompression surgery,” “spinal stenosis,” “Oswestry Disability index,” and “degenerative spondylolisthesis” to search the Ovid Embase, Scopus, Pubmed, Ovid Medline, Google Scholar and Cochrane library databases. Relevant terms or synonyms other than keywords were utilized to conduct a comprehensive search in accordance with the pre-specified eligibility criteria. All the searched articles were exported and cited through Endnote. The search strategy was limited to medical literature in English, published from 2012 to 2018. In cases of an unavailability of the full text or incomplete data, the corresponding author was contacted.

Eligibility Criteria

The study types included in our research were randomized controlled trials and both prospective and retrospective cohort studies. However, case reports, letters to the editor, commentaries, cross-sectional surveys, and documentaries were excluded but used only to bridge and link the outcomes of our study with past medical research for discussion. Moreover, studies in non-English literature, studies which assessed the outcome in pathologies other than intracranial aneurysms, studies without definitive numbers or values, experimental animal trials, and studies with figurative or graphical results presentation without any particular numerical values were also excluded from this research. Two independent authors retrieved the required data in accordance with the mentioned eligibility criteria. Any disagreement was resolved by a collaborative discussion.

Data Collection

Studies were assessed and data were extracted by two independent reviewers according to PRISMA guidelines. Data were collected and compiled on a predefined evidence table. Articles were selected on the basis of relevance to the topic, appropriate sample size, sampling technique, and randomization. The collected data include author, year of publication, sample size, study design, Oswestry disability index (ODI), visual analog pain score (VAS) for back pain, VAS for leg pain, and the statistical results of the study (relative risk, confidence interval, and p-value). Any disagreement was resolved with a collaborative consensus.

Quality Assessment and Risk of Bias

To assess the quality of extracted data, the Newcastle-Ottawa scale was used for cohort studies and the Jadad scale (also known as the Oxford quality scoring system) was utilized for randomized controlled trials. To avoid any risk of bias, we evaluated data using the Cochrane risk of bias tool, which analyzed the data in terms of allocation concealment, blinding, sequence generation, selective outcome reporting, and incomplete outcome data.

Data Analysis and Primary Outcomes

The data was entered on RevMan version five (The Nordic Cochrane Centre, The Cochrane Collaboration, 2014, Copenhagen, Denmark) and analyzed using a forest plot for a visual estimation of the meta-analysis. The test for heterogeneity was also done (p-value <0.05). A fixed effect model with the inverse variance method was used to obtain the overall mean difference estimates and the 95% confidence interval (CI) in order to assess the effect of fusion surgery with decompression and decompression surgery alone on the Oswestry disability index and visual analog scale for back and leg pain. A p-value of less than 0.05 was considered significant.

Results*Extracted Studies Characteristics*

We selected 76 articles (which included cohorts, case-control studies, randomized trials, and reviews) on the basis of relevant titles and abstracts after a systematic review. After going through the abstracts of selected articles, 19 duplicates were excluded from the selected pool. The remaining 57 articles were screened for the required information. Another pool of 28 articles was removed after a screening of titles and abstracts. Removal was on the basis of non-English language literature and the unavailability of the full text of articles. The balance 29 full-text articles were assessed in accordance with the eligibility criteria out of which nine articles were finalized for quantitative synthesis [12-20].

Overall Outcomes

A total of 2929 patients were analyzed in the fusion with decompression group while 2779 patients were analyzed in the decompression group. Decompression with fusion was found to be 2.55 times better as compared to decompression alone in terms of the Oswestry Disability index (mean difference -1.73, CI 95%, {-3.05, -0.40}). The decompression with fusion group was also found 2.1 times (p=0.01) superior than the decompression alone group in terms of visual analog pain scale for back pain (mean difference -3.0, CI 95%, (-5.76, -0.24)) and 1.4 times (p=0.03) superior for leg pain (mean difference -2.35, CI 95%, (-2.35, 0.740)). The overall effect showed fusion with decompression as a 3.54 times better surgical technique over decompression alone for spinal stenosis (p=0.0004) (mean difference -2.02, CI 95%, (-3.13, -

0.90)). The detailed meta-analysis forest plot is shown in Figure 1. The heterogeneity test was also performed and the overall heterogeneity in the meta-analysis was only 16% (Chi-square=19.09, difference=16, p-value=0.14). The funnel plot for heterogeneity is shown in Figure 2.




FIGURE 1: Descriptive representation of the meta-analysis through a forest plot




FIGURE 2: Presentation of heterogeneity through a funnel plot

Discussion

Our meta-analysis compared the outcome of D versus D+F in terms of ODI, VAS (leg pain and back pain). Studies conducted in the past suggest that a radiographically detectable instability of the posterior elements of the spine require fusion surgery along with decompression for favorable postoperative outcomes [21-22]. Similarly, in our meta-analysis, we found fusion with decompression to be better than decompression alone in terms of ODI index and visual analog pain scale.

However, in contrast to our findings, a meta-analysis study showed that the addition of fusion did not result in improved clinical outcomes compared to D over a follow-up period of two years [23]. Similar to our findings, Li et al. concluded that D+F showed less favorable outcomes in terms of ODI, length of hospital stay, and blood loss in comparison with D alone (using coflex). They did not find any significant difference in VAS and major device-related complications, which is in contrast to our findings [24]. Another international study showed that both the D and D+F groups resulted in reduced VAS LP and BP [25]. Aihara et al. reported higher postoperative scores in the D alone group. However, their results were not statistically significant. Also, less blood loss and shorter post-operative hospitalization were observed in the D alone group [26].

Similar to our findings, high rates of satisfaction and decreased leg pain scores were observed in patients with lumbar degenerative spondylolisthesis, who underwent D+F rather than D alone [27]. However, another study suggested that the degree of satisfaction two years after surgery was slightly higher in patients who underwent D alone [28].

Conclusions

On the basis of a meta-analysis of the recent medical literature, authors concluded that decompression with fusion is a 3.5 times better surgical technique compared to decompression alone for spinal stenosis in terms of Oswestry disability index and visual analog pain scale for back pain and leg pain.

Additional Information

Disclosures

Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Deasy J: Acquired lumbar spinal stenosis. JAAPA. 2015, 28:19-23. [10.1097/01.JAA.0000462052.47882.fd](https://doi.org/10.1097/01.JAA.0000462052.47882.fd)
2. Oba H, Tsutsumimoto T, Yui M, Kamanaka T, Ohta H, Kosaku H, Misawa H: A prospective study of recovery from leg numbness following decompression surgery for lumbar spinal stenosis. J Orthop Sci. 2017, 22:670-675. [10.1016/j.jos.2017.04.004](https://doi.org/10.1016/j.jos.2017.04.004)
3. Wu AM, Tong TJ, Wang XY: A rethink of fusion surgery for lumbar spinal stenosis. J Evid Based Med. 2016, 9:166-169. [10.1111/jebm.12215](https://doi.org/10.1111/jebm.12215)
4. Weinstein JN, Lurie JD, Tosteson TD, et al.: Surgical versus nonsurgical treatment for lumbar

- degenerative spondylolisthesis. *N Engl J Med.* 2007, 356:2257-2270. [10.1056/NEJMoa070302](https://doi.org/10.1056/NEJMoa070302)
5. Kelleher MO, Timlin M, Persaud O, Rampersaud YR: Success and failure of minimally invasive decompression for focal lumbar spinal stenosis in patients with and without deformity. *Spine.* 2010, 35:E981-E987. [10.1097/BRS.0b013e3181c46fb4](https://doi.org/10.1097/BRS.0b013e3181c46fb4)
6. Ulrich NH, Burgstaller JM, Pichierri G, et al.: Decompression surgery alone versus decompression plus fusion in symptomatic lumbar spinal stenosis: a Swiss prospective multicenter cohort study with 3 years of follow-up. *Spine.* 2017, 17:S164. [10.1016/j.spinee.2017.07.245](https://doi.org/10.1016/j.spinee.2017.07.245)
7. Donnarumma P, Tarantino R, Nigro L, Rullo M, Messina D, Diacinti D, Delfini R: Decompression versus decompression and fusion for degenerative lumbar stenosis: analysis of the factors influencing the outcome of back pain and disability. *J Spine Surg.* 2016, 2:52-58. [10.21037/2Fjss.2016.03.07](https://doi.org/10.21037/2Fjss.2016.03.07)
8. Moisi M, Fisahn C, Tkachenko L, et al.: Unilateral laminotomy with bilateral spinal canal decompression for lumbar stenosis: a technical note. *Cureus.* 2016, 8:e623. [10.7759/2Fcureus.623](https://doi.org/10.7759/2Fcureus.623)
9. Kalff R, Ewald C, Waschke A, Gobisch L, Hopf C: Degenerative lumbar spinal stenosis in older people: current treatment options. *Dtsch Arztebl Int.* 2013, 110:613-624. [10.3238/2Farztebl.2013.0613](https://doi.org/10.3238/2Farztebl.2013.0613)
10. Ghogawala Z, Benzel EC, Amin-Hanjani S, et al.: Prospective outcomes evaluation after decompression with or without instrumented fusion for lumbar stenosis and degenerative Grade I spondylolisthesis. *J Neurosurg Spine.* 2004, 1:267-272. [10.3171/spi.2004.1.3.0267](https://doi.org/10.3171/spi.2004.1.3.0267)
11. Chen Z, Xie P, Feng F, Chhantyal K, Yang Y, Rong L: Decompression alone versus decompression and fusion for lumbar degenerative spondylolisthesis: a meta-analysis. *World Neurosurg.* 2017, 111:e165-e177. [10.1016/j.wneu.2017.12.009](https://doi.org/10.1016/j.wneu.2017.12.009)
12. Austevoll IM, Gjestad R, Brox JI, et al.: The effectiveness of decompression alone compared with additional fusion for lumbar spinal stenosis with degenerative spondylolisthesis: a pragmatic comparative non-inferiority observational study from the Norwegian Registry for Spine Surgery. *Eur Spine J.* 2017, 26:404-413. [10.1007/s00586-016-4683-1](https://doi.org/10.1007/s00586-016-4683-1)
13. Davis R, Auerbach JD, Bae H, Errico TJ: Can low-grade spondylolisthesis be effectively treated by either coflex interlaminar stabilization or laminectomy and posterior spinal fusion? Two-year clinical and radiographic results from the randomized, prospective, multicenter US investigational device exemption trial. *J Neurosurg Spine.* 2013, 19:174-184. [10.3171/2013.4.spine12636](https://doi.org/10.3171/2013.4.spine12636)
14. Forsth P, Olafsson G, Carlsson T, et al.: A randomized, controlled trial of fusion surgery for lumbar spinal stenosis. *N Engl J Med.* 2016, 374:1413-1423. [10.1056/NEJMoa1513721](https://doi.org/10.1056/NEJMoa1513721)
15. Forsth P, Michaelsson K, Sanden B: Does fusion improve the outcome after decompressive surgery for lumbar spinal stenosis?: A two-year follow-up study involving 5390 patients. *Bone Joint J.* 2013, 97B:960-965. [10.1302/0301-620X.95B7.30776](https://doi.org/10.1302/0301-620X.95B7.30776)
16. Ghogawala Z, Dziura J, Butler WE, et al.: Laminectomy plus fusion versus laminectomy alone for lumbar spondylolisthesis. *N Engl J Med.* 2016, 374:1424-1434. [10.1056/NEJMoa1508788](https://doi.org/10.1056/NEJMoa1508788)
17. Postacchini F, Postacchini R, Menchetti PP, Sessa P, Paolino M, Cinotti G: Lumbar interspinous process fixation and fusion with stand-alone interlaminar lumbar instrumented fusion implant in patients with degenerative spondylolisthesis undergoing decompression for spinal stenosis. *Asian Spine J.* 2016, 10:27-37. [10.4184/asj.2016.10.1.27](https://doi.org/10.4184/asj.2016.10.1.27)
18. Sigmundsson FG, Jönsson B, Strömquist B: Outcome of decompression with and without fusion in spinal stenosis with degenerative spondylolisthesis in relation to preoperative pain pattern: a register study of 1,624 patients. *Spine J.* 2015, 15:638-646. [10.1016/j.spinee.2014.11.020](https://doi.org/10.1016/j.spinee.2014.11.020)
19. Truszczyńska A, Rapała K, Łukawski S, Trzaskoma Z, Tarnowski A, Drzał-Grabiec J, Cabak A: Evaluation of functional outcomes in individuals 10 years after posterior lumbar interbody fusion with corundum implants and decompression: a comparison of 2 surgical techniques. *Med Sci Monit.* 2016, 20:1400-1406. [10.12659/2FMSM.890545](https://doi.org/10.12659/2FMSM.890545)
20. Yagi M, Fujita N, Okada E, et al.: Comparisons of direct costs, outcomes, and cost-utility of decompression surgery with fusion versus decompression alone for degenerative lumbar spondylolisthesis. *J Orthop Sci.* 2018, 23:653-657. [10.1016/j.jos.2018.04.001](https://doi.org/10.1016/j.jos.2018.04.001)
21. Grob D, Humke T, Dvorak J: Degenerative lumbar spinal stenosis. Decompression with and without arthrodesis. *J Bone Joint Surg Am.* 1995, 77:1036-1041.
22. Rompe JD, Eysel P, Zöllner J, Nafe B, Heine J: Degenerative lumbar spinal stenosis. *Neurosurg*

- Rev. 1999, 22:102-106. [10.1007/s101430050040](https://doi.org/10.1007/s101430050040)
23. Chang W, Yuwen P, Zhu Y, Wei N, Feng C, Zhang Y, Chen W: Effectiveness of decompression alone versus decompression plus fusion for lumbar spinal stenosis: a systematic review and meta-analysis. *Arch Orthop Trauma Surg.* 2017, 137:637-650. [10.1007/s00402-017-2685-z](https://doi.org/10.1007/s00402-017-2685-z)
 24. Li AM, Li X, Yang Z: Decompression and coflex interlaminar stabilisation compared with conventional surgical procedures for lumbar spinal stenosis: a systematic review and meta-analysis. *Int J Surg.* 2017, 40:60-67. [10.1016/j.ijssu.2017.02.056](https://doi.org/10.1016/j.ijssu.2017.02.056)
 25. Kim HJ, Park JH, Kim JW, Kang KT, Chang BS, Lee CK, Yeom JS: Prediction of postoperative pain intensity after lumbar spinal surgery using pain sensitivity and preoperative back pain severity. *Pain Med.* 2014, 15:2037-2045. [10.1111/pme.12578](https://doi.org/10.1111/pme.12578)
 26. T Aihara, T Toyone, Y Aoki, T Ozawa, G Inoue, K Hatakeyama, J Ouchi: Surgical management of degenerative lumbar spondylolisthesis: a comparative study of outcomes following decompression with fusion and microendoscopic decompression. *J Musculoskeletal Research.* 2012, 15:1250020. [10.1142/S0218957712500200](https://doi.org/10.1142/S0218957712500200)
 27. Liang HF, Liu SH, Chen ZX, Fei QM: Decompression plus fusion versus decompression alone for degenerative lumbar spondylolisthesis: a systematic review and meta-analysis. *Eur Spine J.* 2017, 26:3084-3095. [10.1007/s00586-017-5200-x](https://doi.org/10.1007/s00586-017-5200-x)
 28. Matsudaira K, Yamazaki T, Seichi A, Takeshita K, Hoshi K, Kishimoto J, Nakamura K: Spinal stenosis in grade I degenerative lumbar spondylolisthesis: a comparative study of outcomes following laminoplasty and laminectomy with instrumented spinal fusion. *J Orthop Sci.* 2005, 10:270-276. [10.1007/s00776-005-0887-7](https://doi.org/10.1007/s00776-005-0887-7)