Cureus
Part of SPRINGER NATURE

Open Access Review Article

From the Triangular Fibrocartilage to the Foveal Concept: The Historical Evolution of Triangular Fibrocartilage Complex Anatomy, Diagnosis, and Repair

Review began 10/28/2026
Review ended 11/06/2026
Published 11/10/2026

© **Copyright** 2026
Ergin et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

DOI: 10.7759/cureus.119428

Musa Ergin [1], Mehmet A. Acar [1]

1. Department of Orthopaedics and Traumatology, Akademi Meram Hospital, Konya, TUR

Corresponding author: Musa Ergin, drmusaergin@gmail.com

Abstract

This review aims to trace the historical evolution of the triangular fibrocartilage complex (TFCC), from early descriptions of wrist trauma and triangular fibrocartilage to the contemporary understanding of the foveal attachment as a critical determinant of distal radioulnar joint (DRUJ) stability. A comprehensive review of medical and historical literature was conducted, including classical medical writings, historical anatomical texts, and landmark anatomical, biomechanical, diagnostic, arthroscopic, and surgical studies. Significant contributions were analyzed chronologically and within their scientific contexts, identifying key conceptual shifts that shaped the modern understanding and treatment of TFCC pathology. The history of TFCC reflects a shift from anatomical description to functional interpretation. Early descriptions of wrist injuries gradually evolved into a more comprehensive understanding of the relevant soft-tissue anatomy. Subsequent anatomical and biomechanical advances established the TFCC as a functional complex involved in ulnocarpal load transfer and DRUJ stability. The development of classification systems provided a morphological framework for traumatic and degenerative lesions, while advances in wrist arthroscopy made it possible to directly and dynamically assess TFCC pathology. Further anatomical and histological investigations refined the understanding of the stabilizing role of the foveal attachment, shifting clinical attention from the tear site to functional instability. This shift redirected surgical treatment from debridement and peripheral suturing to anatomical foveal reattachment and reconstruction in irreparable lesions. The evolution of TFCC knowledge can be understood through three main transitional pathways: from structure to complex, from morphology to function, and from tear repair to stability restoration. The modern foveal concept represents the culmination of successive advances in anatomy, biomechanics, arthroscopy, imaging, and surgery, and provides the conceptual basis for the contemporary management of TFCC injuries and DRUJ instability.

Categories: Orthopedics
Keywords: arthroscopic wrist surgery, distal radioulnar joint (druj), foveal attachment, historical review, tfcc, triangular fibrocartilage complex injury

Introduction And Background

Ulnar-sided wrist pain remains a diagnostic and therapeutic challenge in hand and wrist surgery because of the complex anatomy and functional interdependence of the structures surrounding the distal radioulnar joint (DRUJ) [1,2]. The triangular fibrocartilage complex (TFCC) plays an essential role in ulnocarpal load transmission and DRUJ stability [3,4]. However, the contemporary understanding of the TFCC as an integrated anatomical and functional complex has emerged through successive advances in anatomy, biomechanics, diagnostic techniques, and surgical treatment.

The TFCC comprises the articular disc, dorsal and palmar radioulnar ligaments, meniscus homologue, ulnolunate and ulnotriquetral ligaments, and associated ulnar-sided capsuloligamentous structures. Among these components, the radioulnar ligaments are particularly important for DRUJ stability. Their superficial fibers attach near the ulnar styloid, whereas their deep fibers insert into the ulnar fovea and provide essential restraint to DRUJ translation during forearm rotation [3-5]. This anatomical distinction has important clinical implications. Peripheral TFCC tears with preserved foveal attachment may occur without significant DRUJ instability, whereas disruption of the deep foveal fibers can compromise joint stability. Consequently, tear location alone may not adequately reflect the functional significance of a TFCC injury.

Historically, the structures now recognized as components of the TFCC were not initially understood as a unified functional system [6,7]. Early medical writings described wrist trauma without identifying the complex soft-tissue anatomy involved. Subsequent anatomical investigations progressively characterized the articular disc and its surrounding ligamentous attachments, establishing the morphological foundations of modern TFCC anatomy [6,7]. A major conceptual advancement occurred in 1981 when Palmer and Werner introduced the TFCC as an integrated anatomical and functional complex, emphasizing its contribution to ulnocarpal load transmission and DRUJ stability [8]. Palmer's subsequent classification provided a

systematic framework for distinguishing traumatic and degenerative TFCC lesions [9,10].

The development of wrist arthroscopy, together with increasingly detailed anatomical and histological investigations, further refined the understanding of TFCC injury patterns [11-13]. Direct visualization and probing enabled more precise assessment of peripheral and foveal lesions, while investigations of the deep radioulnar ligament attachments highlighted the importance of foveal integrity in DRUJ stability [12-14]. These developments expanded clinical decision-making beyond the location of a visible tear to include ligament attachment integrity, functional instability, and tissue reparability. Accordingly, arthroscopic debridement, peripheral repair, anatomical foveal reattachment, and ligament reconstruction developed as complementary, lesion-specific treatment strategies rather than successive replacements for one another [15,16].

Although numerous reviews have examined TFCC anatomy, imaging, classification, and surgical management, these developments have frequently been addressed as separate anatomical, technical, or clinical subjects [17-19]. Their integration into a coherent historical narrative remains relatively limited. Understanding the historical evolution of TFCC concepts may help clarify contemporary debates regarding terminology, classification, diagnostic assessment, and surgical decision-making [6,20] (Table *1*).

Period/Year	Contributor (s)	Landmark contribution	Conceptual significance
Classical antiquity	Hippocrates [21]	Early descriptions of wrist trauma and joint reduction	Clinical observation preceding modern anatomical characterization
1975	Kauer [22]	Anatomical investigation of the development of the triangular ligament in humans	Early developmental characterization of the triangular ligament
1978	Mikić [23]	Description of age-related degenerative changes in the triangular fibrocartilage	Recognition of degeneration as a distinct pathological process
1981	Palmer and Werner [8]	Characterization of the TFCC as an integrated anatomical and functional complex	Transition from an isolated anatomical structure to a functional complex
1985	Kuhlmann et al. [24]	Anatomical investigation of the triangular ligament and medial wrist compartment	Further characterization of the ulnar-sided wrist anatomy
1986–1988	Whipple et al. [25]; Roth et al. [26]	Development and clinical application of wrist arthroscopy	Introduction of direct intra-articular visualization and assessment
1989	Palmer [9]	Introduction of the classification of traumatic (Class 1) and degenerative (Class 2) TFCC lesions	Establishment of a systematic morphological classification
1996–2001	Nakamura et al. [11-13]	Functional and histological characterization of TFCC anatomy, origins, and insertions	Transition from morphological description toward functional understanding
2008–2011	Atzei et al. [15,20]; Atzei and Luchetti [27]	Development of arthroscopic foveal repair strategies and classification of foveal TFCC tears in relation to DRUJ instability	Recognition of foveal integrity as a key determinant of DRUJ stability
2014	Bain et al. [16]	Description of anatomical foveal TFCC reconstruction using a tendon graft	Expansion of treatment from primary repair to anatomical reconstruction
2017–2018	Chen [28]; Park et al. [29]	Description of arthroscopically assisted transosseous foveal repair techniques	Development of minimally invasive techniques for restoring foveal attachment
2024	Ma et al. [30]	Systematic review comparing suture-anchor and transosseous suture techniques for arthroscopic foveal repair	Comparative evaluation of contemporary foveal repair strategies

TABLE 1: Historical milestones in the evolution of triangular fibrocartilage complex anatomy, diagnosis, and surgical management.

TFCC: triangular fibrocartilage complex; DRUJ: distal radioulnar joint

The aim of this historical narrative review is to trace the evolution of TFCC knowledge from early anatomical

descriptions to the contemporary understanding of foveal attachment and DRUJ stability. Particular emphasis is placed on three interconnected conceptual transitions: from an isolated anatomical structure to a functional complex, from morphological description to biomechanical interpretation, and from tear-oriented treatment toward lesion-specific management and restoration of stability.

Review

Methodology

This historical narrative review aimed to identify and contextualize the major developments that shaped the contemporary understanding of the TFCC, with particular emphasis on anatomy, biomechanics, diagnosis, classification, arthroscopy, and surgical treatment.

PubMed/MEDLINE and Google Scholar were searched from database inception to May 23, 2026, using combinations of the terms “triangular fibrocartilage,” “TFCC,” “joint disc,” “triangular ligament,” “distal radioulnar joint,” “radioulnar ligament,” “foveal attachment,” “wrist arthroscopy,” “TFCC classification,” and “TFCC repair.” Reference lists of relevant publications were also examined to identify additional historically significant sources.

Publications were selected according to their historical and conceptual significance, with priority given to original studies that introduced or substantially advanced anatomical descriptions, biomechanical concepts, diagnostic approaches, classification systems, arthroscopic techniques, or surgical treatment strategies. Subsequent studies were included when they clarified or expanded upon these landmark contributions. The relevance of identified publications was assessed by multiple authors. No restrictions were imposed regarding publication year or study design. The full texts of included publications were reviewed, including translated versions of relevant non-English historical sources.

The selected literature was synthesized chronologically and thematically to identify three major conceptual transitions: from an isolated anatomical structure to a functional complex, from morphological description to biomechanical interpretation, and from tear-oriented treatment toward lesion-specific management and restoration of stability.

Given the historical narrative design, no quantitative meta-analysis or formal risk-of-bias assessment was performed. Anatomical and cadaveric findings were distinguished from clinical evidence, and the methodological limitations of diagnostic and therapeutic studies were considered when interpreting their findings.

Before the triangular fibrocartilage complex: from ancient wrist trauma to the triangular fibrocartilage

The historical understanding of TFCC evolved gradually from descriptions of wrist trauma to detailed anatomical and pathological investigations. Classical medical writings, including the Hippocratic Corpus, addressed wrist injuries but did not identify the specific soft-tissue structures or injury patterns recognized in contemporary TFCC pathology [21].

Early anatomical investigations focused primarily on the articular disc and triangular ligament rather than an integrated functional complex. Kauer’s developmental study in 1975 examined the relationship between the triangular ligament, ulnar styloid, and adjacent carpal structures [22]. Subsequently, Kuhlmann and colleagues emphasized the anatomical and functional relationships of the triangular ligament within the ulnar-sided wrist [24]. These studies contributed to the anatomical foundations of the modern TFCC concept without establishing its complete biomechanical significance.

An important pathological contribution was made by Mikić in 1978, who examined 180 wrist joints from 100 cadavers and demonstrated progressive age-related degeneration of the triangular fibrocartilage. Disc perforations were absent during the first two decades of life but increased substantially with advancing age [23]. Subsequent anatomical and radiological investigations further characterized the triangular fibrocartilage and surrounding ulnocarpal structures [31].

Collectively, these findings established that structural abnormalities of the triangular fibrocartilage may reflect degenerative processes rather than acute traumatic injury. However, the integrated biomechanical role of the articular disc and its surrounding ligamentous attachments remained incompletely understood. This limitation provided the foundation for the subsequent development of the modern TFCC concept.

From fibrocartilage to complex: the Palmer-Werner era

The introduction of the TFCC concept by Palmer and Werner in 1981 marked a pivotal transition from descriptive anatomy to functional interpretation. Through anatomical dissection and biomechanical testing of 61 specimens, they characterized the articular disc and surrounding ulnar-sided soft tissues as an integrated functional unit rather than an isolated fibrocartilaginous structure. Their findings highlighted the

contribution of the TFCC to ulnocarpal load transmission and DRUJ stability [8].

Subsequent biomechanical investigations further clarified this functional relationship. Biomechanical investigations demonstrated that distal radioulnar alignment influences load distribution across the ulnar wrist [8]. Additional research identified an inverse relationship between positive ulnar variance and articular disc thickness [32], while cadaveric experiments demonstrated that wrist and forearm positioning influenced load transmission through the radius and ulna [33]. Collectively, these anatomical and experimental findings established a biomechanical framework linking TFCC morphology, ulnar variance, and ulnocarpal loading, although their direct clinical implications required further investigation (Table 2).

Year	Landmark study	Principal contribution	Conceptual advance	Lasting clinical significance
1978	Mikić [23]	Demonstrated progressive age-related degeneration and perforation of the triangular fibrocartilage	Structural defect → degenerative pathology	Established that structural abnormalities may represent degeneration rather than acute trauma
1981	Palmer and Werner [8]	Defined the TFCC as an integrated anatomical and functional unit	Structure → complex	Established the foundation of the modern TFCC concept
1989	Palmer [9]	Introduced a systematic classification of traumatic and degenerative TFCC lesions	Complex → clinical morphology	Provided a common language for diagnosis and treatment
1996	Nakamura et al. [12]	Distinguished distal, proximal, and ulnar components and identified the foveal proximal component as the functional radioulnar ligament	Morphology → function	Linked the deep foveal attachment to DRUJ stabilization
2001	Nakamura et al. [13]	Histologically characterized the foveal and basistyloid insertions of the radioulnar ligament	Function → anatomical substrate	Provided the anatomical basis for foveal reattachment
2008	Atzei et al. [15]	Introduced a treatment-oriented framework for peripheral TFCC tears and described arthroscopic foveal reattachment	Visible tear → functional lesion	Integrated foveal integrity, DRUJ instability, and repair strategy
2015	Atzei et al. [34]	Reported all-inside arthroscopic foveal repair using direct DRUJ arthroscopy and suture-anchor fixation	Peripheral repair → anatomical reattachment	Demonstrated restoration of DRUJ stability through targeted foveal repair

TABLE 2: Landmark studies shaping the contemporary understanding and treatment of the TFCC.

TFCC: triangular fibrocartilage complex; DRUJ: distal radioulnar joint

The next significant development was the transformation of this framework into a clinically applicable classification. In 1989, Palmer divided TFCC lesions into two categories: Class 1 traumatic and Class 2 degenerative abnormalities; traumatic lesions were further classified according to their anatomical location [9]. This morphology-based system provided a common language for TFCC pathology and profoundly influenced subsequent diagnosis and treatment. The biological significance of lesion localization has been reinforced by microvascular studies showing that TFCC is predominantly peripheral, while its central portion is essentially devascularized [35].

Arthroscopy and the rise of the foveal concept

The development of wrist arthroscopy marked a major advancement in the understanding of TFCC pathology. During the 1980s, improvements in small-joint instrumentation and arthroscopic techniques facilitated systematic examination of the wrist. Whipple and colleagues described reproducible approaches to wrist arthroscopy, while Roth, Poehling, and Whipple further contributed to its development as a diagnostic and therapeutic procedure [25,26].

Beyond direct visualization, arthroscopy enabled dynamic assessment of TFCC integrity through probing and evaluation of tissue tension. Increasing anatomical knowledge revealed the functional distinction between superficial peripheral attachments and the deep radioulnar ligament fibers inserting into the ulnar fovea. Because these deep fibers contribute substantially to DRUJ stability, apparently similar peripheral tears may have different functional consequences depending on foveal attachment integrity [14,27].

Atzei and Luchetti incorporated this anatomical distinction into the "iceberg" concept, emphasizing that a superficial peripheral lesion visible during radiocarpal arthroscopy may conceal a deeper foveal detachment [27]. Arthroscopic examination subsequently evolved to include complementary diagnostic maneuvers. The trampoline test primarily assesses TFCC tension, whereas the hook test evaluates the integrity of its deep foveal attachment. A clinical study and a cadaveric assessment reported better diagnostic performance of the hook test than the trampoline test for identifying foveal detachment; the cadaveric findings do not, by themselves, establish clinical outcomes [36,37].

Thus, wrist arthroscopy transformed TFCC assessment from predominantly morphological observation toward functional evaluation. The central diagnostic question expanded beyond tear localization to include foveal attachment integrity and DRUJ stability. This conceptual shift established an important foundation for contemporary lesion-specific TFCC management.

Evolution of diagnosis and repair: from tear detection to restoration of stability

The evolution of TFCC diagnosis and treatment has closely paralleled advances in anatomical and biomechanical understanding. Before cross-sectional imaging and arthroscopy, diagnosis relied primarily on clinical examination and conventional radiography. Wrist arthrography subsequently enabled the detection of TFCC perforations through contrast communication between the radiocarpal and distal radioulnar joints, although its ability to characterize tear morphology and functional instability remained limited [38]. Magnetic resonance imaging (MRI) and magnetic resonance arthrography (MRA) subsequently improved the assessment of TFCC injuries. Although these modalities have enhanced the detection of structural abnormalities, particularly with MRA, imaging findings do not directly establish the mechanical stability of the DRUJ [39].

Surgical management evolved in parallel with these diagnostic advances. Arthroscopic debridement provided symptomatic improvement in selected TFCC lesions, as supported by subsequent clinical studies [40,41]. Recognition of regional vascularity and healing potential encouraged repair of suitable peripheral tears rather than routine excision. However, peripheral repair alone may be insufficient when the deep foveal attachment is disrupted, and DRUJ instability persists [14,20]. Thus, debridement, peripheral repair, and foveal reattachment address different injury patterns rather than representing interchangeable treatments.

The recognition of foveal attachment integrity shifted the surgical objective toward restoration of the stabilizing radioulnar ligament insertion. Arthroscopic foveal repair can be performed using transosseous sutures or suture-anchor fixation. Transosseous techniques have been described in technical and clinical reports [28,29], while a systematic review of 26 studies involving 904 patients reported improvements in pain and functional outcomes after both fixation approaches, with follow-up ranging from 6 to 106 months [30]. However, much of the available evidence is derived from observational studies with heterogeneous patient populations, surgical techniques, and follow-up periods. Consequently, the absence of demonstrated superiority of one technique should not be interpreted as evidence of equivalent effectiveness. When the residual tissue is irreparable, ligament reconstruction may be considered to restore DRUJ stability.

These developments illustrate a transition from detecting structural lesions toward evaluating their functional consequences. Contemporary TFCC management therefore requires integration of tear morphology, foveal attachment integrity, tissue reparability, and DRUJ stability. The therapeutic objective is not simply closure of a visible tear, but restoration of a stable and functional wrist.

The modern triangular fibrocartilage complex: lessons from history and unresolved questions

Despite substantial advances in anatomy, imaging, arthroscopy, and surgical repair, the understanding of TFCC continues to evolve. Anatomical boundaries and terminology remain incompletely standardized. A comprehensive review identified inconsistencies in the definitions of TFCC components, particularly the radioulnar ligaments, ulnar attachments, and ulnar collateral ligament, emphasizing the need for standardized anatomical nomenclature [42]. Similar challenges persist in classification. Although the Palmer system remains useful for describing tear morphology, contemporary classification approaches increasingly consider additional anatomical and functional characteristics. Combined tear patterns that do not fit neatly into conventional categories further illustrate the limitations of classification based solely on lesion location [43,44].

Another unresolved issue concerns the relationship between structural abnormalities and clinical symptoms. In an MRI study of 103 asymptomatic volunteers, TFCC abnormalities were identified in 39 wrists, including 23 complete tears [45]. A subsequent pooled analysis demonstrated that the prevalence of TFCC abnormalities increases with age, irrespective of symptoms [46]. These findings emphasize that imaging abnormalities should be interpreted alongside clinical examination. Furthermore, recent diagnostic research suggests that MRI may identify TFCC injuries more reliably than it characterizes the precise location and extent of foveal involvement [47].

The same distinction applies to treatment. The foveal concept advanced TFCC surgery by emphasizing restoration of the stabilizing attachment rather than closure of the visible tear alone. Nevertheless, foveal repair is indicated for selected lesions and does not replace debridement or peripheral repair in other tear patterns. DRUJ stability depends on the interaction of intrinsic and extrinsic stabilizers, including the TFCC, interosseous membrane, pronator quadratus, osseous morphology, and surrounding soft tissues [48]. Recent evidence suggesting that patient-specific factors beyond the repaired TFCC may influence recurrent instability further supports this broader functional perspective [49].

Historically, the central lesson is not that one anatomical model replaces another, but that successive advances expand the understanding of TFCC function and pathology. Knowledge has progressed from isolated structure to functional complex, from morphology to biomechanics, and from tear-oriented treatment toward restoration of stability (Figure *1*).




FIGURE 1: Paradigm shift in the understanding and treatment of the triangular fibrocartilage complex.

TFCC: triangular fibrocartilage complex; DRUJ: distal radioulnar joint

Future developments in dynamic imaging, biomechanical assessment, and individualized treatment strategies may further refine this evolving framework.

Limitations, recommendations, and future perspectives

This historical narrative review provides an integrated perspective on the evolution of TFCC anatomy, diagnosis, and surgical management. However, several limitations should be acknowledged. First, its narrative design does not permit quantitative comparisons or systematic assessment of the strength of evidence. Furthermore, the limited accessibility of early anatomical sources, changes in terminology, and differences between historical and contemporary research methodologies complicate the interpretation of certain developments. Finally, heterogeneity in lesion classification, assessment of DRUJ instability, surgical techniques, study designs, and follow-up duration limits definitive conclusions regarding the comparative effectiveness of different treatment approaches.

Current evidence suggests that TFCC pathology should not be evaluated solely based on tear morphology. Foveal attachment integrity, DRUJ stability, concomitant intra-articular lesions, tissue reparability, and patient-specific characteristics should also be considered. Future studies should therefore adopt reproducible classification systems incorporating both anatomical and functional findings. Correlation of imaging abnormalities with clinical examination and, when indicated, arthroscopic assessment remains essential for distinguishing incidental degenerative changes from clinically significant lesions. Surgical outcomes should be evaluated using standardized measures of DRUJ stability, pain, patient-reported function, complications, and long-term clinical results.

Future research should further clarify the relationship between TFCC anatomy, biomechanical function, and clinical outcomes. Well-designed multicenter prospective studies with adequate sample sizes, standardized outcome measures, and long-term follow-up are needed to compare transosseous foveal repair, suture-anchor fixation, and anatomical reconstruction in appropriately selected patient populations. Such investigations may help establish more reliable, lesion-specific treatment strategies and strengthen the evidence supporting restoration of DRUJ stability as a central objective of contemporary TFCC surgery.

Conclusions

The historical evolution of TFCC reflects a transition from anatomical description to functional understanding. Once regarded primarily as an isolated fibrocartilaginous structure, the TFCC is now recognized as an integrated anatomical and biomechanical complex, with the foveal attachment playing a critical role in DRUJ stability. This evolving understanding has reshaped diagnostic and therapeutic approaches, shifting attention beyond tear morphology toward functional integrity and lesion-specific management. Three major conceptual transitions characterize this progression: from structure to complex, from morphology to function, and from tear-oriented treatment toward restoration of stability. These principles continue to guide contemporary TFCC assessment and treatment while providing a foundation for future anatomical, biomechanical, and clinical research.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Musa Ergin, Mehmet A. Acar

Acquisition, analysis, or interpretation of data: Musa Ergin, Mehmet A. Acar

Drafting of the manuscript: Musa Ergin, Mehmet A. Acar

Critical review of the manuscript for important intellectual content: Musa Ergin, Mehmet A. Acar

Supervision: Mehmet A. Acar

Disclosures

Conflicts of interest: In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

ChatGPT (OpenAI) was used solely for English-language editing and grammatical refinement of the manuscript. The authors reviewed and approved all AI-assisted edits and take full responsibility for the final content.

References

1. Ou Yang O, McCombe DB, Keating C, Maloney PP, Berger AC, Tham SK: Ulnar-sided wrist pain: a prospective analysis of diagnostic clinical tests. ANZ J Surg. 2021, 91:2159-62. 10.1111/ans.17169
2. Jain DK, Wahegaonkar AL: Ulnar-side wrist pain management guidelines: all that hurts is not the TFCC! . Indian J Orthop. 2021, 55:310-7. 10.1007/s43465-020-00319-9
3. Pidgeon TS, Waryasz G, Carnevale J, DaSilva MF: Triangular fibrocartilage complex: an anatomic review . JBJS Rev. 2015, 3:1. 10.2106/JBJS.RVW.N.00053
4. Ahn AK, Chang D, Plate AM: Triangular fibrocartilage complex tears: a review . Bull NYU Hosp Jt Dis. 2006, 64:114-8.
5. Haugstvedt JR, Langer MF, Berger RA: Distal radioulnar joint: functional anatomy, including pathomechanics. J Hand Surg Eur Vol. 2017, 42:338-45. 10.1177/1753193417693170
6. Garcia-Elias M, Domènech-Mateu JM: The articular disc of the wrist. Limits and relations . Acta Anat (Basel). 1987, 128:51-4. 10.1159/000146314
7. Benjamin M, Evans EJ, Pemberton DJ: Histological studies on the triangular fibrocartilage complex of the wrist. J Anat. 1990, 172:59-67.
8. Palmer AK, Werner FW: The triangular fibrocartilage complex of the wrist--anatomy and function . J Hand Surg Am. 1981, 6:153-62. 10.1016/s0363-5023(81)80170-0
9. Palmer AK: Triangular fibrocartilage complex lesions: a classification . J Hand Surg Am. 1989, 14:594-606. 10.1016/0363-5023(89)90174-3
10. Oneson SR, Scales LM, Timins ME, Erickson SJ, Chamoy L: MR imaging interpretation of the Palmer classification of triangular fibrocartilage complex lesions. Radiographics. 1996, 16:97-106. 10.1148/radiographics.16.1.97
11. Nakamura T, Yabe Y: Histological anatomy of the triangular fibrocartilage complex of the human wrist . Ann Anat. 2000, 182:567-72. 10.1016/S0940-9602(00)80106-5
12. Nakamura T, Yabe Y, Horiuchi Y: Functional anatomy of the triangular fibrocartilage complex . J Hand Surg Br. 1996, 21:581-6. 10.1016/s0266-7681(96)80135-5
13. Nakamura T, Takayama S, Horiuchi Y, Yabe Y: Origins and insertions of the triangular fibrocartilage complex: a histological study. J Hand Surg Br. 2001, 26:446-54. 10.1054/jhsb.2001.0562

14. Slutsky DJ: Arthroscopic evaluation of the foveal attachment of the triangular fibrocartilage . Hand Clin. 2011, 27:255-61. 10.1016/j.hcl.2011.05.003
15. Atzei A, Rizzo A, Luchetti R, Fairplay T: Arthroscopic foveal repair of triangular fibrocartilage complex peripheral lesion with distal radioulnar joint instability. Tech Hand Up Extrem Surg. 2008, 12:226-35. 10.1097/BTH.0b013e3181901b1
16. Bain GI, McGuire D, Lee YC, Eng K, Zumstein M: Anatomic foveal reconstruction of the triangular fibrocartilage complex with a tendon graft. Tech Hand Up Extrem Surg. 2014, 18:92-7. 10.1097/BTH.0000000000000044
17. Skalski MR, White EA, Patel DB, Schein AJ, RiveraMelo H, Matcuk GR Jr: The traumatized TFCC: an illustrated review of the anatomy and injury patterns of the triangular fibrocartilage complex. Curr Probl Diagn Radiol. 2016, 45:39-50. 10.1067/j.cpradiol.2015.05.004
18. Srinivasan RC, Shrouder-Henry JJ, Richard MJ, Ruch DS: Open and arthroscopic triangular fibrocartilage complex (TFCC) repair. J Am Acad Orthop Surg. 2021, 29:518-25. 10.5435/JAAOS-D-20-00998
19. von Borstel D, Wang M, Small K, Nozaki T, Yoshioka H: High-resolution 3T MR imaging of the triangular fibrocartilage complex. Magn Reson Med Sci. 2017, 16:3-15. 10.2463/mrms.rev.2016-0011
20. Atzei A: New trends in arthroscopic management of type 1-B TFCC injuries with DRUJ instability . J Hand Surg Eur Vol. 2009, 34:582-91. 10.1177/1753193409100120
21. Hippocrates: On joints. Hippocrates, Volume III: On Wounds in the Head, In the Surgery, On Fractures, On Joints, Mochlicon. Withington ET (ed): Harvard University Press, Cambridge, MA; 1928. 149.
22. Kauer JM: [The development of the triangular ligament in man] . Bull Assoc Anat (Nancy). 1975, 59:893-8.
23. Mikić ZD: Age changes in the triangular fibrocartilage of the wrist joint . J Anat. 1978, 126:367-84.
24. Kuhlmann JN, Vinh T, Guerin-Surville H: [The triangular ligament and the medial compartment of the wrist]. Bull Assoc Anat (Nancy). 1985, 69:305-14.
25. Whipple TL, Marotta JJ, Powell JH 3rd: Techniques of wrist arthroscopy. Arthroscopy. 1986, 2:244-52. 10.1016/s0749-8063(86)80079-2
26. Roth JH, Poehling GG, Whipple TL: Arthroscopic surgery of the wrist. Instr Course Lect. 1988, 37:183-94.
27. Atzei A, Luchetti R: Foveal TFCC tear classification and treatment. Hand Clin. 2011, 27:263-72. 10.1016/j.hcl.2011.05.014
28. Chen WJ: Arthroscopically assisted transosseous foveal repair of triangular fibrocartilage complex . Arthrosc Tech. 2017, 6:e57-64. 10.1016/j.eats.2016.09.004
29. Park JH, Kim D, Park JW: Arthroscopic one-tunnel transosseous foveal repair for triangular fibrocartilage complex (TFCC) peripheral tear. Arch Orthop Trauma Surg. 2018, 138:131-8. 10.1007/s00402-017-2835-3
30. Ma HH, Wang JP, Yang CY: Effectiveness of suture anchor and transosseous suture technique in arthroscopic foveal repair of the triangular fibrocartilage complex: a systematic review. J Orthop Surg Res. 2024, 19:72. 10.1186/s13018-024-04530-4
31. Uchiyama S, Nakatsuchi Y: Anatomical and radiological evaluation of the triangular fibrocartilage complex of the wrist. J Hand Surg Br. 1994, 19:319-24. 10.1016/0266-7681(94)90080-9
32. Palmer AK, Glisson RR, Werner FW: Relationship between ulnar variance and triangular fibrocartilage complex thickness. J Hand Surg Am. 1984, 9:681-2. 10.1016/s0363-5023(84)80013-1
33. af Ekenstam FW, Palmer AK, Glisson RR: The load on the radius and ulna in different positions of the wrist and forearm. A cadaver study. Acta Orthop Scand. 1984, 55:363-5. 10.3109/17453678408992375
34. Atzei A, Luchetti R, Braidotti F: Arthroscopic foveal repair of the triangular fibrocartilage complex . J Wrist Surg. 2015, 4:22-30. 10.1055/s-0035-1544226
35. Bednar MS, Arnoczky SP, Weiland AJ: The microvasculature of the triangular fibrocartilage complex: its clinical significance. J Hand Surg Am. 1991, 16:1101-5. 10.1016/s0363-5023(10)80074-7
36. Atzei A, Luchetti R, Carletti D, Marcovici LL, Cazzoletti L, Barbon S: The hook test is more accurate than the trampoline test to detect foveal tears of the triangular fibrocartilage complex of the wrist. Arthroscopy. 2021, 37:1800-7. 10.1016/j.arthro.2021.03.005
37. Trehan SK, Wall LB, Calfee RP, Shen TS, Dy CJ, Yannascoli SM, Goldfarb CA: Arthroscopic diagnosis of the triangular fibrocartilage complex foveal tear: a cadaver assessment. J Hand Surg Am. 2018, 43:680.e1-5. 10.1016/j.jhsa.2017.12.017
38. Shionova K, Nakamura R, Imaeda T, Makino N: Arthrography is superior to magnetic resonance imaging for diagnosing injuries of the triangular fibrocartilage. J Hand Surg Br. 1998, 23:402-5. 10.1016/s0266-7681(98)80068-5
39. Smith TO, Drew B, Toms AP, Jerosch-Herold C, Chojnowski AJ: Diagnostic accuracy of magnetic resonance imaging and magnetic resonance arthrography for triangular fibrocartilaginous complex injury: a systematic review and meta-analysis. J Bone Joint Surg Am. 2012, 94:824-32. 10.2106/JBJS.J.01775
40. Osterman AL: Arthroscopic debridement of triangular fibrocartilage complex tears . Arthroscopy. 1990, 6:120-4. 10.1016/0749-8063(90)90012-3
41. Saito T, Malay S, Chung KC: A systematic review of outcomes after arthroscopic débridement for triangular fibrocartilage complex tear. Plast Reconstr Surg. 2017, 140:697e-708e. 10.1097/PRS.0000000000003750
42. van der Post AS, Jens S, Daams JG, Obdeijn MC, Maas M, Oostra RJ: The triangular fibrocartilage complex in the human wrist: a scoping review toward uniform and clinically relevant terminology. Clin Anat. 2022, 35:626-48. 10.1002/ca.23880
43. Schmitt R, Kunz AS, Reidler P, Huflage H, Hesse N: Triangular fibrocartilage complex (TFCC) - anatomy, imaging, and classifications with special focus on the CUP classification. Rofo. 2025, 197:759-69. 10.1055/a-2411-8444
44. Minhas S, Kakar S, Wall LB, Goldfarb CA: Foveal triangular fibrocartilage complex tears: recognition of a combined tear pattern. J Hand Surg Am. 2023, 48:1063.e1-6. 10.1016/j.jhsa.2022.03.010
45. Iordache SD, Rowan R, Garvin GJ, Osman S, Grewal R, Faber KJ: Prevalence of triangular fibrocartilage complex abnormalities on MRI scans of asymptomatic wrists. J Hand Surg Am. 2012, 37:98-103. 10.1016/j.jhsa.2011.10.006
46. Chan JJ, Teunis T, Ring D: Prevalence of triangular fibrocartilage complex abnormalities regardless of symptoms rise with age: systematic review and pooled analysis. Clin Orthop Relat Res. 2014, 472:3987-94.

10.1007/s11999-014-3825-1

47. Zhao X, Yu A, Zhao H, Qiu Y: Diagnostic value of MRI in traumatic triangular fibrocartilage complex injuries: a retrospective study. BMC Musculoskelet Disord. 2024, 25:63. 10.1186/s12891-023-07140-z
48. Flores DV, Umpire DF, Rakhra KS, Jibri Z, Belmar GA: Distal radioulnar joint: normal anatomy, imaging of common disorders, and injury classification. Radiographics. 2023, 43:e220109. 10.1148/rg.220109
49. Kim JS, Kim KE, Lee SW, Jeon S, Yang H, Choi YR: Generalized joint laxity increases the risk of recurrence of distal radioulnar joint instability after arthroscopic foveal repair of the triangular fibrocartilage complex. Arthroscopy. 2025, 41:2309-16. 10.1016/j.arthro.2024.10.040