

DOI: 10.7759/cureus.2915

Emergency Management of Anaphylaxis: A High Fidelity Interprofessional Simulation Scenario to Foster Teamwork Among Senior Nursing, Medicine, and Pharmacy Undergraduate Students

Sandra MacDonald 1 , April Manuel 2 , Adam Dubrowski 3 , Natalie Bandrauk 4 , Rebecca Law 5 , Vernon Curran 6 , Young Wah Lee 4

1. School of Nursing, Memorial University of Newfoundland, St. John's, CAN 2. School of Nursing, Memorial University of Newfoundland, Conception Bay South, CAN 3. Emergency Medicine, Memorial University of Newfoundland, St. John's, CAN 4. Faculty of Medicine, Memorial University of Newfoundland, St. John's, CAN 5. School of Pharmacy, Memorial University of Newfoundland, St. John's, CAN 6. Faculty of Medicine, Memorial University of Newfoundland

☑ Corresponding author: Sandra MacDonald, smacdon@mun.ca
Disclosures can be found in Additional Information at the end of the article

Abstract

Nursing, medicine, and pharmacy students have limited opportunities during their undergraduate programs to learn and practice together as an interprofessional team. This has prompted faculty at Memorial University of Newfoundland to explore the use of high fidelity simulated interprofessional education (HF-IPE) to help nursing, medicine, and pharmacy students learn about their roles, develop communication and collaboration skills, and foster teamwork. Research has shown that high fidelity simulated education can promote critical thinking, engage learners, improve confidence, and enhance psychomotor skills; however, there is limited data on the impact of HF-IPE on fostering teamwork. This technical report describes one HF-IPE scenario designed to foster teamwork among senior undergraduate nursing, medicine, and pharmacy students. The scenario is designed to promote an understanding of the roles of nursing, medical, and pharmaceutical professionals in an interprofessional team during the emergency management of an adult patient experiencing acute anaphylaxis. Teamwork and communication skills are emphasized, and students are provided with the opportunity to communicate and collaborate within an interprofessional healthcare team.

Received 05/29/2018 Review began 06/04/2018 Review ended 06/30/2018 Published 07/03/2018

© Copyright 2018

MacDonald et al. This is an open access article distributed under the terms of the Creative Commons

Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Categories: Emergency Medicine, Medical Simulation

Keywords: high-fidelity simulation, interprofessional education, anaphylaxis, nursing, pharmacy, medicine, undergraduate education

Introduction

Teamwork which involves learning with, from, and about one another is an important and requisite skill that must be taught, supported, and nurtured in undergraduate health sciences programs [1-4]. Currently, nursing, medicine, and pharmacy students have few opportunities to practice interprofessional teamwork during their undergraduate education, and they often enter the workforce with little or no experience in working with interprofessional healthcare teams [5-6]. This has been further complicated by a lack of opportunities to practice teamwork

How to cite this article

Macdonald S, Manuel A, Dubrowski A, et al. (July 03, 2018) Emergency Management of Anaphylaxis: A High Fidelity Interprofessional Simulation Scenario to Foster Teamwork Among Senior Nursing, Medicine, and Pharmacy Undergraduate Students. Cureus 10(7): e2915. DOI 10.7759/cureus.2915

behaviors in the clinical setting. These concerns have prompted educators and researchers to search for appropriate and effective teaching and learning strategies to prepare senior students for real-life situations as members of the healthcare team [7].

One effective teaching and learning strategy known to foster teamwork among undergraduate health sciences students is simulated interprofessional education [8-9]. In particular, high fidelity simulated interprofessional education (HF-IPE) has been shown to effectively promote skill acquisition, enhance clinical judgment, and teach students about complex clinical situations [10-14]. HF-IPE can provide an invaluable safety net for learning, allowing students to acquire and develop critical-thinking and decision-making skills without exposing patients or students to unnecessary risk [15-18].

The HF-IPE scenario discussed in this report is designed to create a safe learning environment for undergraduate students to learn with and from one another while managing the care of an adult patient experiencing acute anaphylaxis. This HF-IPE scenario was developed using the Jeffries simulation framework for designing, implementing, and evaluating simulation education. The Jeffries framework includes five major components: the simulation design, learner outcomes, faculty role, student characteristics, and educational practices [19]. The simulation design for this scenario is high fidelity, with learner outcomes including enhanced knowledge of the roles of nursing, medical, and pharmaceutical professionals on the team, improved attitudes towards teamwork, and the demonstration of teamwork behaviors while caring for a patient experiencing anaphylaxis. The learning experience is student-centered, while the faculty play the role of facilitator and observer. The educational practices include building on students' previous knowledge and skills related to teamwork to create new knowledge, behaviors, and skills.

The World Allergy Organization Guidelines for the assessment and management of anaphylaxis are used to guide the simulation, thus creating an awareness of best practice protocols for managing acute anaphylaxis, as well as an opportunity to foster teamwork among undergraduate nursing, medicine, and pharmacy students [20].

Technical Report

There are three healthcare professional education faculties and schools at Memorial University of Newfoundland: nursing, medicine, and pharmacy, all of which are dedicated to the development of undergraduate and postgraduate students. However, currently, there are few opportunities for these groups of undergraduate students to interact face to face as required during common interprofessional activities. The purpose of implementing the HF-IPE scenario described in this technical report is to provide undergraduate nursing, medical, and pharmacy students with an opportunity to build on previous knowledge and skills by participating as effective members of an interprofessional healthcare team during the management of an adult patient experiencing acute anaphylaxis. This scenario is designed to be implemented within any senior level clinical course. The purpose, learning objectives, scope of practice and debriefing points for the scenario are outlined in Figure 1.

Purpose	To participate as an effective member of an interprofessional health care		
r ui pose	team during the emergency management of an adult patient experiencing		
7	acute anaphylaxis.		
Learning	Explain the role of nursing, medicine and pharmacy during the emergency		
Objectives	management of an adult patient experiencing anaphylaxis.		
	Communicate effectively with the interprofessional health care team.		
	Collaborate with the team to manage an adult patient experiencing anaphylaxis.		
	Implement the professional guidelines for the emergency management of		
	an adult patient experiencing anaphylaxis including recognizing the		
	clinical criteria, and acute management.		
Scope of	Nursing; health and physical assessment, oxygen therapy, intramuscular		
Practice	injection of epinephrine, inserting an intravenous catheter, interpreting		
	laboratory and diagnostic tests (e.g. electrocardiogram, arterial blood		
	gases), calculating fluid challenges and basic cardiac life support (BCLS).		
	Medicine; history taking and physical assessment; medical treatment and		
	management of anaphylaxis, calculating fluid challenges and ordering		
	medications and interpreting and diagnosing findings from diagnostic and		
	laboratory tests (e.g. Computed Tomography, chest x-ray) and BCLS.		
	Pharmacy; patient medication assessment, knowledge of potential drug		
	interactions and side effects (e.g. beta-blockers and Epinephrine), drug		
	dosage calculations, recommend and dispense medications as ordered.		
Debriefing	What was your role on the team?		
Points	Do you think the team provided good care? Why or why not?		
	Can you give examples of when the team communicated effectively?		
	Are there any examples of the team coordinating care?		
	Do you think the team collaborated on decisions?		
	Did the team follow the professional guidelines?		

FIGURE 1: Anaphylaxis Teaching Plan

Inputs

- 1. Senior undergraduate nursing, medicine, and pharmacy students
- 2. Nursing, medicine, and pharmacy interprofessional educators
- 3. Adult female/male high fidelity human patient simulator (HPS) programmed to reflect the five stages of the simulation: (1) baseline assessment, (2) early reaction, (3) anaphylaxis, (4) recovery and (5) resolution
- 4. Mock chart: a paper-based chart containing the patient's history and physical exam notes, progress notes, nursing kardex, medication orders, physician orders, vital signs record, medication administration record, laboratory and diagnostics tests, and anaphylaxis protocol
- 5. Equipment: cardiac monitor, pulse oximeter, Venturi mask, intravenous (IV) access, nebulizer and face mask, intravenous fluids, IV set-up tray, mannequin make up for urticaria on abdomen

Processes

The story board developed to guide the implementation of the simulation is divided into five

phases: (1) baseline assessment, (2) early hypotensive reaction, (3) anaphylaxis, (4) recovery, and (5) resolution (Figure 2). Each phase of the simulation includes a description of the situation, assessment data, prompts, and possible actions. Further information for programming the human patient simulator is also provided in the 'Story Board for Programming the Human Patient Simulator' including vital signs and cardiac and respiratory status (Figure 3).

Baseline Assessment Team Meeting

Mrs. Jones is a 50-year-old woman admitted for acute right-sided pleuritic chest pain. She just arrived home from an overnight international flight. She has a history of hypertension, angina and diabetes (Type II). Medications are Ramipril, Metoprolol, Metformin Rosuvastatin. She is allergic to shellfish and wears an allergy band. Received Intravenous (IV) fluids overnight. CT-PA finished 10 minutes ago. Wheezing since the CT and now has pruritus and urticarial lesions on her abdomen. She is receiving O2 28% via Venturi Mask (VM) but the mask is not positioned properly. Oxygen saturation (SaO₂) is 94 % but she is less alert than on admission. Heart Rate (HR) 90, Blood Pressure (BP) 110/70, Respiratory Rate (RR) 32, Oxygen Saturation (SaO₂) 94%, Temperature (T) 37.7

(SaC₂) 94%, 1 emperature (1) 37.7

Actions
Team Meeting
Chart Review
Review Anaphylaxis Protocol

Review Management Reposition VM

Recovery

Three to five minutes after Epinepherine, BP remains low, tongue swollen and wheezing continues. Team discusses possible drug interactions. Vital signs slowly improving: HR 150, B/P 110/80, RR 32, SaO₂92%

"My tongue still feels a little swollen, but I am not so itchy."

"I still feel a little short of breath."

Actions
Continue fluid bolus
2nd dose Epinephrine
Review ABGs if ordered

Early Hypotensive Reaction

Team discusses a possible allergic reaction to CT dye and the proposed medical treatment. Mrs. Jones is initially able to answer simple questions but becomes more confused. Face flushed, hives and urticarial lesions present on trunk. Complaining of feeling "itchy". Mrs. Jones has shallow rapid breathing with audible, expiratory wheezes. Coarse rhonchi are heard in both lower lobes. Electrocardiogram (ECG) shows sinus tachycardia.

HR 110, BP 90/70, RR 30, SaO₂ 92% *Actions*

Recognize Action.
Assess

Allergic

Reaction

Place in Trendelenburg position Possible Actions

Order Arterial Blood Gases (ABGs) Establish a 2nd intravenous access Administer 1-2 liters of 0.9% (isotonic) saline rapidly with 5-10 mL/kg in the first 5-10 minutes

Initiate

Administer Epinephrine

Administer Epinephrine IM HR 150, BP 80/60, RR 26,

Sa O₂ 90%

"My lips feel swollen."
"I feel itchy all over."

Anaphylaxis

Actions
Discuss plan
Review ABGs (if ordered)
Increase oxygen via facemask
Administer 1st dose Epinephrine
Possible Actions
Diphenhydramine
Solu-Cortef



Resolution

Mrs. Jones showing clinical improvement and the urticarial lesions are resolving.

"It is not as itchy and my tongue feels better"

BP 120/80, HR 95, RR 24, SaO₂ 94%

FIGURE 2: Story Board for Acute Anaphylaxis

VS - vital signs; HR - heart rate; BP - blood pressure; RR - respiratory rate; T - temperature; CT - computed tomography; ECG - electrocardiogram; SaO2 - oxygen saturation; ABG - arterial blood gas; IM - intramuscular; VM - Venturi mask, and IV - intravenous.

Vital Signs	Prompts	Interventions	Possible Outcomes		
	Rasalina Assassman	t and Team Meeting (10 – 15 minu			
HR 90 BP 110/70 RR 32 T 37.7 Sa O ₂ 94% O ₂ 28% via VM	Point out red patches on abdomen Team should discuss possible allergic reaction	Team Meeting (10 = 13 mina Team Meeting Chart Review Assess circulation, airway, breathing, mental status, and skin Review emergency protocol	Risk assessed Plan developed Protocol reviewed Medications reviewed /ordered Recognized beta-		
Wheezing heard in lower lobes	to contrast dye	Reposition VM Review medication interactions	blockers can effectEpinephrine		
Early Hypotensive Reaction (10 – 15 minutes)					
HR 110 BP 90/70 RR 30 Sa O ₂ 92% Expiratory wheezes, shallow rapid breathing, Coarse rhonchi	"My stomach feels itch like my allergy." Face flushed, hives and urticarial lesions present on trunk.	Assess patient 2 nd IV and ABGs (if ordered), Fluid bolus & Epinepherine Trendelenburg position Calculate & administer 1-2 liters of 0.9% saline rapidly with 5-10 mL/kg in the first 5-10 minutes	Decide on care 2 nd IV & ABGs (if ordered) IV fluid bolus administered Called for help		
	Anaph	nylaxis (10 - 15 minutes)			
HR 150, BP 80/60, RR 30 Sa O ₂ 90% 3-5 minutes post Epinephrine Sa O ₂ 94%	"My lips feel swollen." "I feel itchy all over"	Increase O ₂ 8 L/min Administer IM Epinephrine in the mid-anterolateral aspect of the thigh, 0.01 mg/kg to maximum does of 0.5 mg IM Diphenhydramine & Solu- Cortef (if ordered)	SaO ₂ increasing BP rising 1 st dose Epinepehrine IV fluid bolus Review ABGs Oxygen increased		
Recovery (10 – 15 minutes)					
HR 150 BP 110/80, RR 32 Sa O ₂ 92% Wheezing and rhonchi reduced	"My tongue still feels a little swollen, but I am not so itchy."	Assess patient Review IV fluids 2 nd dose Epinephrine	Continue fluid bolus 2 nd Epipherine B/P slowly rising Allergic reaction resolving		
		esolution (5 minutes)			
BP 120/80 HR 95 RR 24 Sa O ₂ 94%	"I am not as itchy and my tongue feels better."	Monitor patient's blood pressure, heart rate and function, respiratory status and oxygenation. Must adminster 2 nd dose Epinepherine to resolve. ABGs reviewed if ordered			

FIGURE 3: Story Board for Programming Human Patient Simulator

VS - vital signs; HR - heart rate; BP - blood pressure; RR - respiratory rate; T - temperature; CT - computed tomography; ECG - electrocardiogram; SaO2 - oxygen saturation; ABG - arterial blood gas; IM - intramuscular; VM - Venturi mask, and IV - intravenous.

Discussion

Teamwork is an important and requisite skill that must be taught, supported, and nurtured in nursing, medicine, and pharmacy undergraduate education programs. Although there are three health professional education faculties and schools at Memorial University of Newfoundland, nursing, medicine, and pharmacy, currently there are few opportunities for these groups of undergraduate students to interact face-to-face during common interprofessional activities. This is not unique to Memorial, and similar early introductions to teamwork and communication at the undergraduate learners' level are absent in other universities. This has

led faculty at Memorial to develop a HF-IPE scenario to foster teamwork; however, barriers to implementing this scenario exist, including common challenges associated with implementing IPE in the academic setting: difficulties in scheduling and in matching students' scope of practice.

The professional nursing, medicine, and pharmacy courses have rigid and busy schedules with little or no time for interprofessional activities, unless they can be scheduled within a course. In addition, the scope of practice for each of the professions varies depending on the level of the student, e.g., junior or senior. This HF-IPE scenario is designed for senior-level students, but when there are different levels of students participating, certain modifications may be needed. For example, junior medical students may be aware of the practice protocols for anaphylaxis, but they may have little or no experience managing the treatment of acute anaphylaxis; in contrast, senior medical students would be more likely to have both, knowledge of protocols as well as experience in managing a deteriorating patient in the clinical setting. This varying scope of practice could affect the functioning of the team and have an impact on their learning about their roles.

In order to offset some of these limitations, this simulation scenario is developed for seniorlevel students and focuses on fostering teamwork in addition to developing proficiency in managing anaphylaxis in a deteriorating patient. Prior to participating in the simulation, students are given time to review the anaphylaxis protocols and mock chart and ask questions about their role in the team. The focus of this simulation is teamwork, which is also emphasized in the discussion questions that guide the debriefing session: e.g., do you think the team communicated effectively? In this way, this scenario is designed to help educators foster teamwork among senior students and promote an understanding of their role in the interprofessional team during the management of an adult patient experiencing acute anaphylaxis. Teamwork and communication skills are emphasized in this scenario, by providing students with an opportunity to communicate and collaborate face-to-face with an interprofessional healthcare team. Best practice guidelines for the emergency management of an adult patient experiencing acute anaphylaxis are used to guide the simulation, thus creating an awareness of best practice protocols for managing acute anaphylaxis, as well as creating an opportunity for undergraduate nursing, medicine, and pharmacy undergraduate students to learn with, from, and about one other.

Conclusions

The purpose of this technical report is to provide educators who use HF-IPE with a detailed teaching plan and story boards for implementing an HF-IPE scenario to foster teamwork among senior nursing, medicine, and pharmacy undergraduate students. This report is designed to provide the reader with the context, required resources, scenario, learning objectives, and possible outcomes for a high fidelity simulation designed to care for an adult patient experiencing acute anaphylaxis. Teamwork and communication skills are emphasized while students are provided with the opportunity to learn with, from, and about the nursing, medical, and pharmaceutical members of the interprofessional healthcare team. Best practice guidelines for the emergency management of an adult patient experiencing acute anaphylaxis are used to guide teamwork behaviors during the simulation.

Additional Information

Disclosures

Human subjects: All authors have confirmed that this study did not involve human participants or tissue. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform

disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- 1. Kvan T: Evaluating learning environments for interprofessional care. J Interprof Care. 2013, 27:33-36. 10.3109/13561820.2013.791673
- Robertson J, Bandali K: Bridging the gap: enhancing interprofessional education using simulation. J Interprof Care. 2008, 22:499-508. 10.1080/13561820802303656
- Thistlethwaite J, Kumar K, Moran M, Saunders R, Carr S: An exploratory review of prequalification interprofessional education evaluations. J Interprof Care. 2015, 29:292-297. 10.3109/13561820.2014.985292
- 4. Welllmon R, Lefebvre KM, Ferry D: Effects of high-fidelity simulation on physical therapy and nursing students' attitudes toward interprofessional learning and collaboration. J Nurs Educ. 2017, 56:456-465. 10.3928/01484834-20170712-03
- Hayden J, Keegan M, Kardong-Edgren S, Smiley RA: Reliability and validity testing of the Creighton competency evaluation instrument for use in the NCSBN National Simulation Study. Nurs Educ Perspect. 2014, 35:244-252. 10.5480/13-1130.1
- Murphy J, Nimmagadda J: Partnering to provide simulated learning to address interprofessional education collaborative core competencies. J Interprof Care. 2015, 29:258-259. 10.3109/13561820.2014.942779
- 7. Chen S, Huang T, LiaoI I, Liu C: Development and validation of the simulation learning effectiveness inventory. J Adv Nurs. 2015, 71:2444-2453. 10.1111/jan.12707
- Lin YC, Chan TF, Lai CS, et al.: The impact of an interprofessional problem-based learning curriculum of clinical ethics on medical and nursing students' attitudes and ability of interprofessional collaboration: a pilot study. Kaohsiung J Med Sci. 2013, 29:505-511. 10.1016/j.kjms.2013.05.006
- 9. Tullmann DF, Shilling AM, Goeke LH, Wright EB, Littlewood KE: Recreating simulation scenarios for interprofessional education: an example of educational interprofessional practice. J Interprof Care. 2013, 27:426-428. 10.3109/13561820.2013.790880
- Paige JT, Garbee DD, Kozmenko V, et al.: Getting a head start: high-fidelity, simulation-based operating room team training of interprofessional students. J Am Coll Surg. 2014, 218:140-149. 10.1016/j.jamcollsurg.2013.09.006
- 11. Alfes CM, Steiner SL, Manacci C: Critical care transport training: new strides in simulating the austere environment. Air Med J. 2015, 34:186-187.
- 12. Aqel AA, Muayyad MA: High-fidelity simulation effects on CPR knowledge, skills, acquisition, and retention in nursing students. Worldviews Evid Based Nurs. 2014, 11:394-400. 10.1111/wvn.12063
- 13. Braude P, Reedy G, Dasgupta D, Dimmock V, Jaye P, Birns J: Evaluation of a simulation training program for geriatric medicine. Age Ageing. 2015, 44:677-682. 10.1093/ageing/afv049
- 14. Cheng A, Lockey A, Bhanji F, Lin Y, Hunt EA, Lang E: The use of high-fidelity manikins for advanced life support training: A systematic review and meta-analysis. Resuscitation. 2015, 93:142-149. 10.1016/j.resuscitation.2015.04.004
- 15. Donoghue A, Nishisaki A: High-fidelity in simulation education: only a part of the answer . Resuscitation. 2015, 93:A3-4. 10.1016/j.resuscitation.2015.05.004
- 16. Wang CL, Chnnugounder S, Hippe D, et al.: Comparative effectiveness of hands-on versus computer simulation-based training for contrast media reactions and teamwork skills. J Am Coll Radiol. 2017, 14:103-110. 10.1016/j.jacr.2016.07.013
- 17. Cooper J, Taqueti V: A brief history of the development of mannequin simulators for clinical education and training. BMJ Qual Saf. 2005, 13:i11-i18. 10.1136/qshc.2004.009886
- 18. Leonard B, Shuhaibar E, Chen R: Nursing student perceptions of intraprofessional team education using high-fidelity simulation. J Nurs Educ Pract. 2010, 49:628-631. 10.3928/01484834-20100730-06

- 19. Adamson K: A systematic review of the literature related to the NLN/Jeffries Simulation Framework. Nurs Educ Perspect. 2015, 36:281-291. 10.5480/15-1655
- Simons F, Ardusso LR, Bilò MB, et al.: 2012 Update: World Allergy Organization guidelines for the assessment and management of anaphylaxis. Curr Opin Allergy Clin Immunol. 2012, 12:389-399. 10.1097/ACI.0b013e328355b7e4